

Optimal control of the convection-diffusion equation using stabilized finite element methods

Roland Becker · Boris Vexler

Received: 6 March 2006 / Published online: 3 March 2007
© Springer-Verlag 2007

Abstract In this paper we analyze the discretization of optimal control problems governed by convection-diffusion equations which are subject to pointwise control constraints. We present a stabilization scheme which leads to improved approximate solutions even on coarse meshes in the convection dominated case. Moreover, the in general different approaches “optimize-then-discretize” and “discretize-then-optimize” coincide for the proposed discretization scheme. This allows for a symmetric optimality system at the discrete level and optimal order of convergence.

Mathematics Subject Classification (2000) Optimal control · Stabilized finite elements · Error estimates · Pointwise inequality constraints

1 Introduction

This paper is concerned with the discretization of optimal control problems using stabilized finite element methods. We consider the following optimal control problem governed by a linear convection diffusion reaction equations on a

R. Becker
Laboratoire de Mathématiques Appliquées, Université de Pau et des Pays de l'Adour,
BP 1155, 64013 PAU Cedex, France
e-mail: roland.becker@univ-pau.fr

B. Vexler (✉)
Johann Radon Institute for Computational and Applied Mathematics (RICAM),
Austrian Academy of Sciences, Altenberger Straße 69, 4040 Linz, Austria
e-mail: boris.vexler@oeaw.ac.at

polygonal domain $\Omega \subset \mathbb{R}^d$, $d = 2, 3$:

$$\text{Minimize } J(q, u) = \frac{1}{2} \|u - u_d\|_{L^2(\Omega)}^2 + \frac{\alpha}{2} \|q\|_{L^2(\Omega)}^2, \quad u \in V, q \in Q_{ad} \quad (1)$$

subject to

$$-\varepsilon \Delta u + \beta \cdot \nabla u + \sigma u = f + q \quad \text{in } \Omega, \quad (2)$$

$$u = 0 \quad \text{on } \partial\Omega. \quad (3)$$

The state variable u is searched for in the space $V = H_0^1(\Omega)$ and the admissible set $Q_{ad} \subset Q = L^2(\Omega)$ is given by pointwise box constraints, i.e.,:

$$Q_{ad} = \{q \in L^2(\Omega) : a \leq q \leq b \quad \text{a.e. in } \Omega\},$$

where $a, b \in \mathbb{R} \cup \{\pm\infty\}$, $a < b$.

For the data of the problem we assume: $f, u_d \in L^\infty(\Omega)$, $\sigma \in L^\infty(\Omega)$ with $\sigma \geq \sigma_0 > 0$ and $\alpha > 0$. To simplify the notation we set throughout $\sigma = 1$. Further we assume β to be a constant vector of length $\|\beta\|$.

It is well known that for convection dominated problems standard finite element discretizations applied to the Eqs. (2) and (3) lead to strongly oscillatory solutions unless the mesh size h is sufficiently small with respect to the ratio between ε and $\|\beta\|$. Several methods are known to improve the approximation properties of the pure Galerkin discretization and to reduce the oscillatory behavior, see e.g., [6, 14, 16, 21, 22].

In [9] the authors apply the SUPG method (streamline upwind Petrov Galerkin method, see e.g., [16]) to the optimal control problem (1)–(3). They discuss two different approaches to the discretization of the optimal control problem: “optimize-then-discretize” and “discretize-then-optimize”. In the “optimize-then-discretize” approach first the necessary optimality conditions are established on the continuous level consisting of the state, adjoint and the optimality equations, and then these equations are discretized using a stabilized finite element scheme, e.g., SUPG. In the “discretize-then-optimize” approach the state equation is discretized and then the optimality system for the finite dimensional optimization problem is derived. It is well known that these two approaches lead to the same discretization scheme provided a pure Galerkin discretization is used. However, in the presence of stabilization terms these approaches may differ. In [9] it is shown by numerical computations that for the SUPG discretization the “optimize-then-discretize” approach leads to better asymptotic convergence properties. However, the “discretize-then-optimize” approach has the important advantage of consistency of the state and the adjoint equations on the discrete level which is reflected in the fact that the corresponding optimality system is symmetric. In [10] the authors propose a method with similar behavior for a SUPG-like discretization, based on a non-symmetric modification of the Lagrangian.

In this paper we analyze a stabilization method, which leads to symmetric optimality systems and has optimal order of convergence. For the resulting discretization scheme the approaches “optimize-then-discretize” and “discretize-then-optimize” coincide. The presented method uses standard finite element discretization with stabilization based on local projections (called LPS-method), see [14] for convection diffusion reaction equations and for the Stokes-, Oseen- and Navier–Stokes equations see [2, 4, 5]. The control space is likewise discretized by first-order finite elements.

The main contribution of this paper is the a priori error analysis of the discretization of the optimal control problem (1)–(3) by stabilized finite elements. For both the unconstrained case ($Q_{ad} = Q$) and the constrained case ($Q_{ad} \neq Q$) we obtain the estimate of order $\mathcal{O}(h^{3/2})$ for the L^2 -error in the control, state and the adjoint state.

For the a priori error analysis of pure Galerkin finite element discretizations for optimal control problems with pointwise inequality constraints, we refer e.g., to [1, 11, 15, 20]. In [24] discretization by stabilized finite element for an optimal control problem governed by Stokes equations is analyzed.

Our results are optimal for the following two reasons: first, it is well known that stabilized finite elements leads to optimal order of convergence of $\mathcal{O}(h^{3/2})$ in $L^2(\Omega)$ -norm for the convection diffusion reaction equation (2) and (3) on general quasi-uniform meshes, see e.g., [26]. Second, the presence of control constraints leads to the fact that the optimal control \bar{q} is in general not in $H^2(\Omega)$ and only $\mathcal{O}(h^{3/2})$ convergence can be expected for the piecewise (bi)linear discretization of the control space, see e.g., [23].

The outline of the paper is as follows: in the next section we discuss the precise formulation of the optimal control problem and the optimality conditions. In Sect. 3 we describe the discretization of (1)–(3) and formulate our main results. In Sect. 4 we prove an error estimate for the optimal control problem without control constraints. The extension to the case with pointwise control constraints is presented in Sect. 5. In Sect. 6 we present numerical results.

2 Optimal control problem

In this section we discuss the optimality conditions for the optimal control problem (1)–(3).

A weak solution $u \in V = H_0^1(\Omega)$ of the state equation (2) and (3) is determined by

$$a(u, v) = (f + q, v) \quad \forall v \in V$$

using the bilinear form $a: V \times V \rightarrow \mathbb{R}$ given by:

$$a(u, v) = \varepsilon(\nabla u, \nabla v) + (\beta \cdot \nabla u, v) + (u, v).$$

Throughout the paper we make the following assumption on the domain $\Omega \subset \mathbb{R}^d$:

Assumption 1 There exists $s \in \mathbb{R}, s > d$ such that the weak solution $v \in H_0^1(\Omega)$ of

$$-\varepsilon \Delta v + \beta \cdot \nabla v + v = g \quad \text{in } \Omega, \tag{4}$$

$$v = 0 \quad \text{on } \partial\Omega \tag{5}$$

satisfies the a priori error estimates:

$$\|v\|_{W^{2,p}(\Omega)} \leq c \|g\|_{L^p(\Omega)} \quad \text{for all } g \in L^p(\Omega), \text{ for all } p, 2 \leq p \leq s.$$

Remark 1 For $d = 2$ a sufficient condition for the above assumption is the convexity of the domain Ω , see e.g., [13]. For $d = 3$ one needs an additional angle condition, see e.g., [17].

Theorem 1 Under the above assumption the optimal control problem (1)–(3) admits a unique solution $(\bar{q}, \bar{u}) \in (Q_{ad} \cap W^{1,\infty}(\Omega)) \times (H_0^1(\Omega) \cap W^{2,s}(\Omega))$.

The proof of this theorem follows standard arguments, see e.g., [12, 19, 25].

Remark 2 In the case $Q = Q_{ad}$, i.e., in the absence of control constraints, the optimal control \bar{q} has better regularity properties, i.e., $\bar{q} \in W^{2,s}(\Omega)$.

We denote by $S: Q \rightarrow H_0^1(\Omega) \cap H^2(\Omega)$ the solution operator of the state equations (2) and (3) and introduce the reduced cost functional $j: Q \rightarrow \mathbb{R}$ by:

$$j(q) = J(q, Sq).$$

This allows us to eliminate the state equation and to reformulate the optimization problem as:

$$\text{Minimize } j(q), \quad q \in Q_{ad}.$$

The reduced cost functional j is continuously differentiable and its derivatives are given in the following lemma:

Lemma 1 *There holds:*

$$j'(q)(\delta q) = (z, \delta q) + \alpha(q, \delta q),$$

where $z \in H_0^1(\Omega) \cap W^{2,s}(\Omega)$ is the solution of the adjoint equation:

$$-\varepsilon \Delta z - \beta \cdot \nabla z + z = u - u_d \quad \text{in } \Omega, \tag{6}$$

$$z = 0 \quad \text{on } \partial\Omega, \tag{7}$$

where $u = Sq$ is the associated state to q ; z is called the adjoint state.

Proof Using standard arguments, see e.g., [25], we obtain the above formula for the derivatives of j with $z \in H_0^1(\Omega)$. Due to the fact that $u \in H^2(\Omega) \hookrightarrow L^\infty(\Omega)$ and consequently $u - u_d \in L^\infty(\Omega)$, Assumption 1 implies that $z \in W^{2,s}(\Omega)$.

Due to the fact that Q_{ad} is convex the necessary and sufficient optimality condition for this problem can be formulated as follows:

$$j'(\bar{q})(\delta q - \bar{q}) \geq 0 \quad \forall \delta q \in Q_{ad}. \tag{8}$$

Remark 3 Using Lemma 1 condition (8) can be rewritten as:

$$(\bar{z} + \alpha \bar{q}, \delta q - \bar{q}) \geq 0 \quad \forall \delta q \in Q_{ad},$$

where \bar{z} is the associated adjoint state to \bar{q} . This condition together with the state equation (2) and (3) and the adjoint equation (6) and (7) builds the optimality system for optimal control problem (1)–(3).

In the sequel we will need the gradient $\nabla j(q) \in L^2(\Omega)$ of the reduced cost functional j , which is given as the Riesz representative of $j'(q)(\cdot)$ by:

$$\nabla j(q) = z + \alpha q.$$

Obviously, there holds $\nabla j(\bar{q}) \in W^{1,\infty}(\Omega)$.

Since j is quadratic, the second derivative $j''(q)(\delta q, \tau q)$ does not depend on q and there holds:

$$j''(q)(\delta q, \delta q) \geq \alpha \|\delta q\|_{L^2(\Omega)}^2 \quad \forall \delta q \in Q. \tag{9}$$

3 Discretization and main results

In this section we present the discretization of the optimal control problem (1)–(3) and formulate our main result concerning the asymptotic convergence properties.

For the discretization of the optimal control problem (1)–(3) we consider a family $\{\mathcal{T}_h\}_{h>0}$ of two- or three dimensional meshes consisting of (open) cells K which are either triangles, tetrahedra, quadrilaterals, or hexahedra and constitute a non-overlapping covering of the computational domain Ω . The mesh parameter h is defined as a cell-wise constant function by setting $h_K = h_K$ and h_K is the diameter of K . We use the symbol h also for the maximal cell size, i.e.,

$$h = \max_{K \in \mathcal{T}_h} h_K. \tag{10}$$

The family of meshes $\{\mathcal{T}_h\}_{h>0}$ is assumed to fulfill the standard conditions of quasi-uniformity and shape-regularity, see e.g., [8]. In addition, we require that

the mesh \mathcal{T}_h is organized in a patch-wise manner. This means that it results from a coarser regular mesh \mathcal{T}_{2h} by an uniform refinement. By a patch P of elements we denote a group of cells in \mathcal{T}_h which results from a common coarser cell in \mathcal{T}_{2h} .

For each node x of the mesh \mathcal{T}_h we denote by $\mathcal{N}_h(x) \subset \Omega$ the ‘‘neighborhood’’ of x , which is the union of all cells K with $x \in \partial K$. For quasi-uniform, shape-regular meshes there exists a constant $c_{\mathcal{N}}$ such that

$$\text{diam}(\mathcal{N}_h(x)) \leq c_{\mathcal{N}}h. \tag{11}$$

On the mesh \mathcal{T}_h we define finite element spaces $Q_h \subset H^1(\Omega)$ and $V_h \subset H_0^1(\Omega)$ consisting of linear or bilinear shape functions, see e.g., [16]. In addition we define the space of cell-wise constant functions on patches V_{2h}^{disc} .

In the sequel, we will need an L^2 -orthogonal projection operator $\pi_h: L^2(\Omega) \rightarrow V_{2h}^{\text{disc}}$. This operator has the following approximation and stability properties for $P \in \mathcal{T}_{2h}$:

$$\|\pi_h v\|_{L^2(P)} \leq c\|v\|_{L^2(P)} \quad \forall v \in L^2(P), \tag{12}$$

$$\|v - \pi_h v\|_{L^2(P)} \leq ch\|\nabla v\|_{L^2(P)} \quad \forall v \in H^1(P). \tag{13}$$

For a positive stabilization parameter δ we introduce a stabilization (bilinear, symmetric) form $s_h^\delta: V_h \times V_h \rightarrow \mathbb{R}$ by:

$$s_h^\delta(u_h, v_h) = \delta(\beta \cdot \nabla u_h - \pi_h(\beta \cdot \nabla u_h), \beta \cdot \nabla v_h - \pi_h(\beta \cdot \nabla v_h)),$$

and the discrete semilinear form $a_h: V_h \times V_h \rightarrow \mathbb{R}$ by:

$$a_h(u_h, v_h) = a(u_h, v_h) + s_h^\delta(u_h, v_h). \tag{14}$$

Then, the discretized optimal control problem is formulated as follows:

$$\text{Minimize } J(q_h, u_h), \quad q \in Q_{ad,h} = Q_{ad} \cap Q_h, \quad u_h \in V_h \tag{15}$$

subject to

$$a_h(u_h, v_h) = (f + q_h, v_h) \quad \forall v_h \in V_h. \tag{16}$$

This problem possesses a unique solution $(\bar{q}_h, \bar{u}_h) \in Q_{ad,h} \times V_h$. Similarly to the continuous case we introduce a discrete solution operator $S_h: Q \rightarrow V_h$ defined by:

$$a_h(S_h q, v_h) = (f + q, v_h) \quad \forall v_h \in V_h$$

and the discrete reduced cost functional

$$j_h(q) = J(q, S_h q).$$

The optimality condition for the discretized problem reads:

$$j'_h(\bar{q}_h)(\delta q_h - \bar{q}_h) \geq 0 \quad \forall \delta q_h \in Q_{ad,h}. \tag{17}$$

The derivatives of j_h are given in the following lemma:

Lemma 2 *There holds:*

$$j'_h(q)(\delta q) = (z_h, \delta q) + \alpha(q, \delta q),$$

where $z_h \in V_h$ is the solution of the adjoint equation:

$$a_h(v_h, z_h) = (u_h - u_d, v_h) \quad \forall v_h \in V_h. \tag{18}$$

where $u_h = S_h q$ is the associated discrete state to q . The solution z_h is the associated discrete adjoint state.

The proof follows the same lines as \hat{A} in the continuous case.

Remark 4 We note that we obtain the same discrete adjoint equation as (18) if we directly discretize the continuous adjoint equation by the same (stabilized) discretization scheme. This property relies on the symmetry of the stabilization term and leads to the fact that the approaches “optimize-then-discretize” and “discretize-then-optimize” coincide in our case.

We note that as in the continuous case the second derivative $j''_h(q)$ does depend on q and we have similarly to (9):

$$j''_h(q)(\delta q, \delta q) \geq \alpha \|\delta q\|_{L^2(\Omega)}^2 \quad \forall \delta q \in Q. \tag{19}$$

In the following theorem we formulate our main result for the optimal control problem without control constraints.

Theorem 2 *Let (\bar{q}, \bar{u}) be the solution of the optimal control problem (1)–(3) with $Q_{ad} = Q$ and (\bar{q}_h, \bar{u}_h) be the solution of the discretized problem (15) and (16) with $Q_{ad,h} = Q_h$, then the following estimate holds:*

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \tau(\delta) h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^2 \|\bar{q}\|_{H^2(\Omega)},$$

where

$$\tau(\delta) = (\delta^{1/2} \|\beta\| + \min(\delta^{-1/2}, \|\beta\| \varepsilon^{-1/2}) h + h + \varepsilon^{1/2}). \tag{20}$$

The proof of this theorem will be given in the next section. As a direct consequence of this theorem we give a rule for choosing the stabilization parameter δ . A similar rule is known for the SUPG or LPS discretization of convection-diffusion-reaction equations. The stabilization parameter is chosen in dependence of the Peclet number:

$$Pe = \frac{h \|\beta\|}{\varepsilon}$$

by

$$\delta = \begin{cases} 0, & \text{if } Pe < 1, \\ \frac{h}{\|\beta\|}, & \text{if } Pe \geq 1. \end{cases} \tag{21}$$

This choice leads to the following corollary

Corollary 1 *Let (\bar{q}, \bar{u}) be the solution of the optimal control problem (1)–(3) with $Q_{ad} = Q$ and (\bar{q}_h, \bar{u}_h) be the solution of the discretized problem (15) and (16) with $Q_{ad,h} = Q_h$. Let moreover δ be chosen as in (21), then the following estimates hold:*

1. if $Pe < 1$

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \varepsilon^{1/2} h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^2 \|\bar{q}\|_{H^2(\Omega)},$$

2. if $Pe \geq 1$

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \|\beta\|^{1/2} h^{3/2} (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^2 \|\bar{q}\|_{H^2(\Omega)}.$$

Remark 5 In the case that $\beta \neq \text{Const}$ or in the case of varying cell diameter h_K the Peclet number has to be defined locally, i.e., cell-wise. Then, the stabilization parameter δ is also chosen locally.

Remark 6 We note, that our error estimate is proportional to $\frac{1}{\alpha}$. This is a typical situation if the coercivity constant of j coincides with the regularization parameter α , see (19). However, this is not always the case.

Let us consider a variant of the optimal control problem (15) and (16) with a finite dimensional control space $Q = \mathbb{R}^n$:

$$\text{Minimize } J(q, u) = \frac{1}{2} \|u - u_d\|_{L^2(\Omega)}^2 + \frac{\alpha}{2} \|q\|_Q^2, \quad u \in V, \quad q \in Q_{ad}$$

subject to

$$\begin{aligned} -\varepsilon \Delta u + \beta \cdot \nabla u + \sigma u &= f + \sum_{i=1}^n q_i g_i && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where $g_i \in L^\infty(\Omega)$ are linearly independent functions. This problem possesses a solution also in the case $\alpha = 0$. Moreover, there holds:

$$j''(\bar{q})(\delta q, \delta q) \geq \gamma \|\delta q\|_Q^2 \quad \text{with some } \gamma \geq \alpha.$$

For this problem similar error estimates can be obtained which are proportional to $\frac{1}{\gamma}$.

In order to formulate our main result for the control constrained case we need an additional assumption. We group the cells of the mesh \mathcal{T}_h in two classes $\mathcal{T}_h = \mathcal{T}_h^1 \cup \mathcal{T}_h^2$ with $\mathcal{T}_h^1 \cap \mathcal{T}_h^2 = \emptyset$ as follows: The cell K belongs to \mathcal{T}_h^1 if and only if one of the following conditions is satisfied:

- (a) $\bar{q} \equiv a$ on K ,
- (b) $\bar{q} \equiv b$ on K ,
- (c) $a < \bar{q} < b$ on K

The set \mathcal{T}_h^2 is given by $\mathcal{T}_h^2 = \mathcal{T}_h \setminus \mathcal{T}_h^1$ and consists of the cells which lie “close to the free boundary between the active and the inactive sets”.

Assumption 2 We assume that

$$\left| \bigcup_{K \in \mathcal{T}_h^2} K \right| \leq ch.$$

A similar assumption is used in [20] and in [24].

Remark 7 This assumption is valid if the boundary of the level sets

$$\{x : \bar{q}(x) = a\} \quad \text{and} \quad \{x : \bar{q}(x) = b\}$$

consists of a finite number of rectifiable curves.

In addition we assume $-\infty < a < b < +\infty$ and introduce the inactive set in optimum:

$$\Omega_I = \{x \in \Omega : a < \bar{q}(x) < b\}.$$

Note, that $\bar{q} = -\frac{1}{\alpha} \bar{z}$ on Ω_I and therefore $\bar{q}|_{\Omega_I} \in H^2(\Omega_I)$. Using this subdomain Ω_I we define a norm that will be needed later in the paper:

$$\|q\|_{2,ad} = \left(\|q\|_{W^{1,\infty}(\Omega)}^2 + \|\nabla^2 q\|_{L^2(\Omega_I)}^2 \right)^{1/2}.$$

In the following theorem we formulate our main result for the optimal control problem with control constraints.

Theorem 3 *Let (\bar{q}, \bar{u}) be the solution of the optimal control problem (1)–(3) and (\bar{q}_h, \bar{u}_h) be the solution of the discretized problem (15) and (16). Let moreover Assumption 2 be fulfilled. Then the following estimate holds:*

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \tau(\delta) h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^{3/2} \|\bar{q}\|_{2,ad}$$

with $s > d$ from Assumption 1 and $\tau(\delta)$ defined in (20).

The proof of this theorem is given in Sect. 5.

For the choice of δ according to (21) we obtain the following corollary:

Corollary 2 *Let (\bar{q}, \bar{u}) be the solution of the optimal control problem (1)–(3) and (\bar{q}_h, \bar{u}_h) be the solution of the discretized problem (15) and (16). Let moreover Assumption 2 be fulfilled and δ be chosen according to (21). Then the following estimates hold:*

1. if $Pe < 1$

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \varepsilon^{1/2} h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^{3/2} \|\bar{q}\|_{2,ad},$$

2. if $Pe \geq 1$

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \|\beta\|^{1/2} h^{3/2} (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^{3/2} \|\bar{q}\|_{2,ad}.$$

Remark 8 Theorems 2 and 3 provide error estimates for the error in the control variable. The corresponding estimates for the state and the adjoint variable have similar structure since there holds:

$$\|\bar{u} - \bar{u}_h\|_{L^2(\Omega)} + \|\bar{z} - \bar{z}_h\|_{L^2(\Omega)} \leq ch^2 (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + c\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)}.$$

4 Error analysis for the unconstrained problem

In this section we consider the unconstrained case, i.e., $Q_{ad} = Q$ and prove Theorem 2. To this end we need a special interpolation operator:

Lemma 3 *There is an interpolation operator $I_h : V \cap H^2(\Omega) \rightarrow V_h$ with the following properties: there holds for all $v \in V \cap H^2(\Omega)$*

$$(i) \quad (v - I_h v, w_h) = 0 \quad \forall w_h \in V_{2h}^{disc}, \tag{22}$$

$$(ii) \quad \|v - I_h v\|_{L^2(\Omega)} + h \|\nabla(v - I_h v)\|_{L^2(\Omega)} \leq ch^2 \|\nabla^2 v\|_{L^2(\Omega)}. \tag{23}$$

A constructive proof of this lemma can be found in [4].

In the sequel, we will need an assertion of the term $s_h^\delta(I_h v, I_h v)$. This assertion is given in the following lemma.

Lemma 4 *For any $v \in V \cap H^2(\Omega)$ the following estimate holds:*

$$|s_h^\delta(I_h v, I_h v)| \leq c \delta \|\beta\|^2 h^2 \|v\|_{H^2(\Omega)}^2.$$

Proof We start by

$$s_h^\delta(I_h v, I_h v) = s_h^\delta(v + I_h v - v, v + I_h v - v) \leq 2(s_h^\delta(v, v) + s_h^\delta(v - I_h v, v - I_h v))$$

For the first term we use the interpolation property of π_h , see (13):

$$\begin{aligned} s_h^\delta(v, v) &= \delta \|\beta \cdot \nabla v - \pi_h(\beta \cdot \nabla v)\|_{L^2(\Omega)}^2 \leq c \delta h^2 \|\nabla(\beta \cdot \nabla v)\|_{L^2(\Omega)}^2 \\ &\leq c \delta h^2 \|\beta\|^2 \|v\|_{H^2(\Omega)}^2. \end{aligned}$$

For the second term we use the L^2 -stability of π_h and obtain:

$$\begin{aligned} s_h^\delta(v - I_h v, v - I_h v) &\leq c \delta \|\beta \cdot \nabla(v - I_h v)\|_{L^2(\Omega)}^2 \\ &\leq c \delta \|\beta\|^2 \|\nabla(v - I_h v)\|_{L^2(\Omega)}^2 \leq c \delta \|\beta\|^2 h^2 \|v\|_{H^2(\Omega)}^2. \end{aligned}$$

This completes the proof.

In the following lemma we give an error estimate for the discretization of the state equation with an additional perturbation in the right hand side. To this end we first introduce a norm:

$$\|v\|^2 = \|v\|_{L^2(\Omega)}^2 + \|\varepsilon^{1/2} \nabla v\|_{L^2(\Omega)}^2 + s_h^\delta(v, v).$$

Lemma 5 *Let for $q \in Q$, $u = Sq \in V \cap H^2(\Omega)$ be the associated solution of the state equation (2) and (3), and for $p \in Q$, $w_h = S_h p \in V_h$ be the associated discrete solution, i.e:*

$$a_h(w_h, v_h) = (f + p, v_h) \quad \forall v_h \in V_h.$$

Then the following estimate holds:

$$\|u - w_h\| \leq \|q - p\|_{L^2(\Omega)} + c \tau(\delta) h \|u\|_{H^2(\Omega)},$$

where $\tau(\delta)$ is defined as in (20).

Proof For the error $u - w_h$ there holds a perturbed Galerkin orthogonality relation:

$$a(u - w_h, v_h) = s_h^\delta(w_h, v_h) + (q - p, v_h) \quad \forall v_h \in V_h.$$

We split the error $u - w_h = \eta + \xi$ with

$$\eta = u - I_h u, \quad \xi = I_h u - w_h$$

For the interpolation error η we obtain:

$$\|\eta\|_{L^2(\Omega)} \leq c h^2 \|u\|_{H^2(\Omega)} \tag{24}$$

and

$$|||\eta||| \leq c(h + \varepsilon^{1/2} + \delta^{1/2}\|\beta\|) h \|u\|_{H^2(\Omega)}. \tag{25}$$

For $\xi \in V_h$ we use the perturbed Galerkin orthogonality and obtain:

$$|||\xi|||^2 = a(\xi, \xi) + s_h^\delta(\xi, \xi) = (q - p, \xi) + s_h^\delta(I_h u, \xi) - a(\eta, \xi).$$

For the first term we have:

$$(q - p, \xi) \leq \|q - p\|_{L^2(\Omega)} \|\xi\|_{L^2(\Omega)} \leq \|q - p\|_{L^2(\Omega)} |||\xi|||.$$

For the second term we use Lemma 4:

$$s_h^\delta(I_h u, \xi) \leq (s_h^\delta(I_h u, I_h u))^{1/2} (s_h^\delta(\xi, \xi))^{1/2} \leq \delta^{1/2} h \|\beta\| \|u\|_{H^2(\Omega)} |||\xi|||.$$

For the third term we obtain:

$$|a(\eta, \xi)| \leq |||\eta||| |||\xi||| + |(\beta \cdot \nabla \eta, \xi)|.$$

The last term here can be estimated either using the property (22) of I_h

$$\begin{aligned} |(\beta \cdot \nabla \eta, \xi)| &= |(\eta, \beta \cdot \nabla \xi)| = |(\eta, \beta \cdot \nabla \xi - \pi_h(\beta \cdot \nabla \xi))| \\ &\leq \delta^{-1/2} \|\eta\|_{L^2(\Omega)} \delta^{1/2} \|\beta \cdot \nabla \xi - \pi_h(\beta \cdot \nabla \xi)\|_{L^2(\Omega)} \\ &\leq \delta^{-1/2} \|\eta\|_{L^2(\Omega)} |||\xi||| \end{aligned}$$

or directly:

$$\begin{aligned} |(\beta \cdot \nabla \eta, \xi)| &= |(\eta, \beta \cdot \nabla \xi)| \leq \varepsilon^{-1/2} \|\beta\| \|\eta\|_{L^2(\Omega)} \|\varepsilon^{1/2} \nabla \xi\|_{L^2(\Omega)} \\ &\leq \varepsilon^{-1/2} \|\beta\| \|\eta\|_{L^2(\Omega)} |||\xi|||. \end{aligned}$$

Therefore we have:

$$\begin{aligned} |||\xi||| &\leq \|q - p\|_{L^2(\Omega)} + \min(\delta^{-1/2}, \varepsilon^{-1/2} \|\beta\|) \|\eta\|_{L^2(\Omega)} \\ &\quad + \delta^{1/2} h \|\beta\| \|u\|_{H^2(\Omega)} + |||\eta|||. \end{aligned}$$

Now we use the interpolation estimates (24) and (25) and obtain the desired estimate.

As a direct consequence of the above lemma we obtain an estimate for the error in the adjoint states:

Lemma 6 *Let for $q \in \mathcal{Q}$, $z \in V \cap H^2(\Omega)$ be the associated adjoint solution, i.e the solution of (6) and (7), and for $p \in \mathcal{Q}$, $y_h \in V_h$ denote the associated adjoint discrete solution, then the following estimate holds:*

$$|||z - y_h||| \leq \|q - p\|_{L^2(\Omega)} + c \tau(\delta) h (\|u\|_{H^2(\Omega)} + \|z\|_{H^2(\Omega)}),$$

where $\tau(\delta)$ is defined as in (20).

The proof of this lemma is obtained similarly to the proof of Lemma 5.

Now, we come to the proof of Theorem 2.

Proof (Proof of Theorem 2) Let $p_h \in Q_h$ be arbitrary. We obtain from (19):

$$\begin{aligned} \alpha \|p_h - \bar{q}_h\|_{L^2(\Omega)}^2 &\leq j_h''(\bar{q}_h)(p_h - \bar{q}_h, p_h - \bar{q}_h) \\ &= j_h'(p_h)(p_h - \bar{q}_h) - j_h'(\bar{q}_h)(p_h - \bar{q}_h). \end{aligned}$$

Due to $Q_{ad} = Q$ and $Q_{ad,h} = Q_h$ we have:

$$j_h'(\bar{q}_h)(p_h - \bar{q}_h) = 0 = j'(\bar{q})(p_h - \bar{q}_h). \tag{26}$$

Hence,

$$\alpha \|p_h - \bar{q}_h\|_{L^2(\Omega)}^2 \leq j_h'(p_h)(p_h - \bar{q}_h) - j'(\bar{q})(p_h - \bar{q}_h).$$

We use Lemmas 1 and 2 and obtain:

$$\alpha \|p_h - \bar{q}_h\|_{L^2(\Omega)}^2 \leq (y_h, p_h - \bar{q}_h) - (\bar{z}, p_h - \bar{q}_h),$$

where \bar{z} is the associated adjoint state to \bar{q} and y_h is the associated discrete adjoint state to p_h . Using Lemma 6 we obtain:

$$\|p_h - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{1}{\alpha} \|\bar{q} - p_h\|_{L^2(\Omega)} + \frac{c}{\alpha} \tau(\delta) h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}),$$

where $\tau(\delta)$ is defined as in (20). Due to $Q_{ad} = Q$ we have that

$$\bar{q} = \frac{1}{\alpha} \bar{z} \in V \cap H^2(\Omega).$$

Therefore we can choose $p_h \in Q_h$ as the pointwise interpolation of \bar{q} with

$$\|\bar{q} - p_h\|_{L^2(\Omega)} \leq c h^2 \|\bar{q}\|_{H^2(\Omega)}.$$

Using this fact we obtain the desired estimate:

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} \tau(\delta) h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}) + \frac{c}{\alpha} h^2 \|\bar{q}\|_{H^2(\Omega)}.$$

5 Error analysis for the problem with control constraints

In this section we provide the error estimate for the control constrained case. In the case $Q_{ad} \neq Q$ we can not use the same argument as for the proof of Theorem 2 since the condition (26) does not hold anymore. To overcome this difficulty we will construct a special interpolation $p_h \in Q_{ad,h}$ of the solution $\bar{q} \in W^{1,\infty}(\Omega)$ which fulfills the following condition:

$$j'(\bar{q})(r - p_h) \geq 0 \quad \forall r \in Q_{ad}. \tag{27}$$

In the following lemma we use this condition in order to get an error estimate.

Lemma 7 *Let (\bar{q}, \bar{u}) be the solution of the optimal control problem (1)–(3) and (\bar{q}_h, \bar{u}_h) be the solution of the discretized problem (15) and (16). Let moreover $p_h \in Q_{ad,h}$ fulfill condition (27). Then the following estimate holds:*

$$\|\bar{q} - \bar{q}_h\|_{L^2(\Omega)} \leq \frac{1}{\alpha} \|\bar{q} - p_h\|_{L^2(\Omega)} + \frac{c}{\alpha} \tau(\delta) h (\|\bar{u}\|_{H^2(\Omega)} + \|\bar{z}\|_{H^2(\Omega)}),$$

where $\tau(\delta)$ is defined as in (20).

Proof As in the unconstrained case there holds:

$$\begin{aligned} \alpha \|p_h - \bar{q}_h\|_{L^2(\Omega)}^2 &\leq j'_h(\bar{q}_h)(p_h - \bar{q}_h, p_h - \bar{q}_h) \\ &= j'_h(p_h)(p_h - \bar{q}_h) - j'_h(\bar{q}_h)(p_h - \bar{q}_h). \end{aligned}$$

We use the discrete optimality condition (17) and the condition (27) with $r = \bar{q}_h$. We obtain:

$$-j'_h(\bar{q}_h)(p_h - \bar{q}_h) \leq 0 \leq -j'(\bar{q})(p_h - \bar{q}_h).$$

Hence,

$$\alpha \|p_h - \bar{q}_h\|^2 \leq j'_h(p_h)(p_h - \bar{q}_h) - j'(\bar{q})(p_h - \bar{q}_h).$$

Then we proceed as in the proof of Theorem 2 and obtain the desired estimate.

The interpolation $p_h \in Q_{ad,h}$ is constructed as follows: we prescribe the value of p_h at each node x of the mesh \mathcal{T}_h . To this end we use the “neighborhood” $\mathcal{N}_h(x_i)$ of a node x_i defined in Sect. 3. We set

$$p_h(x_i) = \begin{cases} a, & \text{if } \min_{x \in \mathcal{N}_h(x_i)} \bar{q}(x) = a \\ b, & \text{if } \max_{x \in \mathcal{N}_h(x_i)} \bar{q}(x) = b \\ \bar{q}(x_i), & \text{otherwise.} \end{cases} \tag{28}$$

A similar construction can be found in [7].

In the following lemma we show that p_h is well defined if h is small enough and that p_h fulfills condition (27).

Lemma 8 *Let*

$$h < \frac{b - a}{c_{\mathcal{N}} \|\bar{q}\|_{W^{1,\infty}(\Omega)}}. \tag{29}$$

Then p_h is well defined and fulfills condition (27).

Proof Due to the fact that $\bar{q} \in W^{1,\infty}(\Omega)$ there holds:

$$\max_{x \in \mathcal{N}_h(x_i)} \bar{q}(x) - \min_{x \in \mathcal{N}_h(x_i)} \bar{q}(x) \leq \|\bar{q}\|_{W^{1,\infty}(\Omega)} \text{diam}(\mathcal{N}_h(x_i)) \leq \|\bar{q}\|_{W^{1,\infty}(\Omega)} c_{\mathcal{N}} h,$$

with $c_{\mathcal{N}}$ introduced by (11).

Therefore, either the first or the second or the third case applies, provided h fulfills the condition (29). This proves that p_h is well defined.

In order to prove that p_h fulfills (27) we first consider a neighborhood $\mathcal{N}(x_i)$ such that

$$\min_{x \in \mathcal{N}(x_i)} \bar{q}(x) = a.$$

Then there holds:

$$\bar{q}(x) < b \quad \forall x \in \mathcal{N}(x_i)$$

and consequently $\nabla j(\bar{q}) \geq 0$ pointwise on $\mathcal{N}(x_i)$. Therefore we obtain for an arbitrary $r \in Q_{ad}$

$$\nabla j(\bar{q})(r - p_h) = \nabla j(\bar{q})(r - a) \geq 0$$

pointwise on $\mathcal{N}(x_i)$. For the cases

$$\max_{x \in \mathcal{N}(x_i)} \bar{q}(x) = b$$

and

$$\min_{x \in \mathcal{N}(x_i)} \bar{q}(x) > a \quad \text{and} \quad \max_{x \in \mathcal{N}(x_i)} \bar{q}(x) < b$$

we proceed similarly and obtain that $\nabla j(\bar{q})(r - p_h) \geq 0$ pointwise on Ω for all $r \in Q_{ad}$. This completes the proof.

To complete the proof of Theorem 3 we have to provide an estimate for the interpolation error $\|\bar{q} - p_h\|_{L^2(\Omega)}$.

Lemma 9 *Let $p_h \in Q_{ad,h}$ be constructed as in (28) and Assumption 2 be fulfilled, then the following estimate holds:*

$$\|\bar{q} - p_h\|_{L^2(\Omega)} \leq \frac{c}{\alpha} h^{3/2} \|\bar{q}\|_{2,ad},$$

with $s > d$ from Assumption 1.

Proof We start with

$$\|\bar{q} - p_h\|_{L^2(\Omega)}^2 = \sum_{K \in \mathcal{T}_h} \|\bar{q} - p_h\|_{L^2(K)}^2 = \sum_{K \in \mathcal{T}_h^1} \|\bar{q} - p_h\|_{L^2(K)}^2 + \sum_{K \in \mathcal{T}_h^2} \|\bar{q} - p_h\|_{L^2(K)}^2 \tag{30}$$

For the first sum we have:

$$\sum_{K \in \mathcal{T}_h^1} \|\bar{q} - p_h\|_{L^2(K)}^2 \leq \sum_{K \in \mathcal{T}_h^1} ch^4 \|\nabla^2 \bar{q}\|_{L^2(K)}^2 \leq ch^4 \|\nabla^2 \bar{q}\|_{L^2(\Omega_T)}^2, \tag{31}$$

since $\bar{q} \in H^2(K)$ for each cell $K \in \mathcal{T}_h^1$ and p_h is a usual point interpolation on the cells from \mathcal{T}_h^1 .

For a cell $K \in \mathcal{T}_h^2$ we obtain that either

$$(i) \quad \min_{x \in K} \bar{q}(x) = a \quad \text{and} \quad \max_{x \in K} \bar{q}(x) < b,$$

or

$$(ii) \quad \max_{x \in K} \bar{q}(x) = b \quad \text{and} \quad \min_{x \in K} \bar{q}(x) > a.$$

In the first case we obtain that for each node x_i of K (i.e., $K \subset \mathcal{N}_h(x_i)$)

$$\min_{x \in \mathcal{N}_h(x)} \bar{q}(x) = a,$$

and therefore $p(x_i) = a$ by definition (28). Hence, $p_h \equiv a$ on K . This implies:

$$\|\bar{q} - p_h\|_{L^2(K)}^2 = \|\bar{q} - a\|_{L^2(K)}^2.$$

Due to the fact that $\bar{q} \in W^{1,\infty}(\Omega)$ we have:

$$\|\bar{q} - a\|_{L^2(K)}^2 \leq |K| \|\bar{q} - a\|_{L^\infty(K)}^2 \leq |K| \text{diam}(K)^2 \|\bar{q}\|_{W^{1,\infty}(K)}^2.$$

The same estimate is obtained in the case (ii). By summing up and using Assumption 2, we obtain:

$$\sum_{K \in \mathcal{T}_h^2} \|\bar{q} - p_h\|_{L^2(K)}^2 \leq ch^2 \|\bar{q}\|_{W^{1,\infty}(\Omega)}^2 \sum_{K \in \mathcal{T}_h^2} |K| \leq ch^3 \|\bar{q}\|_{W^{1,\infty}(\Omega)}^2 \tag{32}$$

Inserting (31) and (32) in (30) we obtain the desired estimate.

To complete the proof of Theorem 3 we combine the results from Lemmas 7, 8 and 9.

6 Numerical example

In this section we present a numerical example elucidating our results. To this end we consider the optimization problem (1)–(3) on the unit square $\Omega = (0, 1)^2$ with the following parameters:

$$\varepsilon = 10^{-3}, \quad \beta = (-1, -2)^t, \quad \sigma = 1,$$

$$f = 1, \quad u_d = 1,$$

and

$$\alpha = 0.1, \quad a = 0.5, \quad b = 10.$$

The state and the control variables are discretized by bilinear finite elements as described in Sect. 3. The discretized control constraint problems are solved by primal dual active set method, see, e.g., [3, 18].


Figure 1 shows the optimal control \bar{q}_h computed on the mesh \mathcal{T}_h with $h = 2^{-6}\sqrt{2}$, i.e with Peclet number $Pe \approx 50$ with and without stabilization. It turns out that the solution of the stabilized problem has slight oscillations only in the boundary layer, whereas the solution of the discrete problem without stabilization shows strong oscillations almost in the whole domain.

Since the exact solution of the optimal control problem under consideration is not known, we show the evolution of the values of the cost functional $J(\bar{q}_h, \bar{u}_h)$ for a sequence of uniformly refined meshes \mathcal{T}_h , for h tending to zero. From this sequence, we compute the approximative order of convergence, see Table 1. The observed mean order of convergence is $r \approx 1.73$.

Table 1 Evolution of the values of the cost functional $J(\bar{q}_h, \bar{u}_h)$ for a sequence of uniformly refined meshes \mathcal{T}_h , for h tending to zero

$h/\sqrt{2}$	$J(\bar{q}_h, \bar{u}_h)$	$J(\bar{q}_h, \bar{u}_h) - J(\bar{q}_{2h}, \bar{u}_{2h})$	Order
2^{-2}	0.293169	–	–
2^{-3}	0.273468	$-1.97006e-2$	–
2^{-4}	0.265199	$-8.26853e-3$	1.25254
2^{-5}	0.262427	$-2.77226e-3$	1.57657
2^{-6}	0.261613	$-8.13916e-4$	1.76811
2^{-7}	0.261409	$-2.04352e-4$	1.99383
2^{-8}	0.261360	$-4.86628e-5$	2.07016
2^{-9}	0.261346	$-1.44169e-5$	1.75505

Fig. 1 Optimal control \bar{q}_h (level sets) computed on the mesh \mathcal{T}_h with $h = 2^{-6}\sqrt{2}$, $Pe \approx 50$ without stabilization (*top*) and with stabilization (*bottom*). The boundaries between active and inactive sets are indicated by the *black lines*



7 Conclusion

In this paper we have derived a priori error estimates for a stabilized finite element discretization of optimal control problems governed by convection-diffusion equations subject to pointwise control constraints. The method has optimal convergence behavior and has the commutativity property of discretization and optimization, which is very convenient from an algorithmic point of view. No modification of the Lagrangian is necessary. The presented analysis can be used to analyze related stabilization methods as the one proposed in [6].

References

1. Arada, N., Casas, E., Tröltzsch, F.: Error estimates for a semilinear elliptic optimal control problem. *Comput. Optim. Approx.* **23**, 201–229 (2002)
2. Becker, R., Braack, M.: A finite element pressure gradient stabilization for the stokes equations based on local projections. *Calcolo* **38**(4), 137–199 (2001)
3. Bergounioux, M., Ito, K., Kunisch, K.: Primal-dual strategy for constrained optimal control problems. *SIAM J. Control Optim.* **37**(4), 1176–1194 (1999)
4. Braack, M.: Solution of complex flow problems: Mesh- and model adaptation with stabilized finite elements. *Habilitationsschrift, Institut für Angewandte Mathematik, Universität Heidelberg* (2005)
5. Braack, M., Burman, E.: Local projection stabilization for the Oseen problem and its interpretation as a variational multiscale method. *SIAM J. Numer. Anal.* **43**(6), 2544–2566 (2006)

6. Burman, E., Hansbo, P.: Edge stabilization for Galerkin approximations of convection–diffusion–reaction problems. *Comput. Methods Appl. Mech. Eng.* **193**(15–16), 1437–1453 (2004)
7. Casas, E., Tröltzsch, F.: Error estimates for linear-quadratic elliptic control problems. In: Barbu, V., et al. (ed.) *Analysis and Optimization of Differential Systems*, pp. 89–100. Kluwer, Boston (2003)
8. Ciarlet, P.G.: The finite element method for elliptic problems. In: *Classics Appl. Math.*, vol. 40. SIAM, Philadelphia (2002)
9. Collis, S.S., Heinkenschloss, M.: Analysis of the streamline upwind/Petrov Galerkin method applied to the solution of optimal control problems. *CAAM TR02-01* (2002)
10. Dede, L., Quarteroni, A.: Optimal control and numerical adaptivity for advection-diffusion equations. *M2AN* **39**(5), 1019–1040 (2005)
11. Falk, R.: Approximation of a class of optimal control problems with order of convergence estimates. *J. Math. Anal. Appl.* **44**, 28–47 (1973)
12. Fursikov, A.V.: Optimal control of distributed systems: theory and applications. In: *Transl. Math. Monogr.*, vol. 187. AMS, Providence (1999)
13. Grisvard, P.: *Singularities in Boundary Value Problems*. Springer, Masson, Paris, Berlin (1992)
14. Guermond, J.L.: Stabilization of Galerkin approximations of transport equations by subgrid modeling. *Modél. Math. Anal. Numér.* **36**(6), 1293–1316 (1999)
15. Hinze, M.: A variational discretization concept in control constrained optimization: the linear-quadratic case. *Comput. Optim. Appl.* **30**(1), 45–61 (2005)
16. Johnson, C.: *Numerical Solution of Partial Differential Equations by Finite Element Method*. Cambridge University Press, Cambridge (1987)
17. Kozlov, V.A., Maz'ya, V.G., Rossmann, J.: Spectral problems associated with corner singularities of solutions to elliptic equations. In: *Mathematical Surveys and Monographs*, vol. 85. American Mathematical Society, Providence (2001)
18. Kunisch, K., Rösch, A.: Primal-dual active set strategy for a general class of constrained optimal control problems. *SIAM J. Optim.* **13**(2), 321–334 (2002)
19. Lions, J.L.: Optimal control of systems governed by partial differential equations. In: *Grundlehren Math. Wiss.*, vol. 170. Springer, Berlin (1971)
20. Meyer, C., Rösch, A.: Superconvergence properties of optimal control problems. *SIAM J. Control Optim.* **43**(3), 970–985 (2004)
21. Quarteroni, A., Valli, A.: Numerical approximation of partial differential equations. In: *Springer Series in Computational Mathematics*, vol. 23. Springer, Berlin (1994)
22. Roos, H.G., Stynes, M., Tobiska, L.: Numerical methods for singularly perturbed differential equations. In: *Springer Series in Computational Mathematics*, vol. 24. Springer, Berlin (1996)
23. Rösch, A.: Error estimates for linear-quadratic control problems with control constraints. *Optim. Methods Softw.* **21**(1), 121–134 (2005)
24. Rösch, A., Vexler, B.: Optimal control of the stokes equations: a priori error analysis for finite element discretization with postprocessing. *SIAM J. Numer. Anal.* **44**(5), 1903–1920 (2006)
25. Tröltzsch, F.: *Optimale Steuerung partieller Differentialgleichungen*. Friedr. Vieweg & Sohn Verlag, Wiesbaden (2005)
26. Zhou, G.: How accurate is the streamline diffusion finite element method? *Math. Comp.* **66**(217), 31–44 (1997)