

# FINITE ELEMENT ERROR ANALYSIS FOR MEASURE-VALUED OPTIMAL CONTROL PROBLEMS GOVERNED BY A 1D WAVE EQUATION WITH VARIABLE COEFFICIENTS

PHILIP TRAUTMANN<sup>1</sup>, BORIS VEXLER<sup>2</sup>, AND ALEXANDER ZLOTNIK<sup>3</sup>

ABSTRACT. This work is concerned with the optimal control problems governed by a 1D wave equation with variable coefficients and the control spaces  $\mathcal{M}_T$  of either measure-valued functions  $L^2(I, \mathcal{M}(\Omega))$  or vector measures  $\mathcal{M}(\Omega, L^2(I))$ . The cost functional involves the standard quadratic tracking terms and the regularization term  $\alpha \|u\|_{\mathcal{M}_T}$  with  $\alpha > 0$ . We construct and study three-level in time bilinear finite element discretizations for this class of problems. The main focus lies on the derivation of error estimates for the optimal state variable and the error measured in the cost functional. The analysis is mainly based on some previous results of the authors. The numerical results are included.

**Keywords.** Wave equation, optimal control, measure-valued control, vector measure control, finite element method, stability, error estimates.

**MSC subject classifications.** 65M60, 49K20, 49M05, 49M25, 49M29.

## 1. INTRODUCTION

This work is concerned with the discretization and numerical analysis of optimal control problems involving a 1D linear wave equation with variable coefficients and controls taking values in certain measure spaces. The combination of variable coefficients and irregular data leads to significant technical problems.

Motivated by industrial applications as well as applications in the natural sciences, in which one is interested to place actuators in form of point sources in an optimal way, see, e.g., [4, 9] or in the reconstruction of point sources from given measurements, see, e.g., [32, 42], measure valued optimal control problems involving PDEs gained attention in the last years. These problems can be translated into optimization problems in terms of the coordinates and coefficients of the point sources. However, these optimization problems are non-convex since the solution of the state equation (PDE) depends in a non-linear way on the coordinates of the point sources. Thus one has to deal with multiple local minima. Several authors suggested to cast the control problem resp. inverse problem in form of an optimization problem over a suitable measure space  $\mathcal{M}_T$  involving a convex regularization functional  $R$  which favors point sources as solutions. In our case we introduce the following problem formulation involving the 1D wave equation

$$\begin{aligned}
 J(u) = F(y) + R(u) &\rightarrow \min_{u \in \mathcal{M}_T} \\
 \text{subject to } \rho \partial_{tt} y - \partial_x(\kappa \partial_x y) &= u \quad \text{for } (t, x) \in I \times \Omega = (0, T) \times (0, L)
 \end{aligned} \tag{1.1}$$

with additional initial and boundary conditions. The functional  $F$  is given by a quadratic tracking functional involving  $y|_{I \times \Omega}$ ,  $y(T, \cdot)|_{\Omega}$  and  $\partial_t y(T, \cdot)|_{\Omega}$ . The regularization functional  $R$  and the control space  $\mathcal{M}_T$  are chosen in a way such that  $\mathcal{M}_T$  contains point sources of the desired form and  $R$  promotes controls of such a form, i.e. linear combinations of point sources with time-dependent

---

<sup>1</sup>DEPARTMENT OF MATHEMATICS AND SCIENTIFIC COMPUTING, UNIVERSITY OF GRAZ, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA.

<sup>2</sup>ZENTRUM MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, BOLTZMANNSTRASSE 3, 85748 GARCHING BEI MÜNCHEN, GERMANY.

<sup>3</sup>DEPARTMENT OF MATHEMATICS AT FACULTY OF ECONOMIC SCIENCES, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, MYASNITSKAYA 20, 101000 MOSCOW, RUSSIA.

*E-mail addresses:* philip.trautmann@uni-graz.at, vexler@ma.tum.de, azlotnik2008@gmail.com.

intensities or more general controls with a small spatial support. Since problem (1.1) is convex, one does not need to deal with several local minima. However, it is not longer guaranteed that the solution consists of a sum of point sources. We enforce such controls via the regularization functional  $R$ . Problems of the form (1.1) (also involving other PDEs) have been analysed from theoretical, numerical and algorithmic points of view, see [12, 11, 17, 18, 43, 31, 32, 13, 14, 7, 42, 16, 15]. Optimal control problems governed by the linear wave equation were discussed in several different aspects, see [33, 29, 30, 34, 48, 23, 22, 38, 39, 27, 28]. In our particular case we consider the control spaces  $\mathcal{M}_T$  of measure-valued functions  $L^2(I, \mathcal{M}(\Omega))$  and vector measures  $\mathcal{M}(\Omega, L^2(I))$  with  $R(u) = \alpha \|u\|_{\mathcal{M}_T}$ . These two different choices imply different structural properties of the optimal controls. A typical non-regular element from the space  $\mathcal{M}(\Omega, L^2(I))$  is given by

$$u = \sum_{i=1}^n u_i(t) \delta_{x_i}, \quad u_i \in L^2(I), \quad x_i \in \Omega, \quad (1.2)$$

where  $\delta_{x_i}$  are the Dirac delta functions. Point sources of such type with fixed positions and time-dependent intensities are of interest in acoustics or geology, see [32, 42]. If one is interested in controls involving moving point sources of the form

$$u = \sum_{i=1}^n u_i(t) \delta_{x_i(t)}, \quad u_i \in L^2(I), \quad x_i : I \rightarrow \Omega \text{ is measurable}, \quad (1.3)$$

then the control space  $L^2(I, \mathcal{M}(\Omega))$  rather than  $\mathcal{M}(\Omega, L^2(I))$  is more appropriate. The space  $\mathcal{M}(\Omega, L^2(I))$  and the functional  $\|\cdot\|_{\mathcal{M}(\Omega, L^2(I))}$  are also related to the term directional sparsity resp. joint sparsity, see [24, 20].

For the discretization of optimal control problem (1.1), we discretize the state equation by space-time finite element method as introduced in [47]. Related methods are also discussed and analyzed in [1, 21], see also [2]. The measure-valued control is not directly discretized, cf. the variational control discretization from [25]. However, there exists optimal controls consisting of Dirac measures in the spatial grid points which can be computed, see also [12, 31]. The numerical analysis of the control problem is based on FEM error estimates for the second order hyperbolic equations from [47] and techniques developed in [12, 31]. It requires to overcome significant technical difficulties caused by non-smoothness of controls and states. To the best of our knowledge, this is the first paper providing such numerical analysis for the studied control problems.

The problem like (1.1) for a parabolic/heat state equation is analyzed for the case  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  in [31] and for the case  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  in [12]. In particular, in both papers the authors prove existence of optimal controls and derive optimality conditions and FEM error estimates. Our analysis is partly based on these results of [31]. In [32] a problem similar to (1.1) involving the linear wave equation with constant coefficients as state equation is analyzed. In particular, existing regularity results for a Dirac right-hand side are extended to sources from  $\mathcal{M}(\Omega, L^2(I))$ . Based on these regularity results existence of optimal controls is proved as well as optimality conditions are derived in the 3D case.

Now we briefly sum up the contents of this work. First of all we collect and partially prove required existence and regularity results for the linear wave equation in the 1D setting. In particular, we check that the notions of a weaker solution defined in [47] and more commonly used very weak solution, e.g. [36], are equivalent. Most importantly we prove that the solution of the linear wave equation with variable coefficients from  $H^1(\Omega)$  for any source term  $u \in \mathcal{M}(\Omega, L^2(I))$  is an element of  $\mathcal{C}(\bar{I}, H_0^1(\Omega)) \cap \mathcal{C}^1(\bar{I}, L^2(\Omega))$  provided that the initial data have relevant regularity. The proof is based on a non-standard energy type bound in space, not only in time, cf. [35, 19]. In [32] the same result is proved for the wave equation with constant coefficients using duality techniques. This proof in [32] provides also corresponding results for multidimensional case but can not be directly extended for treating variable coefficients. This is due to the fact that it uses estimates of the solution of the wave equation in the whole space with a Dirac measure on the right hand-side which are proven using the Fourier-and Laplace-transformation or explicit solution formulas.

The existence of optimal controls and the derivation of optimality conditions are discussed on the basis of results from [32, 31]. In the case  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  we prove that the optimal control  $\bar{u}$  belongs to  $\mathcal{C}^1(\bar{I}, \mathcal{M}(\Omega))$ .

Further, the FEM discretization of the state equation is introduced. The state variable  $y_{h,\tau}$  belongs to the space of bilinear finite elements and is defined by the regularized Galerkin method. The resulting numerical scheme is a three-level method in time (i.e., its main equation relates the approximate solution values at three consecutive grid time levels). Moreover, we pose and prove the FEM error estimates in  $\mathcal{C}(\bar{I}, L^2(\Omega))$  for the discrete state equation which we need for the numerical analysis of the control problem. We base this study mainly on the results from [47] concerning error analysis of FEMs for the second order hyperbolic equations in the classes of the data having integer Sobolev or fractional Nikolskii order of smoothness. Note that their sharpness in a strong sense was stated in [46].

Then we consider a semi-discrete optimal control problem in which the continuous state equation is replaced by its discretized version whereas the controls are not discretized. We prove convergence of the discrete optimal controls to the continuous one and derive optimality conditions based on the Lagrange techniques. Most importantly we derive the discrete adjoint state equation. We can conclude that the first-discretize-then-optimize and first-optimize-then-discretize approaches commute. Therefore an analysis of the discrete adjoint state equation including the error estimates in  $\mathcal{C}(\bar{I} \times \bar{\Omega})$  and  $L^2(I, \mathcal{C}_0(\Omega))$  can also be based on techniques from [47]. Then we use results from [31] to represent the numerical error of state variable and of the cost functional in terms of FEM errors of the state equation and the adjoint state equation. Let  $\bar{u}$  and  $\bar{y}$  be the optimal control and the corresponding optimal state, and the variables  $\bar{u}_{\tau,h}$  and  $\bar{y}_{\tau,h}$  be their discrete counterparts. As the main result of this paper we prove the error estimates

$$\|\bar{y} - \bar{y}_{\tau,h}\|_{L^2(I \times \Omega)} = \mathcal{O}((\tau + h)^\alpha), \quad |J(\bar{u}) - J(\bar{u}_{\tau,h})| = \mathcal{O}((\tau + h)^{2/3})$$

where  $\tau$  is the step in time,  $h$  is the maximal step in space and  $\alpha = 1/3$  for  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  or  $\alpha = 2/3$  for  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ . The latter higher order is due to the above mentioned improved regularity results for the state and optimal control. Such estimates are proved for the measure-valued controls in the hyperbolic case for the first time. Similar estimates are impossible in multidimensional settings due to much less fractional Sobolev regularity of optimal states and controls.

Finally we discuss the numerical computation of the discrete control  $\bar{u}_{h,\tau}$ . Based on a control discretization  $u_{h,\tau}$  that given by the sum like (1.2) with  $x_i$  at the spatial grid points and  $u_i$  in the space of linear finite elements, a solution of the semi-discrete control problem can be calculated similarly to [31]. For the actual numerical computation of the optimal control we add the term  $(\gamma/2)\|u\|_{L^2(I \times \Omega)}^2$ ,  $\gamma > 0$ , to (1.1). This regularized problem is solved by a semi-smooth Newton method, see [41]. In a continuation strategy the regularization parameter  $\gamma$  is made sufficiently small. We complete this work with a numerical example for  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ .

The paper is organized in the following way. In Section 2 we introduce the problem setting and the control spaces resp. the regularization functionals. Section 3 is concerned with regularity properties of the linear wave equation with variable coefficients in the 1D setting. In Section 4 the control problem is analyzed from a theoretical point of view. Section 5 deals with discretization of the state equation. Then we obtain stability bounds and error estimates for the discrete state equation in Section 6. Section 7 is concerned with the analysis of the semi-discrete optimal control problem. The next section discusses stability bounds and error estimates for the discrete adjoint state equation. In Sections 9 resp. 10 error estimates for the optimal state and cost functional are derived being the main theoretical results of the study. Section 11 deals with the time stepping formulation of the discrete state equation. In Section 12 we discuss the control discretization with Dirac measures at the grid points. Then we introduce the  $L^2(I \times \Omega)$  regularized problem and describe its solutions by a semi-smooth Newton method. Finally Section 13 provides a numerical example.

## 2. PROBLEM SETTING

We consider optimal control problems of the following form

$$J(y, u) = F(y) + \alpha \|u\|_{\mathcal{M}_T} \rightarrow \min_{u, y} \quad (\mathcal{P})$$

with the parameter  $\alpha > 0$  and the tracking functional

$$F(y) := \frac{1}{2} (\|y - z_1\|_{L^2(I, H_\rho)}^2 + \|y(T) - z_2\|_{H_\rho}^2 + \|\rho \partial_t y(T) - z_3\|_{V_\kappa^*}^2)$$

using  $\mathbf{z} := (z_1, z_2, z_3) \in \mathcal{Y} := L^2(I \times \Omega) \times L^2(\Omega) \times H^{-1}(\Omega)$ , subject to *the state equation* which is an initial-boundary value problem for a 1D linear wave equation with variable coefficients

$$\begin{cases} \rho \partial_{tt} y - \partial_x(\kappa \partial_x y) = u & \text{in } I \times \Omega := (0, T) \times (0, L) \\ y = 0 & \text{on } I \times \partial\Omega \\ y = y^0, \partial_t y = y^1 & \text{in } \{0\} \times \Omega. \end{cases} \quad (2.1)$$

Here, in particular, the initial data  $\mathbf{y} := (y^0, y^1) \in H_0^1(\Omega) \times L^2(\Omega)$ , and  $L > 0$  and  $T > 0$ . The coefficients  $\rho, \kappa \in H^1(\Omega)$  satisfy  $\rho(x) \geq \nu > 0$  and  $\kappa(x) \geq \nu$  on  $\Omega$ .

For brevity we denote  $H = L^2(\Omega)$ ,  $V = H_0^1(\Omega)$ ,  $V^2 = H^2(\Omega) \cap V$  and  $V^3 = \{v \in V | \partial_x(\kappa \partial_x v) \in V\}$  equipped with the norms

$$\|\cdot\|_V = \|\partial_x \cdot\|_H, \quad \|\cdot\|_{V^2} = \|\partial_{xx} \cdot\|_H, \quad \|\cdot\|_{V^3} = \|\partial_x(\kappa \partial_x \cdot)\|_V.$$

Moreover, we utilize the equivalent coefficient-dependent Hilbert norms on  $H, V, V^*$  and  $\mathcal{Y}$

$$\begin{aligned} \|w\|_{H_\rho} &= \|\sqrt{\rho} w\|_H, \quad \|w\|_{V_\kappa} = \|\sqrt{\kappa} \partial_x w\|_H, \quad \|w\|_{V_\kappa^*} = \sup_{\|v\|_{V_\kappa} \leq 1} \langle w, v \rangle_\Omega, \\ \|\mathbf{z}\|_{\mathcal{Y}} &= (\|z_1\|_{L^2(I, H_\rho)}^2 + \|z_2\|_{H_\rho}^2 + \|z_3\|_{V_\kappa^*}^2)^{1/2}, \end{aligned}$$

where  $\langle \cdot, \cdot \rangle_\Omega$  is the duality relation on  $V^* \times V$ .

For the control space  $\mathcal{M}_T$  we consider two choices, either the space of vector measures  $\mathcal{M}(\Omega, L^2(I))$  or the space of weak measurable,  $\mathcal{M}(\Omega)$ -valued functions  $L^2(I, \mathcal{M}(\Omega)) := L_w^2(I, \mathcal{M}(\Omega))$ . Let correspondingly  $\mathcal{C}_T$  be chosen as  $\mathcal{C}_0(\Omega, L^2(I))$  or  $L^2(I, \mathcal{C}_0(\Omega))$  where  $\mathcal{C}_0(\Omega) = \{v \in \mathcal{C}(\bar{\Omega}) | v|_{x=0, L} = 0\}$ . The following identifications of dual spaces hold

$$\mathcal{C}_0(\Omega, L^2(I))^* \cong \mathcal{M}(\Omega, L^2(I)), \quad L^2(I, \mathcal{C}_0(\Omega))^* \cong L^2(I, \mathcal{M}(\Omega)),$$

i.e.  $\mathcal{M}_T = \mathcal{C}_T^*$ , see [12] resp. [31], where more details on the properties of these spaces and the norm  $\|\cdot\|_{\mathcal{M}_T}$  can be found. In particular, the following embeddings hold

$$\mathcal{M}(\Omega, L^2(I)) \hookrightarrow L^2(I, \mathcal{M}(\Omega)) \hookrightarrow L^2(I, V^*). \quad (2.2)$$

## 3. EXISTENCE AND REGULARITY OF THE STATE

**3.1. Weak formulations and preliminary existence, uniqueness and regularity results.** In this section we introduce our solution concepts for the state equation (2.1). We begin with defining a weak formulation of (2.1).

**Definition 3.1.** Let  $(u, y^0, y^1) \in X \times V \times H$  with  $X = L^2(I \times \Omega)$  or  $H^1(I, V^*)$  or  $\mathcal{M}(\Omega, L^2(I))$ . Then  $y \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  is called a *weak solution* of (2.1) if it satisfies the integral identity

$$B(y, v) + (\rho \partial_t y(T), v(T))_H = \int_I \langle u, v \rangle_\Omega dt + (\rho y^1, v(0))_H \quad \forall v \in L^2(I, V) \cap H^1(I, H) \quad (3.1)$$

with the indefinite symmetric bilinear form

$$B(y, v) := -(\rho \partial_t y, \partial_t v)_{L^2(I \times \Omega)} + (\kappa \partial_x y, \partial_x v)_{L^2(I \times \Omega)}, \quad (3.2)$$

and the initial condition  $y(0) = y^0$ .

The right-hand side in (3.1) is well defined for  $X = \mathcal{M}(\Omega, L^2(I))$  too due to embeddings (2.2).

**Remark 3.2.** It is possible (and more common) to suppose that  $v(T) = 0$  in (3.1) when the last term on the left disappears (for example, see [47]). This leads to an equivalent formulation. To check this, it is enough to replace there  $v$  by  $v\beta_\delta$ , where  $\beta_\delta(t) = \min(1, (T-t)/\delta)$ ,  $0 < \delta < T$ . Then  $\partial_t(v\beta_\delta) = (\partial_t v)\beta_\delta - (1/\delta)v\chi_{(T-\delta, T)}$ , where  $\chi_{(T-\delta, T)}$  is the characteristic function of  $(T-\delta, T)$ . Passing to the limit as  $\delta \rightarrow 0$  with the help of the dominated convergence theorem and the properties of  $y$  and  $v$  leads to the result.

Another definition of the weak solution is possible.

**Definition 3.3.** Let  $(u, y^0, y^1) \in X \times V \times H$  with  $X = L^2(I \times \Omega)$  or  $H^1(I, V^*)$  or  $\mathcal{M}(\Omega, L^2(I))$ . A function  $y \in \mathcal{C}(\bar{I}, V) \cap H^2(I, V^*) \hookrightarrow H^1(I, H)$  is called a weak solution of (2.1) if it satisfies

$$\int_I \langle \rho \partial_{tt} y, v \rangle_\Omega + (\kappa \partial_x y, \partial_x v)_H dt = \int_I \langle u, v \rangle_\Omega dt \quad \forall v \in L^2(I, V) \quad (3.3)$$

and  $y(0) = y^0$  as well as  $\partial_t y(0) = y^1$ .

**Proposition 3.4.** *Definitions 3.1 and 3.3 (up to the property  $y \in \mathcal{C}^1(\bar{I}, H)$ ) are equivalent.*

*Proof.* The weak solution from Definition 3.1 has  $\partial_{tt} y \in L^2(I, V^*)$  according to the integral identity (3.1). Then the equivalence of (3.3) and (3.1) can be proved using integration by parts in time and the density of  $\mathcal{C}^\infty(\bar{I}, V)$  in  $L^2(I, V) \cap H^1(I, H)$ , cf. [36, Chapter 1, Theorem 2.1].  $\square$

**Proposition 3.5.** (1) *Let  $(u, y^0, y^1) \in X \times V \times H$  with  $X = L^2(I \times \Omega)$  or  $H^1(I, V^*)$ . Then (2.1) has a unique weak solution satisfying  $y \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H) \cap H^2(I, V^*)$  and*

$$\|y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, H)} + \|\partial_{tt} y\|_{L^2(I, V^*)} \leq c (\|u\|_X + \|y\|_{V \times H}). \quad (3.4)$$

Hereafter  $c > 0$ ,  $c_1 > 0$ , etc., are independent of  $y$  and the data.

In the case  $X = H^1(I, V^*)$  there even holds  $y \in \mathcal{C}^2(\bar{I}, V^*)$  as well as

$$\|\partial_{tt} y\|_{\mathcal{C}(\bar{I}, V^*)} \leq c (\|u\|_{H^1(I, V^*)} + \|y\|_{V \times H}).$$

(2) *Let  $(u, y^0, y^1) \in X \times V^2 \times V$  with  $X = L^2(I, V)$  or  $H^1(I, H)$ . Then the weak solution  $y$  satisfies  $y \in \mathcal{C}(\bar{I}, V^2) \cap \mathcal{C}^1(\bar{I}, V) \cap H^2(I, H)$  and*

$$\|y\|_{\mathcal{C}(\bar{I}, V^2)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_{tt} y\|_{L^2(I, H)} \leq c (\|u\|_X + \|y\|_{V^2 \times V}). \quad (3.5)$$

In the case  $X = H^1(I, H)$  there even holds  $y \in \mathcal{C}^2(\bar{I}, H)$  as well as

$$\|\partial_{tt} y\|_{\mathcal{C}(\bar{I}, H)} \leq c (\|u\|_{H^1(I, H)} + \|y\|_{V^2 \times V}).$$

Moreover,  $y$  satisfies the equation  $\rho \partial_{tt} y - \partial_x(\kappa \partial_x y) = u$  in  $L^2(I \times \Omega)$ , i.e. it is the strong solution.

*Proof.* For example, see [47, Propositions 1.1 and 1.3].  $\square$

Item 2 ensures the regularity of weak solution for more regular data.

For less regular data  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$  one can use other weak formulations. To state the first of them, we define the integration operator  $(\mathcal{I}_t v)(t) := \int_0^t v(s) ds$  and its adjoint  $(\mathcal{I}_t^* v)(t) := \int_t^T v(s) ds$  on  $\bar{I}$ .

**Definition 3.6.** Let  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$ . A function  $y \in \mathcal{C}(\bar{I}, H)$  with  $\mathcal{I}_t y \in \mathcal{C}(\bar{I}, V)$  is called a *weaker solution* of (2.1) if it satisfies

$$\begin{aligned} & \int_I -(\rho y, \partial_t v)_H + (\kappa \partial_x \mathcal{I}_t y, \partial_x v)_H dt + (\rho y(T), v(T))_H \\ &= \int_I \langle u, \mathcal{I}_t^* v \rangle_\Omega dt + (\rho y^0, v(0))_H + \langle \rho y^1, (\mathcal{I}_t^* v)(0) \rangle_\Omega \quad \forall v \in L^2(I, V) \cap H^1(I, H). \end{aligned} \quad (3.6)$$

As in the case of Definition 3.1, it is sufficient to take  $v(T) = 0$  in (3.6), cf. Remark 3.2.

**Proposition 3.7.** *Let  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$ . Then there exists a unique weaker solution  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  and it satisfies the bound*

$$\|y\|_{\mathcal{C}(\bar{I}, H)} + \|\mathcal{I}_t y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, V^*)} \leq c (\|u\|_{L^2(I, V^*)} + \|y\|_{H \times V^*}).$$

*Proof.* See [47, Proposition 1.2]. □

We infer that there are other weak formulations of (2.1) for solutions  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$ . One can use the concept of very weak solutions.

**Definition 3.8.** Let  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$ . A function  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  satisfying

$$\begin{aligned} \int_I (y, \rho \partial_{tt} v - \partial_x(\kappa \partial_x v))_H dt - (\rho y(T), \partial_t v(T))_H + \langle \rho \partial_t y(T), v(T) \rangle_\Omega \\ = \int_I \langle u, v \rangle_\Omega dt - (\rho y^0, \partial_t v(0))_H + \langle \rho y^1, v(0) \rangle_\Omega \end{aligned} \quad (3.7)$$

for any  $v \in L^2(I, V^2) \cap H^2(I, H) \hookrightarrow H^1(I, V)$  is called a *very weak solution* of (2.1).

Actually, these two last solution concepts are equivalent for the considered data spaces.

**Theorem 3.9.** *Definitions 3.6 and 3.8 are equivalent.*

*Proof.* First of all, we consider the auxiliary integrated in  $t$  problem (2.1):

$$\begin{cases} \rho \partial_{tt} \tilde{y} - \partial_x(\kappa \partial_x \tilde{y}) = \mathcal{I}_t u + \rho y^1 & \text{in } I \times \Omega \\ \tilde{y} = 0 & \text{on } I \times \partial\Omega \\ \tilde{y} = 0, \partial_t \tilde{y} = y^0 & \text{in } \{0\} \times \Omega \end{cases} \quad (3.8)$$

for  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$ . Thus, we have  $\mathcal{I}_t u \in H^1(I, V^*)$ . According to Proposition 3.5 problem (3.8) has a unique weak solution  $\tilde{y} \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$ . Moreover, we set  $y = \partial_t \tilde{y}$ . Thus the weak formulation of (3.8) involving  $\tilde{y}$  coincides with the weaker formulation of (2.1) involving  $y$ . Furthermore there holds  $y = \partial_t \tilde{y} \in \mathcal{C}(\bar{I}, H)$  and

$$\partial_t y = \partial_{tt} \tilde{y} = (1/\rho)(\mathcal{I}_t u + \partial_x(\kappa \partial_x \tilde{y}) + \rho y^1) \in \mathcal{C}(\bar{I}, V^*). \quad (3.9)$$

Now we take any  $v \in \mathcal{C}^\infty(\bar{I}, V^2)$  and test (3.6) with  $-\partial_t v$  in the role of  $v$ :

$$\begin{aligned} \int_I (\rho y, \partial_{tt} v)_H - (\kappa \partial_x \mathcal{I}_t y, \partial_x \partial_t v)_H dt - (\rho y(T), \partial_t v(T))_H \\ = \int_I \langle u, -\mathcal{I}_t^*(\partial_t v) \rangle_\Omega dt - (\rho y^0, \partial_t v(0))_H + \langle \rho y^1, -(\mathcal{I}_t^* \partial_t v)(0) \rangle_\Omega. \end{aligned}$$

Next we rearrange a term on the left integrating by parts in  $x$  and  $t$ :

$$- \int_I (\kappa \partial_x \mathcal{I}_t y, \partial_x \partial_t v)_H dt = \int_I (\mathcal{I}_t y, L \partial_t v)_H dt = - \int_I (y, Lv)_H dt + ((\mathcal{I}_t y)(T), Lv(T))_H \quad (3.10)$$

with  $Lv := \partial_x(\kappa \partial_x v)$ . Since  $\mathcal{I}_t y \in \mathcal{C}(\bar{I}, V)$ , we get

$$\begin{aligned} \int_I (y, \rho \partial_{tt} v - \partial_x(\kappa \partial_x v))_H dt - (\rho y(T), \partial_t v(T))_H + \langle \partial_x(\kappa \partial_x \mathcal{I}_t y)(T), v(T) \rangle_\Omega \\ = \int_I \langle u, v \rangle_\Omega dt - \langle (\mathcal{I}_t u)(T), v(T) \rangle_\Omega - (\rho y^0, \partial_t v(0))_H + \langle \rho y^1, v(0) \rangle_\Omega - \langle \rho y^1, v(T) \rangle_\Omega. \end{aligned}$$

Since formula (3.9) implies that

$$\langle \rho \partial_t y(T), \varphi \rangle_\Omega = ((\mathcal{I}_t u)(T), \varphi)_H + \langle \partial_x(\kappa \partial_x \mathcal{I}_t y), \varphi \rangle_\Omega + \langle \rho y^1, \varphi \rangle_\Omega \quad \text{for any } \varphi \in V,$$

by the density of  $\mathcal{C}^\infty(\bar{I}, V^2)$  in  $L^2(I, V^2) \cap H^2(I, H)$  we find that  $y$  is a very weak solution of (2.1).

Now let  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  be a very weak solution of (2.1). Then we take any  $v \in \mathcal{C}^\infty(\bar{I}, V^2)$  and test (3.7) with  $\mathcal{I}_t^* v$ . Thus, we get

$$\begin{aligned} \int_I (y, -\rho \partial_t v)_H - (y, \partial_x(\kappa \partial_x \mathcal{I}_t^* v))_H dt + (\rho y(T), v(T))_H \\ = \int_I \langle u, \mathcal{I}_t^* v \rangle_\Omega dt + (\rho y^0, v(0))_H + \langle \rho y^1, (\mathcal{I}_t^* v)(0) \rangle_\Omega \end{aligned} \quad (3.11)$$

and then

$$\int_I (y, -\rho \partial_t v)_H + (\mathcal{I}_t y, -\partial_x(\kappa \partial_x v))_H dt + (\rho y(T), v(T))_H = \int_I \langle \mathcal{I}_t u + \rho y^1, v \rangle_\Omega dt + (\rho y^0, v(0))_H.$$

The last equation yields that  $L\mathcal{I}_t y = -\rho \partial_t y + \mathcal{I}_t u + \rho y^1 \in \mathcal{C}(\bar{I}, V^*)$ . Thus  $\mathcal{I}_t y \in \mathcal{C}(\bar{I}, V)$  and we can transform a term on the left in (3.11) by replacing  $v$  by  $\mathcal{I}_t^* v$  in (3.10):

$$\int_I (y, -\partial_x(\kappa \partial_x \mathcal{I}_t^* v))_H dt = \int_I (\kappa \partial_x \mathcal{I}_t y, \partial_x v)_H dt.$$

Then the density of  $\mathcal{C}^\infty(\bar{I}, V^2)$  in  $L^2(I, V) \cap H^1(I, H)$  shows that  $y$  is a weaker solution of (2.1).  $\square$

Moreover, there is the concept of solutions by transposition.

**Definition 3.10.** Let  $(u, y^0, y^1) \in L^2(I, V^*) \times H \times V^*$ . A *solution by transposition*  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  of (2.1) is defined by

$$\int_I (\rho y, \phi)_H dt - (\rho y(T), p^1)_H + \langle \rho \partial_t y(T), p^0 \rangle_\Omega = \int_I \langle u, p \rangle_\Omega dt + \langle \rho y^1, p(0) \rangle_\Omega - (\rho y^0, \partial_t p(0))_H \quad (3.12)$$

for all  $(\phi, p^0, p^1) \in L^2(I \times \Omega) \times V \times H$  where  $p \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  is the weak solution of *the adjoint problem*

$$\begin{cases} \rho \partial_{tt} p - \partial_x(\kappa \partial_x p) = \rho \phi & \text{in } I \times \Omega \\ p = 0 & \text{on } I \times \partial \Omega \\ p = p^0, \partial_t p = p^1 & \text{in } \{T\} \times \Omega \end{cases} \quad (3.13)$$

**Proposition 3.11.** *Definitions 3.8 and 3.10 are equivalent too.*

*Proof.* For  $\phi \in H^1(I, H)$  or  $L^2(I, V)$ ,  $p^0 \in V^2$  and  $p^1 \in V$  there holds  $p \in \mathcal{C}(\bar{I}, V^2) \cap \mathcal{C}^1(\bar{I}, V) \cap H^2(I, H)$ , see Proposition 3.5. Due to the density of  $H^1(I, H)$  resp.  $L^2(I, V)$  in  $L^2(I \times \Omega)$  as well as  $V^2$  in  $V$  and  $V$  in  $H$  a very weak solution is a solution by transposition. Now let  $p \in \mathcal{C}^\infty(\bar{I}, V^2)$  and set  $\phi = \partial_{tt} p - (1/\rho) \partial_x(\kappa \partial_x p) \in \mathcal{C}^\infty(\bar{I}, H)$ ,  $p^0 = p(T) \in V^2$  and  $p^1 = \partial_t p(T) \in V^2$ . Thus  $p$  is the solution of (3.13). Then the density of  $\mathcal{C}^\infty(\bar{I}, V^2)$  in  $L^2(I, V^2) \cap H^2(I, H)$  implies that a solution by transposition is a very weak solution.  $\square$

**Remark 3.12.** For  $(u, y^0, y^1) \in L^2(I, H) \times V \times H$ , the weaker solution coincides with the weak one.

**3.2. Existence and regularity of the state.** In this section we study the existence, uniqueness and regularity of solution of the state equation for measure valued source terms. We will carry out the analysis for both control spaces. We use the distinct properties of each space in order to show improved regularity of the state.

**3.2.1. The control space  $\mathcal{M}(\Omega, L^2(I))$ .** The space  $\mathcal{M}(\Omega, L^2(I))$  is not so broad as  $L^2(I, \mathcal{M}(\Omega))$  and contains no moving point sources but contains the standing  $\delta$ -sources (1.2). Therefore, we expect that the state has better regularity properties in this case and prove that  $y \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$ . The proof will be based on a priori bound and a density argument. First we state the following density result.

**Lemma 3.13.** *Let  $u \in \mathcal{M}(\Omega, L^2(I))$ . Then there exists a sequence  $\{u_n\} \subset C_c^\infty(\Omega, L^2(I))$  such that*

$$u_n \rightharpoonup^* u \text{ in } \mathcal{M}(\Omega, L^2(I)) \text{ as } n \rightarrow \infty, \quad \|u_n\|_{\mathcal{M}(\Omega, L^2(I))} \leq \|u\|_{\mathcal{M}(\Omega, L^2(I))} \quad \forall n \geq 1. \quad (3.14)$$

*Proof.* We denote by  $X$  the locally convex space  $\mathcal{M}(\Omega, L^2(I))$  endowed with its weak-star topology and define the absolutely convex set

$$E = \{u \in C_c^\infty(\Omega, L^2(I)) \mid \|u\|_{L^1(\Omega, L^2(I))} \leq 1\} \subset X.$$

Assume that (3.14) is wrong. Then there exists  $u_0 \in \mathcal{M}(\Omega, L^2(I))$ ,  $\|u_0\|_{\mathcal{M}(\Omega, L^2(I))} = 1$  such that  $u_0 \notin \bar{E}$  where  $\bar{E}$  is the closure of  $E$  in  $X$ . Owing to the corollary of a theorem on the separation of convex sets [26, Ch. III, Theorem 6] there exists  $v \in \mathcal{C}_0(\Omega, L^2(I))$  such that

$$|\langle u, v \rangle_{\mathcal{M}(\Omega, L^2(I)), \mathcal{C}_0(\Omega, L^2(I))}| \leq 1 \quad \forall u \in E, \quad 1 < \langle u_0, v \rangle_{\mathcal{M}(\Omega, L^2(I)), \mathcal{C}_0(\Omega, L^2(I))} \leq \|v\|_{\mathcal{C}_0(\Omega, L^2(I))}. \quad (3.15)$$

On the other hand,  $C_c^\infty(\Omega, L^2(I))$  is dense in  $L^1(\Omega, L^2(I))$  thus

$$\sup_{u \in E} |\langle u, v \rangle_{\mathcal{M}(\Omega, L^2(I)), \mathcal{C}_0(\Omega, L^2(I))}| = \|v\|_{\mathcal{C}_0(\Omega, L^2(I))}$$

that contradicts (3.15). Thus  $u_0 \in \bar{E}$ . Since  $\mathcal{C}_0(\Omega, L^2(I))$  is separable, the weak-star topology on  $E$  is metrizable. Therefore the closure of  $E$  is equal to its sequential closure, see [5, Theorem 3.28, Corollary 3.30]. Note that clearly  $C_c^\infty(\Omega, L^2(I)) \subset L^2(I, V)$ .  $\square$

Preliminarily we prove the following crucial a priori bound.

**Lemma 3.14.** *Let  $(u, y^0, y^1) \in L^1(\Omega, L^2(I)) \times V \times H$  and  $y$  be the corresponding strong solution of problem (2.1). Then  $y$  satisfies the following a priori bound*

$$\|y\|_{C(\bar{I}, V)} + \|\partial_t y\|_{C(\bar{I}, H)} + \|\kappa \partial_x y\|_{C(\bar{\Omega}, L^2(I))} + \|\partial_t y\|_{\mathcal{C}_0(\Omega, L^2(I))} \leq c(\|u\|_{L^1(\Omega, L^2(I))} + \|y\|_{V \times H}). \quad (3.16)$$

*Proof.* We first remind the energy equality for problem (2.1)

$$\|\sqrt{\rho} \partial_t y(t)\|_H^2 + \|\sqrt{\kappa} \partial_x y(t)\|_H^2 = \|\sqrt{\rho} y^1\|_H^2 + \|\sqrt{\kappa} \partial_x y^0\|_H^2 + 2\mathcal{I}_t(u(t), \partial_t y(t))_H \quad \text{on } I.$$

After setting

$$E(t) := \|\partial_t y(t)\|_H^2 + \|\partial_x y(t)\|_H^2, \quad E^0 := \|y^1\|_H^2 + \|\partial_x y^0\|_H^2, \quad c_0 := \max(\|\rho\|_{L^\infty(\Omega)}, \|\kappa\|_{L^\infty(\Omega)}),$$

the energy equality implies

$$\nu \|E\|_{C(\bar{I})} \leq c_0 E^0 + 2 \max_{\theta \in \bar{I}} \left| \int_\Omega \int_0^\theta u \partial_t y \, dt dx \right| \leq c_0 E^0 + 2\nu^{-1} \|u\|_{L^1(\Omega, L^2(I))} \|\sqrt{\rho \kappa} \partial_t y\|_{\mathcal{C}_0(\Omega, L^2(I))}. \quad (3.17)$$

We also multiply the equation in (2.1) by  $-2\kappa \partial_x y$  and integrate over  $I$ . Integration by parts in  $t$  yields the equality

$$\rho \kappa \partial_x \left( \|\partial_t y\|_{L^2(I)}^2 \right) + \partial_x \left( \|\kappa \partial_x y\|_{L^2(I)}^2 \right) = 2\rho \kappa \left( \partial_t y(T) \partial_x y(T) - y^1 \partial_x y^0 \right) - 2(u, \kappa \partial_x y)_{L^2(I)} \quad \text{on } \Omega. \quad (3.18)$$

We define a function  $P := \rho \kappa \|\partial_t y\|_{L^2(I)}^2 + \|\kappa \partial_x y\|_{L^2(I)}^2$  on  $\Omega$ . Since the left-hand side of (3.18) equals  $\partial_x P - (\partial_x(\rho \kappa)) \|\partial_t y\|_{L^2(I)}^2$ , taking the modulus and integrating over any  $(a, b) \subset \Omega$  we derive

$$\begin{aligned} \|\partial_x P\|_{L^1(a, b)} &\leq c_0^2 (E(T) + E^0) + 2\|u\|_{L^1(\Omega, L^2(I))} \|\kappa \partial_x y\|_{C(\bar{\Omega}, L^2(I))} + \|(\partial_x(\rho \kappa))\| \|\partial_t y\|_{L^2(I)}^2 \|P\|_{L^1(a, b)} \\ &\leq c_0^2 (\|E\|_{C(\bar{I})} + E^0) + 2\|u\|_{L^1(\Omega, L^2(I))} \|P\|_{C(\bar{\Omega})}^{1/2} + \nu^{-2} \|\partial_x(\rho \kappa)\|_{L^1(a, b)} \|P\|_{C(\bar{\Omega})}. \end{aligned} \quad (3.19)$$

Let  $x_0 \in \bar{\Omega}$  be such that  $\|P\|_{C(\bar{\Omega})} = P(x_0)$  hold and let now  $[a, b] \ni x_0$ . Then the mean value theorem for integrals implies

$$\|P\|_{C(\bar{\Omega})} \leq (b - a)^{-1} \|P\|_{L^1(a, b)} + \|\partial_x P\|_{L^1(a, b)}. \quad (3.20)$$

By the above definitions we clearly have

$$\|P\|_{L^1(\Omega)} \leq c_0^2 \|E\|_{L^1(I)} \leq c_0^2 T \|E\|_{C(\bar{I})}. \quad (3.21)$$



Inserting (3.19) into (3.20) and using (3.21), we obtain

$$\begin{aligned} \|P\|_{\mathcal{C}(\bar{\Omega})} &\leq c_0^2(1 + T(b-a)^{-1})(\|E\|_{\mathcal{C}(\bar{I})} + E^0) + 2\|u\|_{L^1(\Omega, L^2(I))} \|P\|_{\mathcal{C}(\bar{\Omega})}^{1/2} \\ &\quad + \nu^{-2}(b-a)^{1/2} \|\rho\kappa\|_{H^1(\Omega)} \|P\|_{\mathcal{C}(\bar{\Omega})}. \end{aligned} \quad (3.22)$$

Owing to (3.17) we can write

$$\nu\|E\|_{\mathcal{C}(\bar{I})} \leq c_0^2 E^0 + 2\nu^{-1}\|u\|_{L^1(\Omega, L^2(I))} \|P\|_{\mathcal{C}(\bar{\Omega})}^{1/2}. \quad (3.23)$$

Using this in (3.22) and choosing a small enough  $(a, b)$  such that  $\nu^{-2}(b-a)^{1/2} \|\rho\kappa\|_{H^1(\Omega)} \leq 1/2$ , we derive

$$\|P\|_{\mathcal{C}(\bar{\Omega})} \leq c_1(E^0 + \|u\|_{L^1(\Omega, L^2(I))}^2). \quad (3.24)$$

Inserting the last bound in (3.23), we also get

$$\|E\|_{\mathcal{C}(\bar{I})} \leq c_2(E^0 + \|u\|_{L^1(\Omega, L^2(I))}^2).$$

Finally, this yields bound (3.16).  $\square$

**Remark 3.15.** Lemma 3.14 remains valid for  $\rho, \kappa \in W^{1,1}(\Omega)$ . Owing to the absolute continuity of the Lebesgue integral we have  $\|\partial_x(\rho\kappa)\|_{L^1(a,b)} \leq \mu(b-a)$ , where  $\lim_{\theta \rightarrow +0} \mu(\theta) = 0$ , thus one can replace  $(b-a)^{1/2} \|\rho\kappa\|_{H^1(\Omega)}$  by  $\mu(b-a)$  in (3.22) and below in the proof.

**Theorem 3.16.** *Let  $(u, y^0, y^1) \in \mathcal{M}(\Omega, L^2(I)) \times V \times H$ . Then there exists a unique weak solution  $y$  and it satisfies the bound*

$$\|y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, H)} \leq c(\|u\|_{\mathcal{M}(\Omega, L^2(I))} + \|\mathbf{y}\|_{V \times H}). \quad (3.25)$$

*Proof.* 1. Let first  $u = 0$ . According to Proposition 3.5 there exists a unique weak solution  $y$  of (2.1) for any  $\mathbf{y} \in V \times H$  and it satisfies

$$\|y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, H)} \leq c\|\mathbf{y}\|_{V \times H}.$$

2. Now it suffices to consider the case  $y^0 = y^1 = 0$ . Let first  $u \in \mathcal{M}(\Omega, H^1(I)) \hookrightarrow H^1(I, V^*)$  since  $\partial_t u \in \mathcal{M}(\Omega, L^2(I)) \hookrightarrow L^2(I, V^*)$ . Then according to Proposition 3.5 there exists a unique weak solution  $y \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  of (2.1) and it satisfies bound (3.4). Moreover, it is also a weaker solution.

So it remains to prove the bound

$$\|y\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y\|_{\mathcal{C}(\bar{I}, H)} \leq c\|u\|_{\mathcal{M}(\Omega, L^2(I))} \quad \text{for } u \in \mathcal{M}(\Omega, H^1(I)). \quad (3.26)$$

To this end, according to Lemma 3.13 we approximate  $u$  by functions  $\{u_n\} \subset L^2(I, V)$  satisfying (3.14). The strong solution  $y_n$  of (2.1) corresponding to  $u = u_n$  satisfies the bound like (3.16) and in particular

$$\|y_n\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t y_n\|_{\mathcal{C}(\bar{I}, H)} \leq c\|u_n\|_{\mathcal{M}(\Omega, L^2(I))} \leq c\|u\|_{\mathcal{M}(\Omega, L^2(I))}.$$

Therefore there exists a subsequence of  $\{y_n\}$  (not relabelled) and  $\tilde{y} \in L^\infty(I, V) \cap W^{1,\infty}(I, H)$  such that  $y_n$  converges to  $\tilde{y}$  in the weak-star sense of  $L^\infty(I, V) \cap W^{1,\infty}(I, H)$ . This is sufficient to pass to the limit in the last bound and in (3.6) for  $y = y_n$ ,  $u = u_n$  and  $v(T) = 0$ , see Remark 3.2. Thus  $\tilde{y}$  both satisfies the bound

$$\|\tilde{y}\|_{L^\infty(\bar{I}, V)} + \|\partial_t \tilde{y}\|_{W^{1,\infty}(\bar{I}, H)} \leq c\|u\|_{\mathcal{M}(\Omega, L^2(I))}$$

and is a weaker solution of (2.1). Due to its uniqueness there holds  $\tilde{y} = y$ , and bound (3.26) is proved.

3. Let now  $u \in \mathcal{M}(\Omega, L^2(I))$  and  $y$  be the corresponding weaker solution of (2.1), see Proposition 3.7. Since  $\mathcal{M}(\Omega, H^1(I))$  is dense in  $\mathcal{M}(\Omega, L^2(I))$ , cf. [32, Proposition 2.1], there exists a sequence  $\{u_n\} \subset \mathcal{M}(\Omega, H^1(I))$  such that  $u_n \rightarrow u$  in  $\mathcal{M}(\Omega, L^2(I))$  as  $n \rightarrow \infty$ . Let  $y_n \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  be the above weak solution of (2.1) corresponding to  $u = u_n$ . Since  $\{u_n\}$  is a Cauchy sequence in

$\mathcal{M}(\Omega, L^2(I))$ ,  $\{y_n\}$  is a Cauchy sequence in  $\mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  too due to bound (3.26) for  $u = u_n$ . Thus  $y_n \rightarrow \hat{y}$  in  $\mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  and

$$\|\hat{y}\|_{\mathcal{C}(\bar{I}, V)} + \|\partial_t \hat{y}\|_{\mathcal{C}(\bar{I}, H)} \leq c \|u\|_{\mathcal{M}(\Omega, L^2(I))}.$$

Then we pass to the limit in (3.1) for  $y = y_n$ ,  $u = u_n$  and  $v(T) = 0$  and see that  $\hat{y}$  is a weak solution of (2.1). Due to uniqueness of the weaker solution we get  $\hat{y} = y$ , and the proof is complete.  $\square$

3.2.2. *The control space  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$ .* Recall that the space  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  contains the moving point sources (1.3). We set

$$H^{(-1)} = V^*, \quad H^{(0)} = H, \quad H^{(1)} = V, \quad H^{(2)} = V^2, \quad H^{(3)} = V^3$$

and introduce the interpolation spaces

$$H^{(\lambda)} := (H^{(\ell)}, H^{(\ell+1)})_{\lambda-\ell, \infty}, \quad \ell := \lfloor \lambda \rfloor,$$

for non-integer  $\lambda \in (-1, 3)$  using the real  $K_{\lambda, q}$ -interpolation method of Banach spaces for  $q = \infty$ , see [3]. Recall that the value  $q = \infty$  leads to the broadest intermediate spaces. Their explicit description in terms of the subspaces in the Nikolskii spaces is known, see [40, 45, 47]. In particular,

$$H^{(\lambda)} = H^{\lambda, 2}(\Omega) \quad \text{for } 0 < \lambda < \frac{1}{2}, \quad H^{(\lambda)} = \tilde{H}^{1/2, 2}(\Omega) \quad \text{for } \lambda = \frac{1}{2}, \quad H^{(\lambda)} = H_0^{\lambda, 2}(\Omega) \quad \text{for } \frac{1}{2} < \lambda < 1,$$

where  $\tilde{H}^{1/2, 2}(\Omega) := \{w \in H^{1/2, 2}(\Omega) \mid \text{ow} \in H^{1/2, 2}(-L, 2L)\}$  and  $\text{ow}$  is the odd extension of  $w$  with respect to  $x = 0, L$  outside  $\Omega$ . It is well known that the last space contains discontinuous but piecewise continuously differentiable functions.

**Lemma 3.17.** *The following embeddings hold*

$$L^2(I, \mathcal{M}(\Omega)) \hookrightarrow L^2(I, H^{(-1/2)}), \quad (3.27)$$

$$\mathcal{C}^1(\bar{I}, \mathcal{M}(\Omega)) \hookrightarrow H^1(I, H^{(-1/2)}). \quad (3.28)$$

*Proof.* Both the embeddings follow from  $\mathcal{M}(\Omega) \hookrightarrow H^{(-1/2)}$ . Due to [3] this last embedding is equivalent to the following inequality, which we show in the rest of the proof:

$$\sup_{0 < h < L} h^{-1/2} \inf_{v \in H} (\|w - v\|_{V^*} + h\|v\|_H) \leq c \|w\|_{\mathcal{M}(\Omega)} \quad \forall w \in \mathcal{M}(\Omega).$$

Equivalently, we have to show, that for any  $0 < h < L$ , there exists  $w_h \in H$  such that

$$\|w - w_h\|_{V^*} \leq c_1 h^{1/2} \|w\|_{\mathcal{M}(\Omega)}, \quad \|w_h\|_H \leq c_1 h^{-1/2} \|w\|_{\mathcal{M}(\Omega)} \quad \forall w \in \mathcal{M}(\Omega). \quad (3.29)$$

Any  $w \in \mathcal{M}(\Omega)$  can be represented as  $w = D_x W$  with  $\|w\|_{\mathcal{M}(\Omega)} = \text{var}_{\bar{\Omega}} W$ , where  $D_x$  is the distributional derivative of a function  $W \in NBV(\Omega)$  [10, Ch. 2]. Here  $NBV(\Omega)$  is the space of normalized functions of bounded variation on  $\bar{\Omega}$  that are continuous from the right at  $x = 0$  and continuous from the left at any  $x \in (0, L]$ . Notice that the following inequalities hold

$$\|W - W(0)\|_{L^\infty(\Omega)} \leq \text{var}_{\bar{\Omega}} W, \quad \sup_{0 < h < L} \|\delta_{-h} W\|_{L^1(h, L)} \leq \text{var}_{\bar{\Omega}} W \quad \forall W \in NBV(\Omega), \quad (3.30)$$

where  $\delta_{-h} W(x) := h^{-1}(W(x) - W(x - h))$  is the backward difference quotient (the latter inequality follows from the definition of the Riemann integral).

We can choose  $W(0) = 0$  and extend  $W(x) = 0$  for  $x < 0$ . Then for  $0 < h < L$  we define the backward average  $W_h(x) := h^{-1} \int_{-h}^0 W(x + \xi) \, \mathbf{d}\xi$  and set  $w_h := \partial_x W_h = \delta_{-h} W$ . Owing to the inequalities

$$\|\varphi\|_H \leq \|\varphi\|_{L^\infty(\Omega)}^{1/2} \|\varphi\|_{L^1(\Omega)}^{1/2} \quad \forall \varphi \in L^\infty(\Omega)$$

and (3.30) we can prove estimates (3.29):

$$\begin{aligned} \|w - w_h\|_{V^*} &\leq \|W - W_h\|_H \leq (\|W\|_{L^\infty(\Omega)} + \|W_h\|_{L^\infty(\Omega)})^{1/2} \left\| \int_{-h}^0 |\delta_\xi W(x)| \, \mathbf{d}\xi \right\|_{L^1(\Omega)}^{1/2} \\ &\leq (2\|W\|_{L^\infty(\Omega)})^{1/2} (h \sup_{0 < h < L} \|\delta_{-h} W\|_{L^1(\Omega)})^{1/2} \leq ch^{1/2} \text{var}_{\bar{\Omega}} W, \end{aligned}$$

$$\|w_h\|_H = \|\delta_{-h}W\|_H \leq (2h^{-1}\|W\|_{L^\infty(\Omega)})^{1/2}\|\delta_{-h}W\|_{L^1(\Omega)}^{1/2} \leq ch^{-1/2}\text{var}_{\bar{\Omega}}W.$$

□

## 4. ANALYSIS OF THE CONTROL PROBLEM

According to Theorem 3.16 and Proposition 3.7 the state equation (2.1) is uniquely solvable for any  $u$  in either  $\mathcal{M}(\Omega, L^2(I))$  or  $L^2(I, \mathcal{M}(\Omega))$  and the solution  $y$  depends continuously on the data. Therefore, we can introduce the linear and bounded operator  $\hat{S}: (u, y^0, y^1) \mapsto (y, y(T), \rho\partial_t y(T))$ . The control-to-state mapping

$$S: \mathcal{M}_T \rightarrow \mathcal{Y}, \quad u \mapsto (y, y(T), \rho\partial_t y(T))$$

is given by  $Su = \hat{S}(u, 0, 0) + \hat{S}(0, y^0, y^1)$  for fixed  $y^0$  and  $y^1$  and it is an affine and bounded operator. So we can rewrite the original control problem ( $\mathcal{P}$ ) in its reduced form

$$j(u) = \frac{1}{2}\|Su - \mathbf{z}\|_{\mathcal{Y}}^2 + \alpha\|u\|_{\mathcal{M}_T} \rightarrow \min_{u \in \mathcal{M}_T}.$$

**Proposition 4.1.** *Problem ( $\mathcal{P}$ ) has a unique solution  $\bar{u} \in \mathcal{M}_T$ .*

*Proof.* The control-to-state operator  $S$  is weak-star-to-strong sequential continuous, i.e., if  $\{u_n\} \subset \mathcal{M}_T$  and  $u_n \rightharpoonup^* u$  in  $\mathcal{M}_T$ , then  $Su_n \rightarrow Su$  in  $\mathcal{Y}$ . The proof of this continuity property is similar to [32, Lemma 6.1] in the case of solutions by transposition resp. very weak solutions. The strong continuity follows from the compact embeddings and well known Aubin-Lions-Lemma. Then the direct method of calculus of variations combined with the sequential Banach-Alaoglu theorem ( $\mathcal{C}_T$  is separable) can be applied to show existence of an optimal control. Additionally the control is unique since the control-to-state operator  $S$  is injective and the data tracking functional is strictly convex. □

Owing to Proposition 3.7 the optimal control  $\bar{u} \in \mathcal{M}_T$  satisfies the inequalities

$$\alpha\|\bar{u}\|_{\mathcal{M}_T} \leq j(\bar{u}) \leq j(0) = \frac{1}{2}\|S(0) - \mathbf{z}\|_{\mathcal{Y}}^2 \leq c(\|y^0\|_H + \|y^1\|_{V^*} + \|\mathbf{z}\|_{\mathcal{Y}})^2 \quad (4.1)$$

and thus

$$\|\bar{u}\|_{\mathcal{M}_T} \leq c(\|\mathbf{y}\|_{H \times V^*} + \|\mathbf{z}\|_{\mathcal{Y}})^2 \leq C. \quad (4.2)$$

Hereafter  $C > 0$  depends on the norms of data.

Next we discuss first order optimality conditions. We introduce the adjoint control-to-solution operator  $S^*: \mathcal{Y} \rightarrow C(\bar{I}, V) \hookrightarrow \mathcal{C}_T$ ,  $(\phi, p^1, p^0) \mapsto p$  where  $p$  is a weak solution of (3.13). This operator is well defined and bounded according to Proposition 3.5.

We also need the operator  $A^{-1}: V^* \rightarrow V$ ,  $f \mapsto w$  where  $w \in V$  is the unique solution of

$$(\kappa\partial_x w, \partial_x v)_H = \langle f, v \rangle_{\Omega} \quad \forall v \in V. \quad (4.3)$$

The next result provides the necessary and sufficient optimality condition for the optimal pair  $(\bar{p}, \bar{u})$ .

**Proposition 4.2.** *An element  $\bar{u} \in \mathcal{M}_T$  is an optimal control of ( $\mathcal{P}$ ) if and only if*

$$-\bar{p} \in \alpha\partial\|\bar{u}\|_{\mathcal{M}_T}, \quad (4.4)$$

or equivalently

$$\langle -\bar{p}, u - \bar{u} \rangle_{\mathcal{C}_T, \mathcal{M}_T} + \alpha\|\bar{u}\|_{\mathcal{M}_T} \leq \alpha\|u\|_{\mathcal{M}_T} \quad \forall u \in \mathcal{M}_T \quad (4.5)$$

where  $\bar{p} = S^*(\bar{y} - z_1, -(\bar{y}(T) - z_2), A^{-1}(\rho\partial_t \bar{y} - z_3))$  with  $(\bar{y}, \bar{y}(T), \rho\partial_t \bar{y}(T)) = \hat{S}(\bar{u}, y^0, y^1)$ .

*Proof.* For  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  the proof of [32, Theorem 7.1] remains valid; for  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  it is similar to [12, Theorem 3.2]. □

To discuss further the properties of the optimal control  $\bar{u}$ , we introduce the Jordan decomposition of a signed measure  $\mu \in \mathcal{M}(\Omega)$ , see [6]. There exists unique elements  $\mu^\pm \in \mathcal{M}(\Omega)^+$  such that  $\mu = \mu^+ - \mu^-$ . Moreover, we recall the polar decomposition of a vector measure  $\mu \in \mathcal{M}(\Omega, L^2(I))$ :  $d\mu = \mu' d|\mu|$ , where  $\mu'$  is the Radon-Nikodym-derivative of  $\mu$  with respect to  $|\mu|$ .

The subgradient condition in Proposition 4.2 implies the following conditions.

**Proposition 4.3.** *Let  $\bar{u} \in \mathcal{M}_T$  be the optimal control of  $(\mathcal{P})$  and  $\bar{p} \in \mathcal{C}_T$  be the corresponding adjoint state. Then there holds  $\|\bar{p}\|_{\mathcal{C}_T} \leq \alpha$ .*

*In the cases  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  and  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  there respectively hold*

$$\text{supp } \bar{u}^\pm(t) \subset \{x \in \Omega \mid \bar{p}(t, x) = \mp \|\bar{p}(t, \cdot)\|_{\mathcal{C}_0(\Omega)}\} \quad \text{for a.a. } t \in I$$

and

$$\text{supp } |\bar{u}| \subset \{x \in \Omega \mid \|\bar{p}(\cdot, x)\|_{L^2(I)} = \alpha\}, \quad \bar{u}' = -\alpha^{-1}\bar{p} \quad \text{in } L^1((\Omega, |\bar{u}|), L^2(I)). \quad (4.6)$$

*Proof.* A detailed discussion of the proof of these results can be found in [12, 31].  $\square$

The regularity of the adjoint state  $\bar{p}$  is now applied to show improved regularity of the optimal control  $\bar{u}$ .

**Theorem 4.4.** *Let  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ ,  $\mathbf{z} \in \mathcal{Y}^1 := L^2(I, V) \times V \times H$ ,  $\mathbf{y} \in V \times H$  and  $\bar{u}$  be the optimal control of  $(\mathcal{P})$ . Then  $\bar{u} \in \mathcal{C}^1(\bar{I}, \mathcal{M}(\Omega))$  and the following bound holds*

$$\|\bar{u}\|_{\mathcal{C}^1(\bar{I}, \mathcal{M}(\Omega))} \leq C = C(\|\mathbf{y}\|_{V \times H}, \|\mathbf{z}\|_{\mathcal{Y}^1}).$$

*Proof.* There holds  $\bar{y} \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  according to Theorem 3.16. Thus, the optimal adjoint state has the following regularity  $\bar{p} \in \mathcal{C}(\bar{I}, V^2) \cap \mathcal{C}^1(\bar{I}, V)$  by Proposition 3.5. We have  $\bar{u} = -\alpha^{-1}\bar{p}|\bar{u}|$  according to (4.6). Moreover, we define the function

$$w = -\alpha^{-1}(\partial_t \bar{p})|\bar{u}|$$

and show that it serves the time derivative of  $\bar{u}$ . For any  $t_0, t \in \bar{I}$  and  $t_0 \neq t$ , we define the difference quotient  $\bar{u}(t_0, t) = (\bar{u}(t) - \bar{u}(t_0))/(t - t_0)$ . Then we consider

$$\begin{aligned} \|\bar{u}(t_0, t) - w(t_0)\|_{\mathcal{M}(\Omega)} &= \alpha^{-1} \sup_{\|\phi\|_{\mathcal{C}_0(\Omega)} \leq 1} \int_{\Omega} (\bar{p}(t_0, t) - \partial_t \bar{p}(t_0)) \phi \, d|\bar{u}| \\ &\leq \alpha^{-1} \|\bar{p}(t_0, t) - \partial_t \bar{p}(t_0)\|_{\mathcal{C}_0(\Omega)} \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \\ &\leq c\alpha^{-1} \|\bar{p}(t_0, t) - \partial_t \bar{p}(t_0)\|_V \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \rightarrow 0 \end{aligned}$$

as  $t \rightarrow t_0$  since  $\bar{p} \in \mathcal{C}^1(\bar{I}, V)$ . Next, quite similarly we get

$$\|w(t) - w(t_0)\|_{\mathcal{M}(\Omega)} \leq c\alpha^{-1} \|\partial_t \bar{p}(t) - \partial_t \bar{p}(t_0)\|_V \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \rightarrow 0$$

as  $t \rightarrow t_0$ . Consequently  $\partial_t \bar{u} = w \in \mathcal{C}(\bar{I}, \mathcal{M}(\Omega))$ . Finally, we bound  $\partial_t \bar{u}$  as follows

$$\begin{aligned} \|\partial_t \bar{u}\|_{\mathcal{C}(\bar{I}, \mathcal{M}(\Omega))} &\leq c\alpha^{-1} \|\partial_t \bar{p}\|_{\mathcal{C}(\bar{I}, V)} \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \\ &\leq c_1 \alpha^{-1} (\|\bar{y} - z_1\|_{L^2(I, V)} + \|\bar{y}(T) - z_2\|_V + \|\rho \partial_t \bar{y}(T) - z_3\|_H) \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \\ &\leq c_2 \alpha^{-1} (\|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} + \|\mathbf{y}\|_{V \times H} + \|\mathbf{z}\|_{\mathcal{Y}^1}) \|\bar{u}\|_{\mathcal{M}(\Omega, L^2(I))} \end{aligned}$$

owing to Proposition 3.5 and Theorem 3.16. Utilizing bound (4.2) for  $\bar{u}$ , we complete the proof.  $\square$

## 5. DISCRETIZATION OF THE STATE EQUATION

We introduce the uniform grid  $t_m = m\tau$  in time with the step  $\tau = T/M$  and a non-uniform grid  $0 = x_0 < x_1 < \dots < x_N = L$  in space with the steps  $h_j = x_j - x_{j-1}$ , where  $M \geq 2$  and  $N \geq 2$ . Let also  $h = \max_{j=1, \dots, N} h_j$ ,  $h_{\min} = \min_{j=1, \dots, N} h_j$  and  $\vartheta = (\tau, h)$ . We assume that the space grid is *quasi-uniform*, i.e.,  $h \leq c_1 h_{\min}$ . Hereafter  $c, c_1, C$ , etc., are grid-independent.

Let  $V_\tau \subset H^1(I)$  and  $V_h \subset V$  be the spaces of piecewise linear finite elements with respect to the introduced grids on  $\bar{I}$  and  $\bar{\Omega}$ .

We approximate the state variable  $y$  by  $y_\vartheta \in V_\vartheta := V_\tau \otimes V_h \subset H^1(I, V)$  and additionally  $\partial_t y(T)$  by  $y_{T_h}^1 \in V_h$ . For  $(u, y^0, y^1) \in \mathcal{M}_T \times H \times V^*$  the discrete state equation has the following form

$$B_\sigma(y_\vartheta, v) + (\rho y_{T_h}^1, v(T))_H = \langle u, v \rangle_{\mathcal{M}_T, \mathcal{C}_T} + \langle \rho y^1, v(0) \rangle_\Omega \quad \forall v \in V_\vartheta, \quad (5.1)$$

$$(\rho y_\vartheta(0), \varphi)_H = (\rho y^0, \varphi)_H \quad \forall \varphi \in V_h, \quad (5.2)$$

involving the indefinite symmetric bilinear form

$$B_\sigma(y, v) := -(\rho \partial_t y, \partial_t v)_{L^2(I \times \Omega)} - (\sigma - \frac{1}{6}) \tau^2 (\kappa \partial_x \partial_t y, \partial_x \partial_t v)_{L^2(I \times \Omega)} + (\kappa \partial_x y, \partial_x v)_{L^2(I \times \Omega)}, \quad (5.3)$$

with the grid independent parameter  $\sigma$ , cf. (3.1). This definition follows [47] but notice carefully that normally  $y_\vartheta$  is uniquely defined by (5.1) with  $v(T) = 0$  and (5.2). To treat general  $v$ , we need  $y_{T_h}^1$ .

**Remark 5.1.** The second term on the right hand-side of (5.3) regularizes the Galerkin (i.e. projection) method with respect to bilinear form (3.2). It is included to ensure unconditional stability for suitable values of  $\sigma$ . Moreover, the term  $-(1/6) \tau^2 (\kappa \partial_x \partial_t y, \partial_x \partial_t v)_{L^2(I \times \Omega)}$  is the error term of the compound trapezoidal rule applied for the calculation of the temporal integral in  $(\kappa \partial_x y, \partial_x v)_{L^2(I \times \Omega)}$ . So that, in particular, for  $\sigma = 0$  in (5.3) this temporal integral is calculated using this rule whereas for  $\sigma = 1/6$  it is not approximated.

Next we recall the inverse inequality

$$\|\varphi\|_{V_\kappa} \leq \alpha_h \|\varphi\|_{H_\rho} \quad \forall \varphi \in V_h \quad (5.4)$$

where the least constant satisfies  $c_1 h^{-1} \leq \alpha_h \leq c_2 h^{-1}$  for the quasi-uniform grid. For  $\sigma \leq 1/4$  we need to state conditions linking the temporal and spatial grids to ensure stability of the numerical method.

**Assumption 5.2.** In what follows, let

$$\text{if } \sigma < \frac{1}{4}, \text{ then } \tau^2 \alpha_h^2 (\frac{1}{4} - \sigma) \leq 1 - \varepsilon_0^2 \text{ for some } 0 < \varepsilon_0 < 1, \quad (5.5)$$

$$\text{if } \sigma \leq \frac{1}{4}, \text{ then } \tau^2 \alpha_h^2 (\frac{1+\varepsilon_1^2}{4} - \sigma) \leq 1 \text{ for some } 0 < \varepsilon_1 \leq 1. \quad (5.6)$$

**Remark 5.3.** The parameters  $\varepsilon_0$  and  $\varepsilon_1$  can be chosen arbitrarily small but then constants in the stability and error estimates for our FEM can tend to infinity.

**Remark 5.4.** As we see below in Section 11, the method is related to well known time-stepping methods, in particular, to the explicit Leap-Frog-method for  $\sigma = 0$ . Then conditions (5.5) and (5.6) reduce to a CFL-type one  $\tau \alpha_h \leq 2\sqrt{1 - \varepsilon_0^2}$ . For  $\sigma = 1/4$  the method is related to the Crank-Nicolson scheme and is unconditionally stable but in a weaker norm than we need to derive our error estimates so that we impose a very weak CFL-type condition  $\tau \alpha_h \leq 2/\varepsilon_1$ .

Below in proofs we utilize the auxiliary squared norms

$$\|\varphi\|_{H_\rho^0}^2 := \|\varphi\|_{H_\rho}^2 + (\sigma - \frac{1}{4}) \tau^2 \|\varphi\|_{V_\kappa}^2, \quad \|y\|_{C_\tau(H_E)}^2 = \max_{1 \leq m \leq M} \left( \frac{1}{\tau} \|y_m - y_{m-1}\|_{H_\rho^0}^2 + \frac{1}{2} \|y_m + y_{m-1}\|_{V_\kappa}^2 \right)$$

for  $\varphi \in V_h$  and  $y \in V_\tau \otimes V_h$ . We need to bound them by standard norms.

**Lemma 5.5.** Under conditions (5.5) and (5.6) the following inequalities hold

$$\varepsilon_0 \|\varphi\|_{H_\rho} \leq \|\varphi\|_{H_\rho^0} \quad \forall \varphi \in V_h, \quad (5.7)$$

$$\|y\|_{C_\tau(V_\kappa)} := \max_{0 \leq m \leq M} \|y(t_m)\|_{V_\kappa} \leq \frac{\sqrt{2}}{\varepsilon_1} \|y\|_{C_\tau(H_E)} \quad \forall y \in V_\vartheta$$

with  $\varepsilon_0 := 1$  for  $\sigma \geq 1/4$  and  $\varepsilon_1 := \sqrt{4\sigma - 1}$  for  $\sigma > 1/4$ .

*Proof.* For  $\sigma \geq 1/4$ , the first inequality is obvious; for  $\sigma < 1/4$  it can be checked by a direct calculation using (5.4). The proof of the second inequality is covered in [47, Corollary 2.1].  $\square$

Now we discuss some properties of  $y_{T_h}^1$  and  $\partial_t y(T)$  that are essential below.

**Proposition 5.6.** Let  $(y_\vartheta, y_{T_h}^1) \in V_\vartheta \times V_h$  be the solution of (5.1)-(5.2). Then there holds

$$(\rho y_{T_h}^1, \varphi)_H = - \left( \kappa \partial_x \int_I y_\vartheta \, dt, \partial_x \varphi \right)_H + \int_I \langle u, \varphi \rangle_\Omega \, dt + (\rho y^1, \varphi)_H \quad \forall \varphi \in V_h. \quad (5.8)$$

*Proof.* This is proved by testing (5.1) with time constant functions  $v = \varphi \in V_h$ .  $\square$

The non-local in time identity (5.8) is convenient for our error analysis but not for the implementation; for the latter issue see Section 11. Identities similar to (5.8) also hold on the continuous level.

**Proposition 5.7.** (1) Let  $y \in \mathcal{C}(\bar{I}, V) \cap \mathcal{C}^1(\bar{I}, H)$  be the weak solution of (2.1) for  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ . Then there holds

$$(\rho \partial_t y(T), \varphi)_H = -\left(\kappa \partial_x \int_I y \, dt, \partial_x \varphi\right)_H + \left\langle \int_I u \, dt, \varphi \right\rangle_\Omega + (\rho y^1, \varphi)_H \quad \forall \varphi \in V. \quad (5.9)$$

(2) Let  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  be the weaker (very weak) solution of (2.1) for  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$ . Then there holds

$$\langle \rho \partial_t y(T), \varphi \rangle_\Omega = -\left(\kappa \partial_x \int_I y \, dt, \partial_x \varphi\right)_H + \int_I \langle u, \varphi \rangle_\Omega \, dt + \langle \rho y^1, \varphi \rangle_\Omega \quad \forall \varphi \in V. \quad (5.10)$$

*Proof.* For  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  identity (5.9) is proved by testing (3.1) with time constant function  $v = \varphi \in V$ . For  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$  we test (3.7) with any  $\varphi \in V^2$  and get

$$\langle \rho \partial_t y(T), \varphi \rangle_\Omega = ((\mathcal{I}_t y)(T), \partial_x(\kappa \partial_x \varphi))_H + \langle u, \varphi \rangle_{\mathcal{M}_T, \mathcal{C}_T}$$

According to Proposition 3.7 we have  $\mathcal{I}_t y \in \mathcal{C}(\bar{I}, V)$ . Thus there holds

$$\langle \rho \partial_t y(T), \varphi \rangle_\Omega = -(\kappa \partial_x (\mathcal{I}_t y)(T), \partial_x \varphi)_H + \langle u, \varphi \rangle_{\mathcal{M}_T, \mathcal{C}_T} + \langle \rho y^1, \varphi \rangle_\Omega.$$

The density of  $V^2$  in  $V$  implies (5.10).  $\square$

For our analysis, we need some projection and interpolation operators. We introduce the standard projectors  $\pi_h^0: H_\rho \rightarrow V_h$  and  $\pi_h^1: \mathcal{V}_\kappa \rightarrow V_h$  defined by

$$(\rho \pi_h^0 w, \varphi)_H = (\rho w, \varphi)_H \quad \forall \varphi \in V_h, \quad (5.11)$$

$$(\kappa \partial_x \pi_h^1 w, \partial_x \varphi)_H = (\kappa \partial_x w, \partial_x \varphi)_H \quad \forall \varphi \in V_h. \quad (5.12)$$

Clearly  $\|\pi_h^0 w\|_{H_\rho} \leq \|w\|_{H_\rho}$  and  $\|\pi_h^1 w\|_{\mathcal{V}_\kappa} \leq \|w\|_{\mathcal{V}_\kappa}$ . Identity (5.2) means that  $y_\vartheta(0) = \pi_h^0 y^0$ .

Moreover the following property holds

$$(w, \tilde{w})_{\mathcal{V}_\kappa} - (\pi_h^1 w, \pi_h^1 \tilde{w})_{\mathcal{V}_\kappa} = (w - \pi_h^1 w, \tilde{w} - \pi_h^1 \tilde{w})_{\mathcal{V}_\kappa} \quad \forall w, \tilde{w} \in V. \quad (5.13)$$

Following [47], we also introduce the regularized  $H_\rho$  projector  $\pi_{h, \sigma_0}^0: V \rightarrow V_h$  defined by

$$(\rho \pi_{h, \sigma_0} w, \varphi)_H + \sigma_0 \tau^2 (\kappa \partial_x \pi_{h, \sigma_0} w, \partial_x \varphi)_H = (\rho w, \varphi)_H \quad \forall \varphi \in V_h. \quad (5.14)$$

with the grid independent parameter  $\sigma_0 \geq \sigma - 1/4$ . Clearly  $\pi_{h, \sigma_0} = \pi_h^0$  for  $\sigma_0 = 0$ .

Let  $i_\tau: \mathcal{C}(\bar{I}) \rightarrow V_\tau$  be the interpolation operator such that  $i_\tau w(t_m) = w(t_m)$  for all  $m = 0, \dots, M$ .

Next we define the operator  $A_h^{-1}: V^* \rightarrow V_h$ ,  $f \mapsto w_h$  where  $w_h \in V_h$  is the unique solution of

$$(\kappa \partial_x w_h, \partial_x \varphi)_H = \langle f, \varphi \rangle_\Omega \quad \forall \varphi \in V_h. \quad (5.15)$$

Clearly  $A_h^{-1} = \pi_h^1 A^{-1}$ , see (4.3) with  $w = A^{-1} f$ , and the norm in  $\mathcal{V}_\kappa^*$  and its discrete counterpart can be written as

$$\|f\|_{\mathcal{V}_\kappa^*} = \|A^{-1} f\|_{\mathcal{V}_\kappa} = \|w\|_{\mathcal{V}_\kappa}, \quad \|f\|_{\mathcal{V}_{\kappa, h}^*} := \|A_h^{-1} f\|_{\mathcal{V}_\kappa} = \|w_h\|_{\mathcal{V}_\kappa} \leq \|w\|_{\mathcal{V}_\kappa}.$$

Moreover, we set  $r_h A^{-1} := A^{-1} - A_h^{-1} = A^{-1} - \pi_h^1 A^{-1}$ . First we note that

$$A^{-1}: H^{(\lambda)} \rightarrow H^{(\lambda+2)}, \quad -1 \leq \lambda \leq 1. \quad (5.16)$$

Then by the standard FEM error analysis [8] and operator interpolation theory we have

$$\|r_h A^{-1} f\|_V = \|w - \pi_h^1 w\|_V \leq ch^{1+\lambda} \|f\|_{H^{(\lambda)}} \quad \forall f \in H^{(\lambda)}, \quad -1 \leq \lambda \leq 0, \quad (5.17)$$

$$\|r_h A^{-1} f\|_H = \|w - \pi_h^1 w\|_H \leq ch^{2+\lambda} \|f\|_{H^{(\lambda)}} \quad \forall f \in H^{(\lambda)}, \quad -1 \leq \lambda \leq 0. \quad (5.18)$$

## 6. STABILITY AND ERROR ESTIMATES FOR THE DISCRETE STATE EQUATION

In this section we present error estimates for the state equation. We begin with an auxiliary result.

**Lemma 6.1.** *For  $\sigma_0 \geq \sigma - 1/4 \geq 0$ , the following estimate holds*

$$\|\pi_{h,\sigma_0} w - \pi_h^0 w\|_{H_\tau^0} \leq c(\tau + h)^\lambda \|w\|_{H^{(\lambda)}} \quad \forall w \in H^{(\lambda)}, \quad \text{for } 1 \leq \lambda \leq 2. \quad (6.1)$$

*Proof.* We recall the well known estimates

$$\|\pi_h^0 w\|_V \leq c \|w\|_V \quad \forall w \in V, \quad (6.2)$$

$$\|w - \pi_h^0 w\|_V \leq ch \|w\|_{V^2} \quad \forall w \in V^2, \quad (6.3)$$

which are valid using the inverse inequality (5.4). We also remind inequality (5.7) and notice also that for  $\sigma_0 \geq 0$  the following additional inequality holds

$$\sqrt{\sigma_0 \tau} \|\varphi\|_{\mathcal{V}_\kappa} \leq \|\varphi\|_{H_\tau^0} \quad \forall \varphi \in V_h. \quad (6.4)$$

Let  $w \in V$  and  $\varphi \in V_h$ . We apply identities (5.11) and (5.14) and get

$$\begin{aligned} (\rho(\pi_{h,\sigma_0}^0 w - \pi_h^0 w), \varphi)_H + \sigma_0 \tau^2 (\kappa \partial_x (\pi_{h,\sigma_0}^0 w - \pi_h^0 w), \partial_x \varphi)_H &= -\sigma_0 \tau^2 (\kappa \partial_x \pi_h^0 w, \partial_x \varphi)_H \\ &= \sigma_0 \tau^2 (\kappa \partial_x (w - \pi_h^0 w), \partial_x \varphi)_H + \sigma_0 \tau^2 \langle \partial_x (\kappa \partial_x w), \varphi \rangle_\Omega. \end{aligned}$$

Now we set  $\varphi = \pi_{h,\sigma_0}^0 w - \pi_h^0 w$ , and from the former equality and estimate (6.2) for  $\lambda = 1$  as well as the latter equality and estimate (6.3) for  $\lambda = 2$  we obtain the estimate

$$\|\pi_{h,\sigma_0}^0 w - \pi_h^0 w\|_{H_\tau^0} \leq c\tau(\tau + h)^{\lambda-1} \|w\|_{H^{(\lambda)}} \quad \text{for } \lambda = 1, 2.$$

By using the  $K_{\lambda,\infty}$ -method, we complete the proof.  $\square$

Now we get a stability bound and error estimates in  $\mathcal{C}(\bar{I}, H) \times \mathcal{V}_{\kappa,h}^*$  for the discrete state equation.

**Proposition 6.2.** *Let  $y$  and  $(y_\vartheta, y_{Th}^1)$  be the solutions to the state equation (2.1) and the discrete state equation (5.1)-(5.2).*

(1) *For  $(u, y^0, y^1) \in L^2(I, V^*) \times V \times V^*$ , the following stability bound holds:*

$$\|y_\vartheta\|_{\mathcal{C}(\bar{I}, H)} + \|\rho y_{Th}^1\|_{\mathcal{V}_{\kappa,h}^*} \leq c (\|u\|_{L^2(I, V^*)} + \|y\|_{V \times V^*}). \quad (6.5)$$

(2) *For  $(u, y^0, y^1) \in L^2(I, H^{(-1/2)}) \times V \times H^{(-1/2)}$ , the following error estimate holds:*

$$\|y - y_\vartheta\|_{\mathcal{C}(\bar{I}, H)} + \|\rho(\partial_t y(T) - y_{Th}^1)\|_{\mathcal{V}_{\kappa,h}^*} \leq c(\tau + h)^{1/3} (\|u\|_{L^2(I, H^{(-1/2)})} + \|y\|_{V \times H^{(-1/2)}}). \quad (6.6)$$

(3) *For  $(u, y^0, y^1) \in H^1(I, H^{(-1/2)}) \times V \times H$ , the higher order error estimate holds:*

$$\|y - y_\vartheta\|_{\mathcal{C}(\bar{I}, H)} + \|\rho(\partial_t y(T) - y_{Th}^1)\|_{\mathcal{V}_{\kappa,h}^*} \leq c(\tau + h)^{2/3} (\|u\|_{H^1(I, H^{(-1/2)})} + \|y\|_{V \times H}). \quad (6.7)$$

*Proof.* 1. According to [47, Theorem 2.1 (1)], the bound

$$\|y_\vartheta\|_{\mathcal{C}(\bar{I}, H)} + \left\| \int_I y_\vartheta \, dt \right\|_V \leq c (\|u\|_{L^2(I, V^*)} + \|y_\vartheta(0)\|_{H_\tau^0} + \|y^1\|_{V^*}) \quad (6.8)$$

is valid for any  $y_\vartheta(0) \in V_h$ . We have  $y_\vartheta(0) = \pi_h^0 y^0$ . In the case  $\sigma \leq 1/4$ , there clearly holds

$$\|\pi_h^0 y^0\|_{H_\tau^0} \leq \|\pi_h^0 y^0\|_{H_\rho} \leq \|y^0\|_{H_\rho}.$$

For  $\sigma > 1/4$ , we alternatively get using (6.1) for  $\lambda = 1$

$$\|\pi_h^0 y^0\|_{H_\tau^0} \leq \|\pi_h^0 y^0 - \pi_{h,\sigma_0}^0 y^0\|_{H_\tau^0} + \|\pi_{h,\sigma_0}^0 y^0\|_{H_\tau^0} \leq c(\tau + h) \|y^0\|_V + \|y^0\|_{H_\rho}$$

for any  $\sigma_0 \geq \sigma - 1/4$ .

We proceed with the bound for  $y_{Th}^1$ . Identity (5.8) and bound (6.8) together with the generalized Minkowski inequality imply

$$\|\rho y_{Th}^1\|_{\mathcal{V}_{\kappa,h}^*} \leq c \left( \left\| \int_I y_\vartheta \, dt \right\|_V + \left\| \int_I u \, dt \right\|_{V^*} + \|\rho y^1\|_{V^*} \right) \leq c_1 (\|u\|_{L^2(I, V^*)} + \|y\|_{V \times H}). \quad (6.9)$$

Finally we derive bound (6.5).

2. Let  $\tilde{y}_\vartheta$  be the solution of equation (5.1) for  $\tilde{y}_\vartheta(0) = \pi_{h,\sigma_0}^0 y^0$ . From [47, Theorem 4.1] we get the error estimate

$$\|y - \tilde{y}_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - \tilde{y}_\vartheta) dt \right\|_V \leq c(\tau + h)^{1/3} (\|u\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{y}\|_{H^{(1/2)} \times H^{(-1/2)}}).$$

In the case  $\sigma \leq 1/4$  we can choose  $\sigma_0 = 0$ , then  $y_\vartheta(0) = \pi_{h,\sigma_0} y^0 = \pi_h^0 y^0$  and  $\tilde{y}_\vartheta = y_\vartheta$ . In the case  $\sigma \geq 1/4$  we can use the stability bound (6.8) and estimate (6.1) to get

$$\|\tilde{y}_\vartheta - y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\tilde{y}_\vartheta - y_\vartheta) dt \right\|_V \leq c \|\pi_{h,\sigma_0}^0 y^0 - \pi_h^0 y^0\|_{H_\tau^0} \leq c_1(\tau + h) \|y^0\|_V. \quad (6.10)$$

Then by subtracting (5.8) from (5.10) and applying identity (5.12) we find

$$\langle \rho(\partial_t y(T) - y_{T_h}^1), \varphi \rangle_\Omega = - \left( \kappa \partial_x \int_I (y - y_\vartheta) dt, \partial_x \varphi \right)_H = - \left( \kappa \partial_x \int_I (\pi_h^1 y - y_\vartheta) dt, \partial_x \varphi \right)_H \quad \forall \varphi \in V_h,$$

consequently

$$\|\rho(\partial_t y(T) - y_{T_h}^1)\|_{V_{\kappa,h}^*} \leq c \left\| \int_I (\pi_h^1 y - y_\vartheta) dt \right\|_V. \quad (6.11)$$

Thus we obtain (6.6).

3. Once again we apply [47, Theorem 4.1] and first get the estimate

$$\|y - \tilde{y}_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - \tilde{y}_\vartheta) dt \right\|_V \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I,H)} + \|\mathbf{y}\|_{V \times H}).$$

Combining it together with (6.10), we derive

$$\|y - y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - y_\vartheta) dt \right\|_V \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I,H)} + \|\mathbf{y}\|_{V \times H}). \quad (6.12)$$

In this proof, we apply this estimate in the case  $u = 0$  only (but in general case below).

In the remaining case  $\mathbf{y} = 0$ , from [47, Theorem 4.1] we also get the higher order error estimate

$$\|y - y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - y_\vartheta) dt \right\|_V \leq c(\tau + h)^{4/3} \|u\|_{H^1(I,H)} \quad \text{for } \forall u \in H^1(I,H). \quad (6.13)$$

Moreover owing to Proposition 3.7 and bound (6.5) (both for  $\mathbf{y} = 0$ ) we have

$$\begin{aligned} \|y - y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - y_\vartheta) dt \right\|_V &\leq \|y\|_{C(\bar{I},H)} + c \|\mathcal{I}t y\|_{C(\bar{I},V)} + \|y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I y_\vartheta dt \right\|_V \\ &\leq c_1 \|u\|_{L^2(I,V^*)} \quad \text{for } \forall u \in L^2(I,V^*). \end{aligned}$$

The last bound and estimate (6.13) imply by the  $K_{1/2,\infty}$ -method:

$$\|y - y_\vartheta\|_{C(\bar{I},H)} + \left\| \int_I (\pi_h^1 y - y_\vartheta) dt \right\|_V \leq c(\tau + h)^{2/3} \|u\|_{B_{1/2}} \quad \forall u \in B_{1/2} := (L^2(I,V^*), H^1(I,H))_{1/2,\infty}.$$

Due to the simple embedding

$$H^1(I, H^{(-1/2)}) = (H^1(I; V^*), H^1(I; H))_{1/2,\infty} \hookrightarrow (L^2(I, V^*), H^1(I, H))_{1/2,\infty}$$

and inequality (6.11) we complete the proof.  $\square$

**Remark 6.3.** A priori stability bound (6.5) implies the unique solvability of the discrete state equation (5.1)-(5.2).

**Remark 6.4.** According to the given proof, for  $\tilde{y}_\vartheta$  in place of  $y_\vartheta$  the norms of  $\mathbf{y}$  in (6.5) and (6.6) can be weakened down to respectively  $\|\mathbf{y}\|_{H \times V^*}$  and  $\|\mathbf{y}\|_{H^{(1/2)} \times H^{(-1/2)}}$ . For  $\sigma \leq 1/4$ , we have  $\tilde{y}_\vartheta = y_\vartheta$ . The same can be shown for  $y_\vartheta$  also for  $\sigma > 1/4$  provided that  $\tau \alpha_h \leq c_0$  with any  $c_0 > 0$ .



## 7. DISCRETE CONTROL PROBLEM

First we introduce the discrete mapping  $\hat{S}_\vartheta: (u, y_0, y_1) \mapsto (y_\vartheta, y_\vartheta(T), \rho y_{T_h}^1)$  and the discrete affine linear control-to-state mapping

$$S_\vartheta: \mathcal{M}_T \rightarrow \mathcal{Y}_\vartheta = V_\vartheta \times V_h \times (\rho \times V_h), \quad u \mapsto (y_\vartheta, y_\vartheta(T), \rho y_{T_h}^1)$$

defined by  $S_\vartheta u = \hat{S}_\vartheta(u, 0, 0) + \hat{S}_\vartheta(0, y_0, y_1)$ , with  $\rho \times V_h = \{\rho\varphi; \varphi \in V_h\}$ . The mapping  $S_\vartheta$  is a composition of

$$u \mapsto \vec{u} = \{\langle u, e_{m,n}^\vartheta \rangle_{\mathcal{M}_T, \mathcal{C}_T}\}_{m,n=1}^{M, N-1}, \quad \mathcal{M}_T \rightarrow \mathbb{R}^{M(N-1)},$$

where  $\{e_{m,n}^\vartheta\}$  is a basis in  $V_\vartheta$ , and  $\vec{u} \mapsto (y_\vartheta, y_\vartheta(T), \rho y_{T_h}^1)$ . The former mapping is bounded due to  $e_{m,n}^\vartheta \in \mathcal{C}_T$  and the latter one is finite dimensional. Thus  $S_\vartheta$  is a bounded operator. Then we consider the following semi-discrete optimal control problem

$$j_\vartheta(u) = \frac{1}{2} \|S_\vartheta u - \mathbf{z}\|_{\mathcal{Y}_h}^2 + \alpha \|u\|_{\mathcal{M}_T} \rightarrow \min_{u \in \mathcal{M}_T} \quad (\mathcal{P}_\vartheta)$$

with the squared semi-norm corresponding to the inner product

$$(\mathbf{z}, \tilde{\mathbf{z}})_{\mathcal{Y}_h} = (\rho z_1, \tilde{z}_1)_{L^2(I \times \Omega)} + (\rho z_2, \tilde{z}_2)_H + (A_h^{-1} z_3, A_h^{-1} \tilde{z}_3)_{\mathcal{V}_k} \quad \forall \mathbf{z}, \tilde{\mathbf{z}} \in \mathcal{Y}.$$

Using the similar argument as in the continuous case it can be shown that  $(\mathcal{P}_\vartheta)$  has a solution  $\bar{u}_\vartheta$  which is not unique in general, and due to the optimality, the stability bound (6.5) and property (5.16) (for  $\lambda = -1$ ) one gets

$$\alpha \|\bar{u}_\vartheta\|_{\mathcal{M}_T} \leq j_\vartheta(\bar{u}_\vartheta) \leq j_\vartheta(0) = \frac{1}{2} \|S_\vartheta(0) - \mathbf{z}\|_{\mathcal{Y}_h}^2 \leq c (\|\mathbf{y}\|_{V \times V^*} + \|z\|_{\mathcal{Y}})^2,$$

cf. (4.1), and consequently

$$\|\bar{u}_\vartheta\|_{\mathcal{M}_T} \leq c (\|\mathbf{y}\|_{V \times V^*} + \|z\|_{\mathcal{Y}})^2 \leq C. \quad (7.1)$$

**Theorem 7.1.** *Let  $\mathbf{z} \in \mathcal{Y}$ ,  $\mathbf{y} \in V \times H^{(-1/2)}$  and  $\bar{u}, \bar{u}_\vartheta \in \mathcal{M}_T$  be the optimal controls of respectively problems  $(\mathcal{P})$  and  $(\mathcal{P}_\vartheta)$ . Then there holds*

$$\bar{u}_\vartheta \rightharpoonup^* \bar{u} \text{ in } \mathcal{M}_T, \quad \|\bar{u}_\vartheta\|_{\mathcal{M}_T} \rightarrow \|\bar{u}\|_{\mathcal{M}_T} \text{ as } \vartheta \rightarrow 0.$$

*Proof.* Owing to (7.1) there exists a sequence  $\{\vartheta_n\}$ ,  $\vartheta_n \rightarrow 0$ , and  $u \in \mathcal{M}_T$  such that  $\bar{u}_{\vartheta_n} \rightharpoonup^* u$  in  $\mathcal{M}_T$  as  $n \rightarrow \infty$ . Next we prove that this implies that

$$\|S_{\vartheta_n} \bar{u}_{\vartheta_n} - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} \rightarrow \|Su - \mathbf{z}\|_{\mathcal{Y}}. \quad (7.2)$$

To this end, we write the chain of inequalities

$$\begin{aligned} \left| \|S_{\vartheta_n} \bar{u}_{\vartheta_n} - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} - \|Su - \mathbf{z}\|_{\mathcal{Y}} \right| &\leq \left| \|S_{\vartheta_n} \bar{u}_{\vartheta_n} - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} - \|Su - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} \right| + \left| \|Su - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} - \|Su - \mathbf{z}\|_{\mathcal{Y}} \right| \\ &\leq \|S_{\vartheta_n} \bar{u}_{\vartheta_n} - Su\|_{\mathcal{Y}_{\vartheta_n}} + \left| \|Su - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} - \|Su - \mathbf{z}\|_{\mathcal{Y}} \right| \\ &\leq \|S_{\vartheta_n} \bar{u}_{\vartheta_n} - S\bar{u}_{\vartheta_n}\|_{\mathcal{Y}_{\vartheta_n}} + \|S\bar{u}_{\vartheta_n} - Su\|_{\mathcal{Y}_{\vartheta_n}} + \left| \|Su - \mathbf{z}\|_{\mathcal{Y}_{\vartheta_n}} - \|Su - \mathbf{z}\|_{\mathcal{Y}} \right|. \end{aligned}$$

The first term on the right in the last inequality converges to zero according to the error estimate (6.6). The convergence of the second term follows from the weak-star-to-strong continuity of  $S: \mathcal{M}_T \rightarrow \mathcal{Y}$  and the stability of  $\pi_h^1$  in  $V$ . Finally, property (5.13) for  $\tilde{w} = w$  implies the convergence of the last term. Then (7.2) and the weak-star lower semicontinuity of  $\|\cdot\|_{\mathcal{M}_T}$  in  $\mathcal{M}_T$  implies

$$j(u) \leq \liminf_{n \rightarrow \infty} j_{\vartheta_n}(\bar{u}_{\vartheta_n}) \leq \limsup_{n \rightarrow \infty} j_{\vartheta_n}(\bar{u}_{\vartheta_n}) \leq \limsup_{n \rightarrow \infty} j_{\vartheta_n}(\bar{u}) = j(\bar{u}).$$

Thus, the uniqueness of  $\bar{u}$  means that  $u = \bar{u}$  and in addition implies the convergence of the whole sequence  $\bar{u}_\vartheta \rightharpoonup^* \bar{u}$  in  $\mathcal{M}_T$  as  $\vartheta \rightarrow 0$ . Moreover, we have  $j_\vartheta(\bar{u}_\vartheta) \rightarrow j(\bar{u})$ . This and (7.2) lead to  $\|\bar{u}_\vartheta\|_{\mathcal{M}_T} \rightarrow \|\bar{u}\|_{\mathcal{M}_T}$ .  $\square$

For convenience we set  $F_h(\mathbf{z}) = (1/2)\|\mathbf{z}\|_{\mathcal{Y}_h}^2$ . In the following the directional derivative of a functional  $g: \mathcal{M}_T \rightarrow \mathbb{R}$  at  $u \in \mathcal{M}_T$  in direction  $\delta u \in \mathcal{M}_T$  is denoted by  $Dg(u)\delta u$ . In the case  $Dg(u) \in \mathcal{M}_T^*$ ,  $g$  is the Gateaux differentiable in  $u$ . Moreover, we make use of the convex subdifferential of  $\|\cdot\|_{\mathcal{M}_T}$ . Let  $\hat{u} \in \mathcal{M}_T$  and  $p \in \mathcal{C}_T$ . Then there holds  $p \in \partial\|\hat{u}\|_{\mathcal{M}_T}$  if and only if

$$\langle p, u - \hat{u} \rangle_{\mathcal{C}_T, \mathcal{M}_T} + \alpha\|\hat{u}\|_{\mathcal{M}_T} \leq \alpha\|u\|_{\mathcal{M}_T} \quad \forall u \in \mathcal{M}_T.$$

An element  $\bar{u}_\vartheta \in \mathcal{M}_T$  is an optimal solution of  $(\mathcal{P}_\vartheta)$  if and only if  $-D((F_h \circ S_\vartheta)(\bar{u}_\vartheta)) \in \alpha\partial\|\bar{u}_\vartheta\|_{\mathcal{M}_T}$ . To calculate  $D((F_h \circ S_\vartheta)(u))$  for  $u \in \mathcal{M}_T$ , we apply the Lagrange technique and define the Lagrange functional by

$$\begin{aligned} L(u, y_\vartheta, y_{T_h}^1, p_\vartheta, p_{0h}^1) &= F_h(y_\vartheta, y_\vartheta(T), y_{T_h}^1) - B_\sigma(y_\vartheta, p_\vartheta) - (\rho y_{T_h}^1, p_\vartheta(T))_H \\ &\quad + \langle u, p_\vartheta \rangle_{\mathcal{M}_T, \mathcal{C}_T} + (\rho y^1, p_\vartheta(0))_H + (\rho(y_\vartheta(0) - y^0), p_{0h}^1)_H \end{aligned}$$

with  $(p_\vartheta, p_{0h}^1) \in V_\vartheta \times V_h$  (where we base on identities (5.1)-(5.2)). We obviously have

$$(F_h \circ S_\vartheta)(u) = L(u, S_\vartheta u, p_\vartheta, p_{0h}^1) \quad \forall (p_\vartheta, p_{0h}^1) \in V_\vartheta \times V_h.$$

Thus there holds

$$D((F_h \circ S_\vartheta)(u))\delta u = D_u L(u, y_\vartheta, y_{T_h}^1, p_\vartheta, p_{0h}^1)\delta u = \langle p_\vartheta, \delta u \rangle_{\mathcal{C}_T, \mathcal{M}_T} \quad \forall \delta u \in \mathcal{M}_T$$

provided that  $(p_\vartheta, p_{0h}^1) \in V_\vartheta \times V_h$  is the solution of the discrete problem

$$\begin{aligned} -D_{y_\vartheta} L(u, y_\vartheta, y_{T_h}^1, p_\vartheta, p_{0h}^1)v &= B_\sigma(v, p_\vartheta) - (\rho(y_\vartheta - z_1), v)_{L^2(I \times \Omega)} \\ &\quad - (\rho(y_\vartheta(T) - z_2), v(T))_H - (\rho v(0), p_{0h}^1)_H = 0 \quad \forall v \in V_\vartheta \end{aligned}$$

and

$$-D_{y_{T_h}^1} L(u, y_\vartheta, y_{T_h}^1, p_\vartheta, p_{0h}^1)\varphi = (\rho\varphi, p_\vartheta(T))_H - (\rho A_h^{-1}(\rho y_{T_h}^1 - z_3), \varphi)_H = 0 \quad \forall \varphi \in V_h.$$

Therefore the discrete optimality system consists of the discrete state equation

$$\begin{aligned} B_\sigma(\bar{y}_\vartheta, v) + (\rho \bar{y}_{T_h}^1, v(T))_H &= \langle \bar{u}_\vartheta, v \rangle_{\mathcal{M}_T, \mathcal{C}_T} + (\rho y^1, v(0))_H \quad \forall v \in V_\vartheta, \\ (\rho \bar{y}_\vartheta(0), \varphi)_H &= (\rho y^0, \varphi)_H \quad \forall \varphi \in V_h, \end{aligned} \quad (7.3)$$

the discrete adjoint state equation

$$\begin{aligned} B_\sigma(v, \bar{p}_\vartheta) - (\rho v(0), \bar{p}_{0h}^1)_H &= (\rho(\bar{y}_\vartheta - z_1), v)_{L^2(I \times \Omega)} + (\rho(\bar{y}_\vartheta(T) - z_2), v(T))_H \quad \forall v \in V_\vartheta, \\ (\rho\varphi, \bar{p}_\vartheta(T))_H &= (\rho A_h^{-1}(\rho \bar{y}_{T_h}^1 - z_3), \varphi)_H \quad \forall \varphi \in V_h \end{aligned} \quad (7.4)$$

and the discrete variational inequality

$$\langle -\bar{p}_\vartheta, u - \bar{u}_\vartheta \rangle_{\mathcal{C}_T, \mathcal{M}_T} + \|\bar{u}_\vartheta\|_{\mathcal{M}_T} \leq \|u\|_{\mathcal{M}_T} \quad \forall u \in \mathcal{M}_T. \quad (7.5)$$

## 8. STABILITY AND ERROR ESTIMATES FOR THE DISCRETE ADJOINT STATE EQUATION

We define the general discrete adjoint state equation

$$B_\sigma(v, p_\vartheta) - (\rho v(0), p_{0h}^1)_H = (\rho(y - z_1), v)_{L^2(I \times \Omega)} + (\rho(y(T) - z_2), v(T))_H \quad \forall v \in V_\vartheta, \quad (8.1)$$

$$(\rho\varphi, p_\vartheta(T))_H = (\rho A_h^{-1}(\rho \partial_t y(T) - z_3), \varphi)_H \quad \forall \varphi \in V_h. \quad (8.2)$$

Here  $y$  is the solution to the state equation (2.1). Clearly identity (8.2) means simply that  $p_\vartheta(T) = A_h^{-1}q_T = \pi_1^h A^{-1}q_T$  with  $q_T := \rho \partial_t y(T) - z_3$ .

Now we get a stability bound and error estimates in  $\mathcal{C}(\bar{I}, H) \times \mathcal{V}_{\kappa, h}^*$  and  $\mathcal{C}_T$  for the discrete adjoint state equation.

**Proposition 8.1.** *Let  $p = S^*(y - z_1, -(y(T) - z_2), A^{-1}(\rho \partial_t y(T) - z_3))$  and  $(p_\vartheta, p_{0h}^1)$  be the solution of the corresponding general discrete adjoint state equation (8.1)-(8.2).*

(1) *If  $y \in \mathcal{C}(\bar{I}, H) \cap \mathcal{C}^1(\bar{I}, V^*)$  and  $\mathbf{z} \in \mathcal{Y}$ , then the following stability bound holds*

$$\|p_\vartheta\|_{\mathcal{C}(\bar{I}, V)} + \|\rho p_{0h}^1\|_{\mathcal{V}_{\kappa, h}^*} \leq c(\|y - z_1\|_{L^2(I \times \Omega)} + \|y(T) - z_2\|_H + \|\rho \partial_t y(T) - z_3\|_{V^*}). \quad (8.3)$$

(2) If  $u \in L^2(I, V^*)$ ,  $\mathbf{z} \in \mathcal{Y}$  and  $\mathbf{y} \in H \times V^*$ , then the following error estimate holds

$$\|p - p_\vartheta\|_{C(\bar{I}, H)} + \|\rho(\partial_t p(0) - p_h^0)\|_{V_{\kappa, h}^*} \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I, V^*)} + \|\mathbf{z}\|_{\mathcal{Y}} + \|\mathbf{y}\|_{H \times V^*}). \quad (8.4)$$

(3) If  $u \in L^2(I, H^{(-1/2)})$ ,  $\mathbf{z} \in \mathcal{Y}^{1/2} := L^2(I, H^{(1/2)}) \times H^{(1/2)} \times H^{(-1/2)}$  and  $\mathbf{y} \in H^{(1/2)} \times H^{(-1/2)}$ , then the following error estimate holds

$$\|p - p_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{1/2}} + \|\mathbf{y}\|_{H^{(1/2)} \times H^{(-1/2)}}). \quad (8.5)$$

(4) If  $u \in H^1(I, H^{(-1/2)})$ ,  $\mathbf{z} \in \mathcal{Y}^{3/2} := L^2(I, H^{(3/2)}) \times H^{(3/2)} \times H^{(1/2)}$  and  $\mathbf{y} \in H^{(3/2)} \times H^{(1/2)}$ , then the following higher order error estimate holds

$$\|p - p_\vartheta\|_{L^2(I, C_0(\Omega))} \leq c(\tau + h)^{4/3} (\|u\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{3/2}} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}}). \quad (8.6)$$

*Proof.* 1. According to [47, Theorem 2.1 (2)] the following energy bound hold

$$\|p_\vartheta\|_{C(\bar{I}, V)} + \|\partial_t p_\vartheta\|_{L^\infty(\bar{I}, H)} \leq c (\|y - z_1\|_{L^2(I \times \Omega)} + \|p_\vartheta(T)\|_V + \|y(T) - z_2\|_H)$$

for any  $p_\vartheta(T) \in V_h$ . Using (6.2),  $A_h^{-1} = \pi_h^1 A^{-1}$  and (5.16) we get

$$\|p_\vartheta(T)\|_V \leq c \|A^{-1} q_T\|_V \leq c_1 \|q_T\|_{V^*}. \quad (8.7)$$

By applying also the counterpart of inequalities (6.9) we derive bound (8.3).

2. The counterpart of the error estimate (6.12) for the adjoint state equation case and bound (8.7) give

$$\begin{aligned} \|p - p_\vartheta\|_{C(\bar{I}, H)} + \left\| \int_I (\pi_h^1 p - p_\vartheta) dt \right\|_V &\leq c(\tau + h)^{2/3} (\|y - z_1\|_{L^2(I, H)} + \|A_h^{-1} q_T\|_V + \|y(T) - z_2\|_H) \\ &\leq c_1 (\tau + h)^{2/3} (\|y\|_{C(\bar{I}, H)} + \|\partial_t y\|_{C^1(\bar{I}, V^*)} + \|\mathbf{z}\|_{\mathcal{Y}}). \end{aligned}$$

Owing to inequality (6.11) and Proposition 3.7 we obtain estimate (8.4).

3. Below we need the multiplicative inequalities

$$\|w\|_{C(\bar{I} \times \bar{\Omega})} \leq c \|w\|_{C(\bar{I}, H)}^{1/2} \|w\|_{C(\bar{I}, V)}^{1/2} \quad \forall w \in C(\bar{I}, V), \quad (8.8)$$

$$\|w\|_{L^2(I, C_0(\Omega))} \leq c \|w\|_{L^2(I, H)}^{1/2} \|w\|_{L^2(I, V)}^{1/2} \quad \forall w \in L^2(I, V). \quad (8.9)$$

Let  $\check{p}_\vartheta$  be the auxiliary solution to (8.1) for  $\check{p}_\vartheta(T) = \pi_h^0 A^{-1} q_T$ . Owing to inequality (8.8) and the stability bounds [47, Theorem 2.1] we get

$$\|p_\vartheta - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c \|(p_\vartheta - \check{p}_\vartheta)(T)\|_{H_T^0}^{1/2} \|(p_\vartheta - \check{p}_\vartheta)(T)\|_V^{1/2}.$$

Consequently, for  $q_T \in H^{(\alpha-2)}$ , by (6.2), (5.17) and (5.18) the following chain of inequalities hold

$$\begin{aligned} \|p_\vartheta - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} &\leq c (\|(p_\vartheta - \check{p}_\vartheta)(T)\|_H^{1/2} \|(p_\vartheta - \check{p}_\vartheta)(T)\|_V^{1/2} + \tau^{1/2} \|(p_\vartheta - \check{p}_\vartheta)(T)\|_V) \\ &\leq c_1 (\|r_h A^{-1} q_T\|_H^{1/2} \|r_h A^{-1} q_T\|_V^{1/2} + \tau^{1/2} \|r_h A^{-1} q_T\|_V) \leq c_2 (\tau + h)^{\alpha-1/2} \|q_T\|_{H^{(\alpha-2)}} \end{aligned}$$

for  $1 \leq \alpha \leq 2$ . Thus it is enough to prove error estimates (8.5) and (8.6) for  $\check{p}_\vartheta$  instead of  $p_\vartheta$ .

According to [47, Theorem 5.3 and estimate (5.18)] we have the error estimate

$$\begin{aligned} \|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} &= \|p - \check{p}_\vartheta\|_{C_\tau(\bar{I}, C(\bar{\Omega}))} := \max_{0 \leq m \leq M} \|(p - \check{p}_\vartheta)(t_m)\|_{C(\bar{\Omega})} \\ &\leq c(\tau + h)^{2(\alpha-1/2)/3} (\|y - z_1\|_{L^2(I, H^{(\alpha-1)})} + \|y(T) - z_2\|_{H^{(\alpha-1)}} + \|q_T\|_{H^{(\alpha-2)}}), \quad \alpha = 1, 2. \end{aligned} \quad (8.10)$$

We emphasize that due to [47, Theorem 4.3 (2) (e)] and (6.1) this estimate holds for  $\check{p}_\vartheta(T) = \pi_h^0 A^{-1} q_T$ .

Inequality (8.8), Proposition 3.5 (applied to the adjoint state problem) and property (5.16) imply the following error estimate for the time interpolation

$$\begin{aligned} \|p - i_\tau p\|_{C(\bar{I} \times \bar{\Omega})} &\leq c(\tau \|\partial_t p\|_{C(\bar{I}, H)})^{1/2} \|(\tau \partial_t)^{\alpha-1} p\|_{C(\bar{I}, V)}^{1/2} \\ &\leq c_1 \tau^{\alpha/2} (\|y - z_1\|_{L^2(I, H^{(\alpha-1)})} + \|y(T) - z_2\|_{H^{(\alpha-1)}} + \|q_T\|_{H^{(\alpha-2)}}), \end{aligned} \quad (8.11)$$

for  $\alpha = 1, 2$ . Owing to estimates (8.10) and (8.11) as well as Propositions 3.7 and 3.5 we get

$$\begin{aligned} \|p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} &\leq c(\tau + h)^{2(\alpha-1/2)/3} (\|y\|_{C(\bar{I}, H^{(\alpha-1)})} + \|\partial_t y\|_{C(\bar{I}, H^{(\alpha-2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{(\alpha-1)}}) \\ &\leq c_1(\tau + h)^{2(\alpha-1/2)/3} (\|u\|_{L^2(I, H^{(\alpha-2)})} + \|\mathbf{y}\|_{H^{(\alpha-1)} \times H^{(\alpha-2)}} + \|\mathbf{z}\|_{\mathcal{Y}^{(\alpha-1)}}), \quad \alpha = 1, 2, \end{aligned} \quad (8.12)$$

where  $\mathcal{Y}^{(\alpha)} := L^2(I, H^{(\alpha)}) \times H^{(\alpha)} \times H^{(\alpha-1)}$ .

By applying the  $K_{1/2, \infty}$ -method, we get (8.5) for  $\check{p}_\vartheta$  in the role of  $p_\vartheta$ .

4. First notice that the multiplicative inequality (8.9), Proposition 3.5 (2) (applied for the adjoint state problem) and property (5.16) imply another error estimate for the time interpolation

$$\begin{aligned} \|p - i_\tau p\|_{L^2(I, C_0(\Omega))} &\leq c(\tau^2 \|\partial_{tt} p\|_{L^2(I, H)})^{1/2} (\tau \|\partial_t p\|_{C(\bar{I}, V)})^{1/2} \\ &\leq c_1 \tau^{3/2} (\|y - z_1\|_{L^2(I, V)} + \|y(T) - z_2\|_V + \|q_T\|_H). \end{aligned}$$

Then Proposition 3.5 (1) leads to

$$\|p - i_\tau p\|_{L^2(I, C_0(\Omega))} \leq c\tau^{3/2} (\|u\|_{H^1(I, V^*)} + \|z\|_{\mathcal{Y}} + \|\mathbf{y}\|_{V \times H}). \quad (8.13)$$

Next we derive the error estimate

$$\|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h)^{4/3} (\|u\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{3/2}} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}}). \quad (8.14)$$

According to [47, Theorem 5.3 and estimate (5.18)] and Propositions 3.7 and 3.5 the following three estimates hold

$$\|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h)^{4/3} \|\mathbf{z}\|_{\mathcal{Y}^{3/2}} \quad \text{for } u = 0, \mathbf{y} = 0, \quad (8.15)$$

$$\|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h) \|y\|_{H^1(I, H)} \leq c_1(\tau + h) (\|u\|_{H^1(I, V^*)} + \|\mathbf{y}\|_{V \times H}) \quad \text{for } \mathbf{z} = 0,$$

$$\begin{aligned} \|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} &\leq c(\tau + h)^{5/3} (\|\partial_{tt} y\|_{L^2(I, H)} + \|\mathbf{y}\|_{V \times H}) \\ &\leq c_1(\tau + h)^{5/3} (\|u\|_{H^1(I, H)} + \|\mathbf{y}\|_{V^2 \times V}) \quad \text{for } \mathbf{z} = 0 \end{aligned}$$

for  $\check{p}_\vartheta(T) = \pi_h^0 A^{-1} q_T$  (for the same reason as above). Then applying the  $K_{1/2, \infty}$ -method to the two last estimates we get

$$\|i_\tau p - \check{p}_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h)^{4/3} (\|u\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}}) \quad \text{for } \mathbf{z} = 0.$$

By combining this estimate and (8.15) we obtain (8.14).

Estimates (8.13) and (8.14) imply

$$\|p - \check{p}_\vartheta\|_{L^2(I, C_0(\Omega))} \leq c(\tau + h)^{4/3} (\|u\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{3/2}} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}})$$

that completes the proof of (8.6) for  $\check{p}_\vartheta$  in the role of  $p_\vartheta$ .  $\square$

**Remark 8.2.** A priori stability bound (6.5) (taken for  $y = 0$ ) implies the unique solvability of the general discrete adjoint state equation (8.1)-(8.2).

## 9. ERROR ESTIMATES FOR THE STATE VARIABLE

We introduce the discrete adjoint control-to-state operator  $S_\vartheta^*: L^2(I \times \Omega) \times V \times H \rightarrow V_\vartheta$ ,  $(\phi, p^1, p^0) \mapsto p_\vartheta$  defined by

$$B_\sigma(v, p_\vartheta) = (\rho\phi, v)_{L^2(I \times \Omega)} - (\rho p^1, v(T))_H \quad \forall v \in V_\vartheta, v(0) = 0$$

with  $p_\vartheta(T) = \pi_h^0 p^0$ . Similarly to bound (8.3) and Remark 8.2 it is well defined and satisfies

$$\|S_\vartheta^*(\phi, p_0, p_1)\|_{C(\bar{I}, V)} \leq c(\|\phi\|_{L^2(I \times \Omega)} + \|p^0\|_V + \|p^1\|_H).$$

Let for brevity  $W, W_h: \mathcal{Y} \rightarrow \mathcal{Y}^*$  be the duality mappings defined by

$$W(y_1, y_2, y_3) = (y_1, -y_2, A^{-1}y_3), \quad W_h(y_1, y_2, y_3) = (y_1, -y_2, A_h^{-1}y_3) \quad \forall (y_1, y_2, y_3) \in \mathcal{Y}.$$

With this notation, the function  $p_\vartheta = S_\vartheta^*(y - z_1, -(y(T) - z_2), A_h^{-1}(\rho\partial_t y(T) - z_3)) = S_\vartheta^*W_h(Su - z)$  solves the general discrete adjoint state equation (8.1)-(8.2).

**Proposition 9.1.** *Let  $\mathbf{z} \in \mathcal{Y}$  and  $\mathbf{y} \in V \times V^*$ . Then the following estimate holds*

$$\|S\bar{u} - S_\vartheta\bar{u}_\vartheta\|_{\mathcal{Y}_h} \leq \|S\bar{u} - S_\vartheta\bar{u}\|_{\mathcal{Y}_h} + C\|S^*W(S\bar{u} - \mathbf{z}) - S_\vartheta^*W_h(S\bar{u} - \mathbf{z})\|_{C_T}^{1/2}. \quad (9.1)$$

*Proof.* We recall that  $\bar{p} = S^*W(S\bar{u} - \mathbf{z})$  and  $\bar{p}_\vartheta = S_\vartheta^*W_h(S_\vartheta\bar{u}_\vartheta - \mathbf{z})$  and test the continuous subgradient condition (4.5) with the discrete optimal control  $\bar{u}_\vartheta$  and the discrete subgradient condition (7.5) with the continuous optimal control  $\bar{u}$ . Then we subtract the first inequality from the second one and get

$$\langle \bar{u} - \bar{u}_\vartheta, \bar{p} - \bar{p}_\vartheta \rangle_{\mathcal{M}_T, C_T} \leq 0.$$

We define  $\hat{p}_\vartheta := S_\vartheta^*W_h(S\bar{u} - \mathbf{z})$ , insert it between  $\bar{p}$  and  $\bar{p}_\vartheta$  and obtain

$$0 \leq \langle \bar{u}_\vartheta - \bar{u}, \bar{p} - \hat{p}_\vartheta \rangle_{\mathcal{M}_T, C_T} + \langle \bar{u}_\vartheta - \bar{u}, \hat{p}_\vartheta - \bar{p}_\vartheta \rangle_{\mathcal{M}_T, C_T}. \quad (9.2)$$

For convenience we introduce the variables  $(\hat{y}_\vartheta, \hat{y}_\vartheta(T), \rho\hat{y}_{T_h}^1) = S_\vartheta\bar{u}$  and remark that the state equations for  $(\bar{y}_\vartheta, \bar{y}_{T_h}^1)$  and  $(\hat{y}_\vartheta, \hat{y}_{T_h}^1)$  have the same initial data. With the help of them we rewrite the second term on the right in (9.2) taking first the difference of the discrete state equations (7.3) and (5.1) (taken for  $(\hat{y}_\vartheta, \hat{y}_{T_h}^1)$ ) for  $v = \hat{p}_\vartheta - \bar{p}_\vartheta$ , next the difference of the discrete adjoint state equations (7.4) and (8.1)-(8.2) (taken for  $\hat{p}_\vartheta$ ) for  $v = \bar{y}_\vartheta - \hat{y}_\vartheta$  and  $\varphi = \bar{y}_{T_h}^1 - \hat{y}_{T_h}^1$  and finally using (5.15)

$$\begin{aligned} \langle \bar{u}_\vartheta - \bar{u}, \hat{p}_\vartheta - \bar{p}_\vartheta \rangle_{\mathcal{M}_T, C_T} &= B_\sigma(\bar{y}_\vartheta - \hat{y}_\vartheta, \hat{p}_\vartheta - \bar{p}_\vartheta) + (\rho(\bar{y}_{T_h}^1 - \hat{y}_{T_h}^1), (\hat{p}_\vartheta - \bar{p}_\vartheta)(T))_H \\ &= (\rho(\bar{y}_\vartheta - \hat{y}_\vartheta), \bar{y} - \bar{y}_\vartheta)_{L^2(I \times \Omega)} + (\rho(\bar{y}_\vartheta - \hat{y}_\vartheta)(T), (\bar{y} - \bar{y}_\vartheta)(T))_H + (\rho(\bar{y}_{T_h}^1 - \hat{y}_{T_h}^1), A_h^{-1}(\rho(\partial_t \bar{y}(T) - \bar{y}_{T_h}^1)))_H \\ &= (S_\vartheta\bar{u}_\vartheta - S_\vartheta\bar{u}, S\bar{u} - S_\vartheta\bar{u}_\vartheta)_{\mathcal{Y}_h}. \end{aligned}$$

Further we easily get

$$\begin{aligned} \langle \bar{u}_\vartheta - \bar{u}, \hat{p}_\vartheta - \bar{p}_\vartheta \rangle_{\mathcal{M}_T, C_T} &= (S\bar{u} - S_\vartheta\bar{u}_\vartheta, S_\vartheta\bar{u}_\vartheta - S_\vartheta\bar{u})_{\mathcal{Y}_h} = (S\bar{u} - S_\vartheta\bar{u}_\vartheta, S\bar{u} - S_\vartheta\bar{u})_{\mathcal{Y}_h} - \|S\bar{u} - S_\vartheta\bar{u}_\vartheta\|_{\mathcal{Y}_h}^2 \\ &\leq \frac{1}{2}\|S\bar{u} - S_\vartheta\bar{u}\|_{\mathcal{Y}_h}^2 - \frac{1}{2}\|S\bar{u} - S_\vartheta\bar{u}_\vartheta\|_{\mathcal{Y}_h}^2. \end{aligned}$$

Thus (9.2) implies

$$\begin{aligned} \|S\bar{u} - S_\vartheta\bar{u}_\vartheta\|_{\mathcal{Y}_h}^2 &\leq 2\langle \bar{u}_\vartheta - \bar{u}, \bar{p} - \hat{p}_\vartheta \rangle_{\mathcal{M}_T, C_T} + \|S\bar{u} - S_\vartheta\bar{u}\|_{\mathcal{Y}_h}^2 \\ &\leq 2(\|\bar{u}_\vartheta\|_{\mathcal{M}_T} + \|\bar{u}\|_{\mathcal{M}_T})\|\bar{p} - \hat{p}_\vartheta\|_{C_T} + \|S\bar{u} - S_\vartheta\bar{u}\|_{\mathcal{Y}_h}^2. \end{aligned}$$

Finally by applying bounds (4.2) and (7.1) we derive (9.1).  $\square$

This proposition is important since it allows one to derive estimates for  $\bar{y} - \bar{y}_\vartheta$  with the help of the above error estimates for the discrete state and adjoint state equations.

**Theorem 9.2.** (1) *Let  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$ ,  $\mathbf{z} \in \mathcal{Y}^{1/2}$  and  $\mathbf{y} \in V \times H$ . Then the following error estimate holds*

$$\|\bar{y} - \bar{y}_\vartheta\|_{L^2(I \times \Omega)} + \|(\bar{y} - \bar{y}_\vartheta)(T)\|_H + \|\rho(\partial_t \bar{y}(T) - \bar{y}_{T_h}^1)\|_{\mathcal{V}_{\kappa, h}^*} \leq C(\tau + h)^{1/3}. \quad (9.3)$$

(2) Let  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ ,  $\mathbf{z} \in \mathcal{Y}^{3/2}$  and  $\mathbf{y} \in H^{(3/2)} \times H^{(1/2)}$ . Then the following higher order error estimate holds

$$\|\bar{y} - \bar{y}_\vartheta\|_{L^2(I \times \Omega)} + \|(\bar{y} - \bar{y}_\vartheta)(T)\|_H + \|\rho(\partial_t \bar{y}(T) - \bar{y}_{Th}^1)\|_{\mathcal{V}_{\kappa, h}^*} \leq C(\tau + h)^{2/3}. \quad (9.4)$$

*Proof.* 1. Let us base on Proposition 9.1. First, Proposition 6.2 (4) implies

$$\|S\bar{u} - S_\vartheta \bar{u}\|_{\mathcal{Y}_h} \leq c(\tau + h)^{1/3} (\|\bar{u}\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{y}\|_{V \times H}).$$

Second, Proposition 8.1 (3) leads to

$$\|S^*W(S\bar{u} - \mathbf{z}) - S_\vartheta^*W_h(S\bar{u} - \mathbf{z})\|_{C_T} \leq c(\tau + h)^{2/3} (\|\bar{u}\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{1/2}} + \|\mathbf{y}\|_{H^{(1/2)} \times H^{(-1/2)}}).$$

Now owing to Proposition 9.1, embedding (3.27) and bound (4.2) for  $\bar{u}$  error estimate (9.3) is proved.

2. First, Proposition 6.2 (3) implies

$$\|S\bar{u} - S_\vartheta \bar{u}\|_{\mathcal{Y}_h} \leq c(\tau + h)^{2/3} \left( \|\bar{u}\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}} \right).$$

Second, Proposition 8.1 (4) leads to

$$\|S^*W(S\bar{u} - \mathbf{z}) - S_\vartheta^*W_h(S\bar{u} - \mathbf{z})\|_{C_T} \leq c(\tau + h)^{4/3} (\|\bar{u}\|_{H^1(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{3/2}} + \|\mathbf{y}\|_{H^{(3/2)} \times H^{(1/2)}}).$$

Now owing to Proposition 9.1, embedding (3.28) and Theorem 4.4 for  $\bar{u}$  error estimate (9.4) is proved too.  $\square$

**Remark 9.3.** Note that our error bounds could be better provided that one would improve the last term on the right in (9.1) by increasing the power 1/2. But this seems a complicated problem.

## 10. ERROR ESTIMATE FOR THE COST FUNCTIONAL

In this section we derive error estimate for the cost functional. We first observe the inequalities

$$j(\bar{u}) \leq j(\bar{u}_\vartheta), \quad j_\vartheta(\bar{u}_\vartheta) \leq j_\vartheta(\bar{u})$$

which can be equivalently rewritten in the form

$$j(\bar{u}) - j_\vartheta(\bar{u}) \leq j(\bar{u}) - j_\vartheta(\bar{u}_\vartheta) \leq j(\bar{u}_\vartheta) - j_\vartheta(\bar{u}_\vartheta). \quad (10.1)$$

Therefore, to bound  $|j(\bar{u}) - j_\vartheta(\bar{u}_\vartheta)|$  below we apply the following result.

**Proposition 10.1.** *Let  $\mathbf{y} \in V \times H$ . Then for any  $u \in \mathcal{M}_T$*

$$\begin{aligned} |j(u) - j_\vartheta(u)| &\leq c \left( \|Su - S_\vartheta u\|_{\mathcal{Y}_h}^2 + \right. \\ &\quad \left. + (\|u\|_{\mathcal{M}_T} + \|\mathbf{y}\|_{V \times H}) \left( \|p - p_\vartheta\|_{C_T} + \|p(0) - p_\vartheta(0)\|_H + h\|\partial_t p(0)\|_H + \|\rho(\partial_t p(0) - p_{0h}^1)\|_{\mathcal{V}_{\kappa, h}^*} \right) \right. \\ &\quad \left. + \|r_h A^{-1}(\rho \partial_t y(T))\|_V^2 + \|r_h A^{-1} z_3\|_V^2 \right) \quad (10.2) \end{aligned}$$

with  $(y, y(T), \rho \partial_t y(T)) = Su$  and the same  $p$  and  $(p_\vartheta, p_{0h}^1)$  as in Proposition 8.1.

*Proof.* Let  $u \in \mathcal{M}_T$ . According to the definitions of the continuous and discrete cost functionals and property (5.13) for  $\tilde{w} = w$  and  $\tilde{w}_h = w_h$  we get

$$\begin{aligned} j(u) - j_\vartheta(u) &= \frac{1}{2} \|Su - \mathbf{z}\|_{\mathcal{Y}}^2 - \frac{1}{2} \|S_\vartheta u - \mathbf{z}\|_{\mathcal{Y}_h}^2 \\ &= \frac{1}{2} (Su - S_\vartheta u, Su + S_\vartheta u - 2\mathbf{z})_{\mathcal{Y}_h} + \frac{1}{2} \|A^{-1}(\rho \partial_t y(T) - z_3)\|_{\mathcal{V}_\kappa}^2 - \frac{1}{2} \|A_h^{-1}(\rho \partial_t y(T) - z_3)\|_{\mathcal{V}_{\kappa, h}}^2 \\ &= -\frac{1}{2} \|Su - S_\vartheta u\|_{\mathcal{Y}_h}^2 + (Su - S_\vartheta u, Su - \mathbf{z})_{\mathcal{Y}_h} + \frac{1}{2} \|r_h A^{-1}(\rho \partial_t y(T) - z_3)\|_{\mathcal{V}_\kappa}^2. \quad (10.3) \end{aligned}$$

We set  $p_{Th} := A_h^{-1}(\rho \partial_t y(T) - z_3)$ .

Owing to the adjoint problem (3.12) with  $(\phi, p^1, p^0) = W(y - z_1, y(T) - z_2, \rho \partial_t y(T) - z_3)$  we get

$$\begin{aligned} (Su, Su - \mathbf{z})_{\mathcal{Y}_h} - (A_h^{-1}(\rho \partial_t y(T)), p_{Th})_{\mathcal{V}_\kappa} &= (\rho y, y - z_1)_{L^2(I \times \Omega)} + (\rho y(T), y(T) - z_2)_H \\ &= \langle u, p \rangle_{\mathcal{M}_T, \mathcal{C}_T} + (\rho y^1, p(0))_H - \langle \rho \partial_t y(T), p^0 \rangle_\Omega - (\rho y^0, \partial_t p(0))_H. \end{aligned}$$

Similarly owing to the general discrete adjoint state equation (8.1)-(8.2) for  $v = y_\vartheta$  and the discrete state equation (5.1)-(5.2) for  $v = p_\vartheta$  and  $\varphi = p_{0h}^1$  we get

$$\begin{aligned} (S_\vartheta u, Su - \mathbf{z})_{\mathcal{Y}_h} - (A_h^{-1}(\rho y_{Th}^1), p_{Th})_{\mathcal{V}_\kappa} &= (\rho y_\vartheta, y - z_1)_{L^2(I \times \Omega)} + (\rho y_\vartheta(T), y(T) - z_2)_H \\ &= B_\sigma(y_\vartheta, p_\vartheta) - (\rho y_\vartheta(0), p_{0h}^1)_H \\ &= \langle u, p_\vartheta \rangle_{\mathcal{M}_T, \mathcal{C}_T} + (\rho y^1, p_\vartheta(0))_H - (\rho y_{Th}^1, p_\vartheta(T))_H - (\rho y^0, p_{0h}^1)_H. \end{aligned}$$

In addition owing to the definitions (8.2) of  $p_\vartheta(T)$  and (5.15) of  $A_h^{-1}$ , we can write

$$(\rho y_{Th}^1, p_\vartheta(T))_H = (\rho y_{Th}^1, p_{Th})_H = (A_h^{-1}(\rho y_{Th}^1), p_{Th})_{\mathcal{V}_\kappa}.$$

Consequently we obtain

$$\begin{aligned} (Su - S_\vartheta u, Su - \mathbf{z})_{\mathcal{Y}_h} &= (Su, Su - \mathbf{z})_{\mathcal{Y}_h} - (S_\vartheta u, Su - \mathbf{z})_{\mathcal{Y}_h} \\ &= \langle u, p - p_\vartheta \rangle_{\mathcal{M}_T, \mathcal{C}_T} - (\rho y^0, \partial_t p(0) - p_{0h}^1)_H + (\rho y^1, p(0) - p_\vartheta(0))_H \\ &\quad + (A_h^{-1}(\rho \partial_t y(T)), p_{Th})_{\mathcal{V}_\kappa} - \langle \rho \partial_t y(T), p^0 \rangle_\Omega. \end{aligned} \quad (10.4)$$

In addition using property (5.13) we derive

$$\begin{aligned} (A_h^{-1}(\rho \partial_t y(T)), p_{Th})_{\mathcal{V}_\kappa} - \langle \rho \partial_t y(T), p^0 \rangle_\Omega &= (A_h^{-1}(\rho \partial_t y(T)), p_{Th})_{\mathcal{V}_\kappa} - (A^{-1}(\rho \partial_t y(T)), p^0)_{\mathcal{V}_\kappa} \\ &= -(r_h A^{-1}(\rho \partial_t y(T)), r_h A^{-1}(\rho \partial_t y(T) - z_3))_{\mathcal{V}_\kappa}. \end{aligned} \quad (10.5)$$

Next, for the term  $(\rho y^0, \partial_t p(0) - p_{0h}^1)_H$  in (10.4) we have

$$\begin{aligned} |(\rho y^0, \partial_t p(0) - p_{0h}^1)_H| &= |(\rho(y^0 - \pi_h^0 y^0), \partial_t p(0) - p_{0h}^1)_H + (\rho \pi_h^0 y^0, \partial_t p(0) - p_{0h}^1)_H| \\ &\leq |(\rho(y^0 - \pi_h^0 y^0), \partial_t p(0))_H| + c \|\pi_h^0 y^0\|_V \|\rho(\partial_t p(0) - p_{0h}^1)\|_{\mathcal{V}_{\kappa, h}^*} \\ &\leq c_1 \|y^0\|_V \left( h \|\partial_t p(0)\|_H + \|\rho(\partial_t p(0) - p_{0h}^1)\|_{\mathcal{V}_{\kappa, h}^*} \right) \end{aligned} \quad (10.6)$$

due to the bounds  $\|y^0 - \pi_h^0 y^0\|_{H_\rho} \leq \|y^0 - \pi_h^1 y^0\|_{H_\rho}$ , (5.18) and (6.2). Clearly also  $|(\rho y^1, p(0) - p_\vartheta(0))_H| \leq \|y^1\|_H \|p(0) - p_\vartheta(0)\|_H$ . Finally from (10.3)-(10.6) we derive (10.2).  $\square$

Now we prove for the cost functional a higher order error estimate than (9.3) for the state variable in the case  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$ .

**Theorem 10.2.** *Let  $\mathcal{M}_T = L^2(I, \mathcal{M}(\Omega))$ ,  $\mathbf{z} \in \mathcal{Y}^{1/2}$  and  $\mathbf{y} \in V \times H$ . Then the following error estimate for the cost functional holds*

$$|j(\bar{u}) - j_\vartheta(\bar{u}_\vartheta)| \leq C(\tau + h)^{2/3}.$$

*Proof.* Let us base on Proposition 10.1 and take any  $u \in L^2(I, \mathcal{M}(\Omega))$ . Owing to Proposition 6.2 (2) we have

$$\|Su - S_\vartheta u\|_{\mathcal{Y}_h} \leq c(\tau + h)^{1/3} (\|u\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{y}\|_{V \times H^{(-1/2)}}).$$

Proposition 8.1 (3) leads to

$$\|p - p_\vartheta\|_{C(\bar{I} \times \bar{\Omega})} \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I, H^{(-1/2)})} + \|\mathbf{z}\|_{\mathcal{Y}^{1/2}} + \|\mathbf{y}\|_{H^{(1/2)} \times H^{(-1/2)}}).$$

Owing to Propositions 3.5(1) (applied to the adjoint state problem) and 3.7 we have

$$\|\partial_t p(0)\|_H \leq \|\partial_t p\|_{C(\bar{I}, H)} \leq c(\|u\|_{L^2(I, V^*)} + \|\mathbf{y}\|_{H \times V^*} + \|\mathbf{z}\|_{\mathcal{Y}})$$

(like in estimates (8.11)-(8.12) for  $\alpha = 1$ ). By using estimate (5.17) for  $\lambda = -1/2$  we obtain

$$\|r_h A^{-1}(\rho \partial_t y(T))\|_V + \|r_h A^{-1} z_3\|_V \leq ch^{1/2} (\|\partial_t y(T)\|_{H^{(-1/2)}} + \|z_3\|_{H^{(-1/2)}}).$$

By collecting all these estimates together with embedding (3.27), Proposition 8.1 (2) to bound  $\|\rho(\partial_t p(0) - p_{0h}^1)\|_{V_{\kappa,h}^*}$  and applying Proposition 10.1, we derive

$$|j(u) - j_\vartheta(u)| \leq c(\tau + h)^{2/3} (\|u\|_{L^2(I, \mathcal{M}(\Omega))} + \|\mathbf{z}\|_{Y^{1/2}} + \|\mathbf{y}\|_{V \times H})^2.$$

Owing to inequalities (10.1) together with bounds (4.2) for  $\bar{u}$  and (7.1) for  $\bar{u}_\vartheta$  the proof is complete.  $\square$

**Remark 10.3.** In the case  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  we know that  $\bar{u} \in H^1(I, M(\Omega))$  (cf. Theorem 4.4). The lack of the corresponding bound  $\|\bar{u}_\vartheta\|_{H^1(I, M(\Omega))} \leq C$  on the discrete level, does not allow us to prove the error estimate  $|j(\bar{u}) - j_\vartheta(\bar{u}_\vartheta)| \leq C(\tau + h)^{4/3}$ . The estimate  $|j(\bar{u}) - j_\vartheta(\bar{u}_\vartheta)| \leq C(\tau + h)^{2/3}$  follows directly from (9.4).

## 11. TIME-STEPPING FORMULATION

In this section we discuss the time-stepping formulation of the discrete state equation (5.1)-(5.2) and the discrete adjoint state equation (7.4). We introduce the piecewise-linear ‘‘hat’’ functions such that  $e_m^\tau(t_k) = \delta_{m,k}$  for any  $k, m = 0, \dots, M$ , where  $\delta_{m,k}$  is the Kronecker delta. We recall that  $e_m^\tau$  are ‘‘half’’ hat functions for  $m = 0, M$ . There holds  $V_\tau = \text{span}\{e_0^\tau, \dots, e_M^\tau\}$ . Similarly, we introduce the spatial hat functions such that  $e_j^h(x_k) = \delta_{j,k}$  for any  $j = 1, \dots, N-1$  and  $k = 0, \dots, N$ ; then  $V_h = \text{span}\{e_1^h, \dots, e_{N-1}^h\}$ .

Then the approximate state variable  $y_\vartheta \in V_\vartheta$  can be represented in the following forms

$$y_\vartheta(t, x) = \sum_{m=0}^M \sum_{j=1}^{N-1} y_{m,j} e_j^h(x) e_m^\tau(t) = \sum_{m=0}^M y_m^h(x) e_m^\tau(t) = \sum_{j=1}^{N-1} y_j^\tau(t) e_j^h(x), \quad (t, x) \in \bar{I} \times \bar{\Omega} \quad (11.1)$$

with  $y_{m,j} \in \mathbb{R}$ ,  $y_m^h \in V_h$  and  $y_j^\tau \in V_\tau$ .

We also define the forward and backward difference and the average in time operators

$$\delta_t v_m = \frac{v_{m+1} - v_m}{\tau}, \quad \bar{\delta}_t v_m = \frac{v_m - v_{m-1}}{\tau},$$

$$B^\tau v_0 = \frac{1}{3}v_0 + \frac{1}{6}v_1, \quad B^\tau v_m = \frac{1}{6}v_{m-1} + \frac{2}{3}v_m + \frac{1}{6}v_{m+1}, \quad 1 \leq m \leq M-1, \quad B^\tau v_M = \frac{1}{6}v_{M-1} + \frac{1}{3}v_M.$$

We define the self-adjoint positive-definite operators  $B_h$  and  $L_h$  acting in  $V_h$  (in other words, the mass and stiffness matrices) such that

$$(B_h \varphi_h, \psi_h)_{V_h} = (\rho \varphi_h, \psi_h)_H, \quad (L_h \varphi_h, \psi_h)_{V_h} = (\kappa \partial_x \varphi_h, \partial_x \psi_h)_H \quad \forall \varphi, \psi \in V_h.$$

For  $w \in V^*$  and  $u \in L^2(I, V^*)$  we define the vectors  $w^h = \{\langle w, e_j^h \rangle_\Omega\}_{j=1}^{N-1}$  and

$$u_m^\vartheta = \frac{1}{\tau} \left\{ \langle u, e_j^h \rangle_\Omega, e_m^\tau \right\}_{L^2(I)}_{j=1}^{N-1}, \quad 1 \leq m \leq M-1, \quad u_m^\vartheta = \frac{2}{\tau} \left\{ \langle u, e_j^h \rangle_\Omega, e_m^\tau \right\}_{L^2(I)}_{j=1}^{N-1}, \quad m = 0, M.$$

We recall the form of the discrete state (11.1).

The forward time-stepping is implemented as follows. The integral identities (5.1)-(5.2) are equivalent to the operator equations

$$(B_h + \sigma \tau^2 L_h) \delta_t \bar{\delta}_t y_{\vartheta, m} + L_h y_{\vartheta, m} = u_m^\vartheta, \quad m = 2, \dots, M-1, \quad (11.2)$$

$$(B_h + \sigma \tau^2 L_h) \delta_t y_{\vartheta, 1} + \frac{\tau}{2} L_h y_{\vartheta, 0} = (\rho y^1)^h + \frac{\tau}{2} u_0^\vartheta, \quad (11.3)$$

$$B_h y_{\vartheta, 0} = (\rho y^0)^h \quad (11.4)$$

followed by the counterpart of (11.3) at time  $T$  for  $y_{T,h}^1$ :

$$B_h y_{T,h}^1 = (B_h + \sigma \tau^2 L_h) \bar{\delta}_t y_{\vartheta, M} - \frac{\tau}{2} L_h y_{\vartheta, M} + \frac{\tau}{2} u_M^\vartheta. \quad (11.5)$$



Next *the adjoint (backward) time-stepping* is implemented in a similar manner. Namely, the integral identities (7.4) are equivalent to the operator equations

$$(B_h + \sigma\tau^2 L_h)\delta_t \bar{\delta}_t p_{\vartheta,m} + L_h p_{\vartheta,m} = B_h B^\tau y_{\vartheta,m} - (\rho z_1)_m^\vartheta, \quad m = M-1, \dots, 1, \quad (11.6)$$

$$-(B_h + \sigma\tau^2 L_h)\bar{\delta}_t p_{\vartheta,M} + \frac{\tau}{2} L_h p_{\vartheta,M} = B_h y_{\vartheta,M} - (\rho z_2)^h + \frac{\tau}{2} (B_h B^\tau y_{\vartheta,M} - (\rho z_1)_M^\vartheta), \quad (11.7)$$

$$L_h p_{\vartheta,M} = B_h y_{T^h}^1 - z_3^h, \quad (11.8)$$

followed by the counterpart of (11.5) for  $p_{0h}^1$ :

$$B_h p_{0h}^1 = (B_h + \sigma\tau^2 L_h)\delta_t p_{\vartheta,0} + \frac{\tau}{2} L_h p_{\vartheta,0} - \frac{\tau}{2} (B_h B^\tau y_{\vartheta,0} - (\rho z_1)_0^\vartheta). \quad (11.9)$$

**Remark 11.1.** For  $\sigma = 1/4$  the three-level time stepping scheme (11.2)-(11.5) is closely related to the well-known two-level Crank-Nicolson method applied to the first order in time system

$$\begin{cases} \partial_t y = v, \quad \rho \partial_t v - \partial_x(\kappa \partial_x y) = u & \text{in } I \times \Omega \\ y = 0 & \text{on } I \times \partial\Omega \\ y = y^0, \quad v = y^1 & \text{in } \{0\} \times \Omega, \end{cases}$$

see [47, Section 8] for details, as well as to the Petrov-Galerkin method described in [30]. After the mass lumping, for  $\sigma = 0$  our method becomes explicit and is related to the Leap-Frog method; moreover, for any  $\sigma$  it becomes close to three-level finite-difference schemes with such weight in time, eg. see [44].

## 12. CONTROL DISCRETIZATION. SOLUTION PROCESS AND $L^2(I \times \Omega)$ -REGULARIZATION

Now we discuss in more detail solving of the semi-discrete optimization problem  $(\mathcal{P}_\vartheta)$  in the case  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$ .

An important point is that we can seek its solution in the form

$$\bar{u}_\vartheta \in \mathcal{M}_\vartheta := V_\tau \otimes \mathcal{M}_h, \quad \mathcal{M}_h := \text{span}\{\delta_{x_1}, \dots, \delta_{x_{N-1}}\} \subset \mathcal{M}(\Omega).$$

To show that, let  $\pi_\tau^0$  be the projector in  $L^2(I)$  on  $V_\tau$ . Note that, for  $\eta \in L^2(I)$ , it satisfies

$$(B^\tau \pi_\tau^0 \eta)_m = \frac{1}{\tau} (\eta, e_m^\tau)_{L^2(I)} \quad \text{for } 1 \leq m \leq M-1, \quad (B^\tau \pi_\tau^0 \eta)_m = \frac{2}{\tau} (\eta, e_m^\tau)_{L^2(I)} \quad \text{for } m = 0, M.$$

Then we define  $\Pi_h: \mathcal{M}(\Omega) \rightarrow \mathcal{M}_h$  by  $\Pi_h w := \sum_{j=1}^{N-1} \langle w, e_j^h \rangle_\Omega \delta_{x_j}$  and  $\Pi_\vartheta = \pi_\tau^0 \Pi_h$ . The following identity holds

$$\langle \Pi_\vartheta u, v \rangle_{\mathcal{M}_T, \mathcal{C}_T} = \langle u, \pi_\tau^0 i_h v \rangle_{\mathcal{M}_T, \mathcal{C}_T} \quad \forall u \in \mathcal{M}_T, v \in \mathcal{C}_T$$

with the interpolation operator  $i_h: \mathcal{C}_0(\Omega) \rightarrow V_h$  such that  $i_h w(x_j) = w(x_j)$  for all  $j = 0, \dots, N$ . In particular, if  $v \in V_\vartheta$ , then

$$\langle \Pi_\vartheta u, v \rangle_{\mathcal{M}_T, \mathcal{C}_T} = \langle u, v \rangle_{\mathcal{M}_T, \mathcal{C}_T},$$

and consequently (like in [31, Lemma 3.11]) we have  $S_\vartheta = S_\vartheta \circ \Pi_\vartheta$  as well as  $\|\Pi_\vartheta u\|_{\mathcal{M}_T} \leq \|u\|_{\mathcal{M}_T}$ . Thus for each solution  $\tilde{u}_\vartheta$  of problem  $(\mathcal{P}_\vartheta)$ , the discrete control  $\Pi_\vartheta \tilde{u}_\vartheta$  satisfies

$$j_\vartheta(\tilde{u}_\vartheta) = j_\vartheta(\Pi_\vartheta \tilde{u}_\vartheta).$$

Therefore  $\Pi_\vartheta \tilde{u}_\vartheta$  is also a solution of  $(\mathcal{P}_\vartheta)$ . This is a justification for solving the fully discrete problem

$$j_\gamma(u_\vartheta) = \frac{1}{2} \|S_\vartheta u_\vartheta - \mathbf{z}\|_{\mathcal{Y}_h}^2 + \alpha \|u_\vartheta\|_{\mathcal{M}(\Omega, L^2(I))} \rightarrow \min_{u_\vartheta \in \mathcal{M}_\vartheta} \quad (12.1)$$

in order to get a solution of  $(\mathcal{P}_\vartheta)$ .

The direct solution of (12.1) by means of a generalized Newton type method is a challenging problem since a proper globalization strategy is needed, see [37]. Thus we propose a solution strategy based on an additional  $L^2(I \times \Omega)$ -regularization of (12.1) with a parameter  $\gamma > 0$  and a continuation method. For high values of  $\gamma$  the corresponding Newton type method converges independently of the initial guess in numerical practice. Thus the continuation strategy can be seen as simple globalization strategy.

On the continuous level we consider the following regularized problem

$$j_\gamma(u) = \frac{1}{2} \|Su - \mathbf{z}\|_{\mathcal{Y}}^2 + \alpha \|u\|_{\mathcal{M}(\Omega, L^2(I))} + \frac{\gamma}{2} \|u\|_{L^2(I \times \Omega)}^2 \rightarrow \min_{u \in L^2(I \times \Omega)}. \quad (12.2)$$

It is possible to formulate a semi-smooth Newton method for this problem on the continuous level which is based on the following necessary and sufficient optimality condition

$$\bar{u}_\gamma(t, x) = -\frac{1}{\gamma} \max \left( 0, 1 - \frac{\alpha}{\|\bar{p}(\cdot, x)\|_{L^2(I)}} \right) \bar{p}(t, x), \quad (t, x) \in I \times \Omega, \quad (12.3)$$

with  $\bar{p} = S^*W_h(S\bar{u}_\gamma - \mathbf{z})$ . Moreover, this semi-smooth Newton method is superlinear convergent. Let  $\bar{u}_\gamma$  and  $\bar{u}$  be the unique solutions of (12.2) and  $(\mathcal{P})$ . Then we have  $\bar{u}_\gamma \rightharpoonup^* \bar{u}$  in  $\mathcal{M}(\Omega, L^2(I))$ , see [31, 41, 24]. This justifies the use of a continuation strategy in  $\gamma$ . The control discretization described above can not be used for (12.2). Instead we propose to use discrete controls from  $V_\vartheta$ , i.e.,

$$u_\vartheta(t, x) = \sum_{m=0}^M \sum_{j=1}^{N-1} u_{m,j} e_m^\tau(t) e_j^h(x) = \sum_{j=1}^{N-1} u_j(t) e_j^h(x) = \sum_{m=0}^M u_m(x) e_m^\tau(t),$$

cf. (11.1). In particular, we solve the following fully discrete regularized problem

$$j_\vartheta^\gamma(u_\vartheta) = \frac{1}{2} \|S_\vartheta(l_\vartheta u_\vartheta) - \mathbf{z}\|_{\mathcal{Y}_h}^2 + \alpha \|u_\vartheta\|_{\mathcal{M}(\Omega, L^2(I)), h} + \frac{\gamma}{2} \|u_\vartheta\|_{L^2(I \times \Omega), h}^2 \rightarrow \min_{u_\vartheta \in V_\vartheta} \quad (12.4)$$

with

$$\|u_\vartheta\|_{\mathcal{M}(\Omega, L^2(I)), h} = \sum_{j=1}^{N-1} d_j \|u_j\|_{L^2(I)}, \quad \|u_\vartheta\|_{L^2(I \times \Omega), h}^2 = \sum_{m=0}^M (B^\tau u_m)^t D (B^\tau u_m)$$

where  $D = \text{diag}(d_1, \dots, d_{N-1})$  is the lumped mass matrix. Moreover, the operator  $l_\vartheta$  defined by

$$(l_\vartheta u_\vartheta, v_\vartheta)_{L^2(I \times \Omega), h} = \sum_{m=0}^M (B^\tau u_m)^t D (B^\tau v_m) \quad \forall u_\vartheta, v_\vartheta \in V_\vartheta.$$

The use of  $D$  allows us to derive the following optimality conditions for (12.4)

$$\bar{u}_{m,j}^\gamma = -\frac{1}{\gamma} \max \left( 0, 1 - \frac{\alpha}{\|\bar{p}_{\vartheta \cdot, j}\|_{L^2(I)}} \right) \bar{p}_{\vartheta m, j}, \quad (12.5)$$

for all  $m$  and  $j$ , with  $\bar{p}_\vartheta = S_\vartheta^* W_h(S_\vartheta \bar{u}_\vartheta - \mathbf{z})$ , cf. (12.3). Based on (12.5) we can set up a semi-smooth Newton method. Since problem (12.4) is a discretization of (12.2), we can expect that this method behaves mesh independently. Let  $\bar{u}_\vartheta^\gamma = \sum_{j=1}^{N-1} u_j(t) e_j^h$  be the solution of (12.4) and we define

$$\tilde{u}_\vartheta^\gamma = \sum_{j=1}^{N-1} d_j u_j(t) \delta_{x_j}.$$

As  $\gamma \rightarrow 0$  the control  $\tilde{u}_\vartheta^\gamma$  tends to a solution of (12.1) justifying the use of this control discretization and the continuation strategy. For more details see [41].

### 13. NUMERICAL RESULTS

In this section, we present results of numerical experiments and consider two examples both involving zero initial data  $y^0 = y^1 = 0$ , the control space  $\mathcal{M}_T = \mathcal{M}(\Omega, L^2(I))$  and the tracking functional

$$F(y) = \frac{1}{2} \|y - z\|_{L^2(I \times \Omega)}^2, \quad z(x) := \frac{1}{\sqrt{2\pi\rho}} e^{-\frac{(x-\lambda)^2}{2\rho^2}}$$

with the time independent desired state  $z$  which is a Gaussian centered at  $x = \lambda$ . We choose  $\rho = 0.1$  and  $\lambda$  as an irrational parameter.

For sufficiently large  $\alpha$  ( $\alpha = 0.1$ ), we expect that the optimal control  $\bar{u}$  consists of one point source with a position close to  $\lambda$ . If the Gaussian would move through the domain, a point source shaped  $\bar{u}$  is not able to follow the center of the Gaussian since  $\mathcal{M}(\Omega, L^2(I))$  contains no moving point sources. The optimal control would rather consist of some additional fixed point sources. This would not lower the regularity of the state whereas a moving point source can cause it.

The domain  $\Omega$  and the time segment  $I$  are discretized by the uniform grids for  $N = 2^{r_h}$  and  $M = 2^{r_\tau}$  where  $r_\tau, r_h = 2, 3, \dots, r^{\max}$  with  $r^{\max} = 10$ . The stability parameter is fixed to its lowest

value  $\sigma = 1/4$  ensuring unconditional stability of the time-stepping method. The discrete control problem is solved for  $r_h = 2, 3, \dots, r^{\max}$  and the fixed  $r_\tau = r^{\max}$  and then vice versa. The solution process has been described above in Section 12. Numerically the desired state  $z$  is replaced by  $i_h z$  for simplicity, moreover the corresponding error  $\mathcal{O}(h^2)$  is negligible. Since the optimal pairs  $(\bar{u}, \bar{y})$  are not known in our examples, we replace them by reference solutions  $(\hat{u}, \hat{y})$  which are taken as the approximate solutions on the finest grid level.

**Example 1.** We first take the constant coefficients  $\rho \equiv 1$  and  $\kappa \equiv 1$  and set  $\lambda = \pi/20$ . We depict the reference solution  $(\hat{u}, \hat{y})$  in Figure 1. As expected, the optimal control  $\hat{u}$  consists only of one point




FIGURE 1. Example 1: the reference solution  $(\hat{u}, \hat{y})$



source positioned in the vicinity of  $\lambda$ . Thus, the state  $\hat{y}$  has a kink at this position. Due to reflections at the boundary,  $\hat{y}$  has also kinks at other positions.

Next, we discuss the convergence results. In Figure 2, we see the convergence rate of  $\|\bar{y}_\sigma - \hat{y}\|_{L^2(I \times \Omega)}$  (left) and the objective functional (right) as  $h$  refines. The state error behaves mostly in a linear way and the rate for the functional is close to two; as usual the latter is approximately the doubled rate of the former, and fortunately both are better than the above proved theoretical rates. In Figure 3, we see the similar results as  $\tau$  refines. The error of the functional stagnates at the last  $\tau$  refinement that is caused by a too coarse space grid. Nevertheless, we observe reduced rates for  $\hat{y}$  much less than two caused by its reduced regularity (kinks).

**Example 2.** Now we take the variable coefficient

$$\kappa(x) = \begin{cases} 1.2 & 0.25 < x \leq 1, \\ 0.2 & 0 \leq x < 0.25 \end{cases}$$

and set  $\lambda = \pi/6$ . Our analysis does not cover discontinuous coefficients, but they are of great importance in applications, for example, in seismic tomography. A jump discontinuity in  $\kappa$  translates to a jump in the wave speed which can be related to two different material characteristics changing at the point of discontinuity. Note that the point of discontinuity is a grid point for all grid levels. The reference solution  $(\hat{u}, \hat{y})$  is displayed in Figure 4. Once again  $\hat{u}$  consists of one Dirac measure with a time-dependent intensity located in the vicinity of  $\lambda = \pi/6$  and thus  $\hat{y}$  has a kink at this position. Moreover, we can clearly see that at  $x = 0.25$  the wave speed changes and the wave propagation becomes slower. In Figure 5 we observe that the error of the state variable converges in a linear way whereas the error measured in the objective functional behaves quadratically. Finally in Figure 6 we

FIGURE 2. Example 1: errors as  $h$  refines and  $M = 2^{10}$ FIGURE 3. Example 1: errors as  $\tau$  refines and  $N = 2^{10}$ 



study the error behaviour for  $\tau$ -refinement and find the similar rates of convergence. So somewhat surprisingly we find that the convergence behavior of the error is comparable to the previous Example 1 with  $\kappa \equiv 1$  that stimulates further possible studies.

#### ACKNOWLEDGEMENTS


Ph. Trautmann and B. Vexler were supported by FWF and DFG through the International Research Training Group IGDK 1754 ‘Optimization and Numerical Analysis for Partial Differential Equations with Nonsmooth Structures’. A. Zlotnik has been funded within the framework of the Academic Fund Program at the National Research University Higher School of Economics in 2016-2017 (grant no. 16-01-0054) and by the Russian Academic Excellence Project ‘5-100’. He also thanks the Technical University of Munich for its hospitality in 2014-2015 years.

#### REFERENCES

- [1] L. BALES AND I. LASIECKA, *Continuous finite elements in space and time for the nonhomogeneous wave equation*, Computers & Mathematics with Applications, 27 (1994), pp. 91 – 102.

FIGURE 4. Example 2: the reference solution  $(\hat{u}, \hat{y})$ FIGURE 5. Example 2: errors as  $h$  refines and  $M = 2^{10}$ 

- [2] W. BANGERTH, M. GEIGER, AND R. RANNACHER, *Adaptive Galerkin finite element methods for the wave equation*, Comput. Methods Appl. Math., 10 (2010), pp. 3–48.
- [3] J. BERGH AND J. LÖFSTRÖM, *Interpolation Spaces. An Introduction*, Springer, Berlin-New York, 1976.
- [4] A. BERMÚDEZ, P. GAMALLO, AND R. RODRÍGUEZ, *Finite element methods in local active control of sound*, SIAM J. Control Optim., 43 (2004), pp. 437–465.
- [5] H. BREZIS, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, New York, 2011.
- [6] V. I. BOGACHEV, *Measure theory. Vol. I, II*, Springer, Berlin, 2007.
- [7] K. BREDIES AND H. K. PIKKARAINEN, *Inverse problems in spaces of measures*, ESAIM Control Optim. Calc. Var., 19 (2013), pp. 190–218.
- [8] S. C. BRENNER AND L. R. SCOTT, *The mathematical theory of finite element methods*, vol. 15 of Texts in Applied Mathematics, Springer, New York, 3 ed., 2008.
- [9] P. BRUNNER, C. CLASON, M. FREIBERGER, AND H. SCHARFETTER, *A deterministic approach to the adapted optode placement for illumination of highly scattering tissue*, Biomed. Opt. Express, 3 (2012), pp. 1732–1743.
- [10] G. BUTAZZO, M. GIAQUINTA, AND S. HILDEBRANDT, *One-dimensional variational problems. An introduction*, Clarendon Press, Oxford, 1998.

FIGURE 6. Example 2: errors as  $\tau$  refines and  $N = 2^{10}$ 

- [11] E. CASAS, C. CLASON, AND K. KUNISCH, *Approximation of elliptic control problems in measure spaces with sparse solutions*, SIAM J. Control Optim., 50 (2012), pp. 1735–1752.
- [12] ———, *Parabolic control problems in measure spaces with sparse solutions*, SIAM J. Control Optim., 51 (2013), pp. 28–63.
- [13] E. CASAS AND K. KUNISCH, *Optimal control of semilinear elliptic equations in measure spaces*, SIAM J. Control Optim., 52 (2014), pp. 339–364.
- [14] ———, *Parabolic control problems in space-time measure spaces*, ESAIM Control Optim. Calc. Var., 22 (2016), pp. 355–376.
- [15] E. CASAS, B. VEXLER, AND E. ZUAZUA, *Sparse initial data identification for parabolic pde and its finite element approximations*, Math. Control Relat. Fields, 5 (2015), pp. 377–399.
- [16] E. CASAS AND E. ZUAZUA, *Spike controls for elliptic and parabolic PDEs*, Systems Control Lett., 62 (2013), pp. 311–318.
- [17] C. CLASON AND K. KUNISCH, *A duality-based approach to elliptic control problems in non-reflexive Banach spaces*, ESAIM Control Optim. Calc. Var., 17 (2011), pp. 243–266.
- [18] ———, *A measure space approach to optimal source placement*, Comp. Optim. Appl., 53 (2011), pp. 155–171.
- [19] C. FABRE AND J.-P. PUEL, *Pointwise controllability as limit of internal controllability for the wave equation in one space dimension.*, Portugaliae Mathematica, 51 (1994), pp. 335–350.
- [20] M. FORNASIER AND H. RAUHUT, *Recovery algorithms for vector-valued data with joint sparsity constraints*, SIAM J. Numer. Anal., 46 (2008), pp. 577–613.
- [21] D. A. FRENCH AND T. E. PETERSON, *A continuous space-time finite element method for the wave equation*, Math. Comp., 65 (1996), pp. 491–506.
- [22] M. GUGAT, A. KEIMER, AND G. LEUGERING, *Optimal distributed control of the wave equation subject to state constraints*, ZAMM Z. Angew. Math. Mech., 89 (2009), pp. 420–444.
- [23] M. GUGAT, E. TRÉLAT, AND E. ZUAZUA, *Optimal Neumann control for the 1D wave equation: finite horizon, infinite horizon, boundary tracking terms and the turnpike property*, Systems Control Lett., 90 (2016), pp. 61–70.
- [24] R. HERZOG, G. STADLER, AND G. WACHSMUTH, *Directional sparsity in optimal control of partial differential equations*, SIAM J. Control Optim., 50 (2012), pp. 943–963.
- [25] M. HINZE, *A variational discretization concept in control constrained optimization: The linear quadratic case*, Comput. Optim. Appl., 30 (2005), pp. 45–61.
- [26] L. V. KANTOROVICH AND G. P. AKILOV, *Functional analysis*, Pergamon Press, Oxford, 1982.
- [27] A. KRÖNER, *Adaptive finite element methods for optimal control of second order hyperbolic equations*, Comput. Methods Appl. Math., 11 (2011), pp. 214–240.
- [28] ———, *Semi-smooth Newton methods for optimal control of the dynamical Lamé system with control constraints*, Numer. Funct. Anal. Optim., 34 (2013), pp. 741–769.
- [29] A. KRÖNER AND K. KUNISCH, *A minimum effort optimal control problem for the wave equation*, Comput. Optim. Appl., 57 (2014), pp. 241–270.
- [30] A. KRÖNER, K. KUNISCH, AND B. VEXLER, *Semismooth Newton methods for optimal control of the wave equation with control constraints*, SIAM J. Control Optim., 49 (2011), pp. 830–858.

- [31] K. KUNISCH, K. PIEPER, AND B. VEXLER, *Measure valued directional sparsity for parabolic optimal control problems*, SIAM J. Control Optim., 52 (2014), pp. 3078–3108.
- [32] K. KUNISCH, P. TRAUTMANN, AND B. VEXLER, *Optimal control of the undamped linear wave equation with measure valued controls*, SIAM J. Control Optim., 54 (2016), pp. 1212–1244.
- [33] K. KUNISCH AND D. WACHSMUTH, *On time optimal control of the wave equation and its numerical realization as parametric optimization problem*, SIAM J. Control Optim., 51 (2013), pp. 1232–1262.
- [34] I. LASIECKA AND J. SOKOLOWSKI, *Sensitivity analysis of optimal control problems for wave equations*, SIAM J. Control Optim., 29 (1991), pp. 1128–1149.
- [35] J. LIONS, *Control of distributed singular systems*, Gauthier-Villars, Paris, 1985.
- [36] J. L. LIONS AND E. MAGENES, *Non-homogeneous boundary value problems and applications*, vol. 1, Springer Berlin, 1972.
- [37] A. MILZAREK AND M. ULBRICH, *A semismooth Newton method with multidimensional filter globalization for  $l_1$ -optimization*, SIAM J. Optim., 24 (2014), pp. 298–333.
- [38] B. S. MORDUKHOVICH AND J.-P. RAYMOND, *Dirichlet boundary control of hyperbolic equations in the presence of state constraints*, Appl. Math. Optim., 49 (2004), pp. 145–157.
- [39] ———, *Neumann boundary control of hyperbolic equations with pointwise state constraints*, SIAM J. Control Optim., 43 (2004/05), pp. 1354–1372.
- [40] S. M. NIKOLSKII, *Approximation of functions of several variables and imbedding theorems*, Springer, Berlin, 1975.
- [41] K. PIEPER, *Finite element discretization and efficient numerical solution of elliptic and parabolic sparse control problems*, PhD thesis, TU Munich, 2015.
- [42] K. PIEPER, P. TRAUTMANN, B. TANG QUOC, AND D. WALTER, *Inverse point source location for the helmholtz equation*. Submitted.
- [43] K. PIEPER AND B. VEXLER, *A priori error analysis for discretization of sparse elliptic optimal control problems in measure space*, SIAM J. Control Optim., 51 (2013), pp. 2788–2808.
- [44] A. A. SAMARSKII, *The theory of difference schemes*, Marcel Dekker, New York-Basel, 2001.
- [45] H. TRIEBEL, *Interpolation theory, function spaces, differential operators*, vol. 18 of North-Holland Mathematical Library, North-Holland Publishing Co., Amsterdam-New York, 1978.
- [46] A. A. ZLOTNIK, *Lower error estimates for three-layer difference methods of solving the wave equation with data from hölder spaces*, Math. Notes, 51 (1992), pp. 321–323.
- [47] ———, *Convergence rate estimates of finite-element methods for second order hyperbolic equations*, in Numerical methods and applications, G. I. Marchuk, ed., CRC Press, Boca Raton, FL, 1994, pp. 155–220.
- [48] E. ZUAZUA, *Optimal and approximate control of finite-difference approximation schemes for the 1D wave equation*, Rend. Mat. Appl. (7), 24 (2004), pp. 201–237.