

## *hp*-FINITE ELEMENTS FOR FRACTIONAL DIFFUSION\*

DOMINIK MEIDNER<sup>†</sup>, JOHANNES PFEFFERER<sup>†</sup>, KLEMENS SCHÜRHOZ<sup>†</sup>, AND  
 BORIS VEXLER<sup>†</sup>

**Abstract.** The purpose of this work is to introduce and analyze a numerical scheme to efficiently solve boundary value problems involving the spectral fractional Laplacian. The approach is based on a reformulation of the problem posed on a semi-infinite cylinder in one more spatial dimension. After a suitable truncation of this cylinder, the resulting problem is discretized with linear finite elements in the original domain and with *hp*-finite elements in the extended direction. The proposed approach yields a drastic reduction of the computational complexity in terms of degrees of freedom and even has slightly improved convergence properties compared to the state-of-the-art discretization using linear finite elements for both the original domain and the extended direction. The performance of the method is illustrated by numerical experiments.

**Key words.** fractional Laplace operator, nonlocal operators, finite elements, *hp*-finite elements, discretization error estimates, anisotropic meshes

**AMS subject classifications.** 35S15, 65R20, 65N12, 65N30

**DOI.** 10.1137/17M1135517

**1. Introduction.** In this work, we are concerned with boundary value problems involving the fractional Laplacian, being the prototype of a nonlocal operator. To be more specific, let  $\Omega \subset \mathbb{R}^d$  for  $d \in \{1, 2, 3\}$  be a bounded, convex, polygonal, or polyhedral domain. We are interested in the solution of the boundary value problem

$$(1.1) \quad (-\Delta)^s u = f \quad \text{in } \Omega,$$

where  $(-\Delta)^s$  denotes the spectral fractional Laplacian of order  $s \in (0, 1)$  defined by the eigenvalues and eigenfunctions of the standard Laplacian with homogeneous Dirichlet boundary conditions, precisely introduced in section 2. The main purpose of this paper is to introduce and analyze a numerical scheme to efficiently solve problem (1.1).

Our approach is based on the following equivalent reformulation of problem (1.1) posed on the semi-infinite cylinder  $\{(x, y) \in \Omega \times (0, \infty)\} \subset \mathbb{R}^{d+1}$ : Let  $u$  be the weak solution of the extended problem

$$\begin{aligned} \operatorname{div}(y^{1-2s} \nabla u) &= 0 && \text{in } \Omega \times (0, \infty), \\ u &= 0 && \text{on } \partial\Omega \times [0, \infty), \\ \partial_{\nu} y^{1-2s} u &= d_s f && \text{on } \Omega \times \{0\}; \end{aligned}$$

see section 2 for more details. Then, the trace  $\mathbf{u} = u(\cdot, 0)$  is the solution of the fractional boundary value problem (1.1).

In contrast to the nonlocal problem (1.1), the extended problem is localized. However, a direct application of a finite element method to the extended problem is not feasible because of the semi-infinite domain. As a remedy, the exponential decay of  $u$  in direction  $y$  toward infinity (see Proposition 2.11) can be employed such that a

---

\*Received by the editors June 20, 2017; accepted for publication (in revised form) June 21, 2018; published electronically August 2, 2018.

<http://www.siam.org/journals/sinum/56-4/M113551.html>

<sup>†</sup>Department of Mathematics, Technical University of Munich, Garching b. Munich 85748, Germany (meidner@ma.tum.de, pfefferer@ma.tum.de, klemens.schuerholz@tum.de, vexler@ma.tum.de).

truncation of the semi-infinite cylinder to  $\Omega \times (0, Y)$  becomes reasonable. The extended problem posed on the truncated cylinder can be discretized using finite elements. However, due to the degenerate/singular nature of the extended problem, anisotropic meshes are favorable in order to obtain an optimally convergent numerical scheme. Moreover, the height  $Y$  of the truncated cylinder needs to be chosen dependent on the mesh parameter to ensure the aforementioned convergence. This approach was already pursued in [31] using a discretization with first degree tensor product finite elements on graded meshes in the extended direction; see also [16, 15, 30, 32] for related results. If  $h_\Omega$  denotes the mesh parameter and  $\mathcal{N}_\Omega$  the number of degrees of freedom in  $\Omega$ , then the approach from [31] yields a discretization error of order  $O(h_\Omega |\ln h_\Omega|^s)$  in the corresponding energy norm associated with (1.1) while solving problems with  $O(\mathcal{N}_\Omega^{1+1/d})$  degrees of freedom.

In this work, we introduce and analyze a discretization of the truncated problem with linear finite elements in the original domain  $\Omega$  and with  $hp$ -finite elements on a geometric mesh in the extended direction. This drastically reduces the computational complexity to  $O(\mathcal{N}_\Omega (\ln \mathcal{N}_\Omega)^2)$  degrees of freedom and even yields a slightly better convergence rate of order  $O(h_\Omega)$ . Especially, when  $\mathcal{N}_\Omega$  is large, the difference between the factor  $(\ln \mathcal{N}_\Omega)^2$  and  $\mathcal{N}_\Omega^{1/d}$  becomes clearly perceptible. For instance, in our numerical experiments we could reduce the number of degrees of freedom by a factor of about 111 to obtain an error of less than  $9 \cdot 10^{-3}$  in the case  $s = 0.8$ ; see section 5 for more details. We also notice that our approach and results are not limited to the spectral fractional Laplacian. They naturally extend by only minor modifications to fractional powers of general second order elliptic operators; see [31, section 7] and Remark 2.3 below.

Let us briefly give an overview of other numerical approaches from the literature to solve boundary value problems involving the fractional Laplacian: Due to the spectral definition of the operator, it seems to be natural to compute an approximating, discrete spectral decomposition of the standard Laplacian in order to get an approximation of the solution of (1.1); see [23, 24, 36]. However, this may result in solving a large number of discrete eigenvalue problems. Another approach to determine an approximation to the solution of problem (1.1) is analyzed in [9]; see also [7, 8, 10] for related results. In that reference,  $(-\Delta)^{-s}$  is represented in terms of Bochner integrals involving  $(I - t^2 \Delta)^{-1}$  for  $t \in (0, \infty)$ . Subsequently, different quadrature formulas to approximate this integral are analyzed which require multiple evaluations of  $(I - t_i^2 \Delta_h)^{-1}$  with  $t_i$  being a quadrature point and  $-\Delta_h$  denoting a finite element discretization to  $-\Delta$ . Especially, using sinc quadrature, this results in solving  $O(\ln \mathcal{N}_\Omega)$  standard problems, which can be done in parallel. The computational effort is comparable to that of the  $hp$ -approach in the present paper after having reformulated the arising linear systems of equations as generalized Sylvester matrix equations, for which efficient solvers are available; see section 5 for details. Numerical approaches for the integral definition of the fractional Laplacian, which is not equivalent to the spectral definition considered in the present paper, can be found in [3, 2, 4, 5, 11, 18, 20, 21, 22].

This paper is organized as follows. In section 2, we state the definition of the fractional Laplacian, formulate the extended problem in detail, and introduce the functional framework needed for the subsequent error analysis. Moreover, in this section, we are concerned with several properties of the solution of the extended problem such as a series representation and corresponding regularity results. The discrete, extended problem posed on the truncated cylinder is formulated at the beginning

of section 3. In the extended direction, we distinguish between graded meshes and  $h$ -FEM and geometric meshes and  $hp$ -FEM; see sections 3.1 and 3.2. The error analysis is given in section 4. Thereby, in section 4.1, we mainly recover the results of [31]. The reason for doing this is twofold. First, we are able to slightly improve the mesh grading condition used in [31]. However, the main reason to analyze the  $h$ -FEM on graded meshes before developing the analysis for the  $hp$ -method considered in section 4.2 is that the techniques we use are almost identical for both cases, but the details are simpler for  $h$ -FEM. Implementation aspects and numerical experiments, which underline the efficiency of our approach, are presented in section 5. In the appendix, we collect different results for special functions defined by the modified Bessel functions of second kind. These are especially needed in section 2 for the discussion of the solution of the extended problem.

Finally, we notice that, in the following,  $c$  denotes a generic constant which will always be independent of the mesh parameter  $h_\Omega$  when we analyze the discretization error.

**2. Continuous problem.** Let  $-\Delta$  be the  $L^2(\Omega)$  realization of the Laplacian with homogeneous Dirichlet boundary conditions. It is well-known that  $-\Delta$  has a compact resolvent and its eigenvalues form a nondecreasing sequence  $0 < \lambda_1 < \lambda_2 \leq \dots \leq \lambda_k \leq \dots$  satisfying  $\lim_{k \rightarrow \infty} \lambda_k = \infty$ . We denote by  $\varphi_k$  the orthonormal  $L^2(\Omega)$  eigenfunctions associated with  $\lambda_k$  fulfilling

$$\int_\Omega \nabla \varphi_k \cdot \nabla v \, dx = \lambda_k \int_\Omega \varphi_k v \, dx \quad \forall v \in H_0^1(\Omega).$$

For any  $s \geq 0$ , we introduce the fractional order Sobolev space

$$\mathbb{H}^s(\Omega) = \left\{ \mathbf{v} \in L^2(\Omega) \mid \|\mathbf{v}\|_{\mathbb{H}^s(\Omega)}^2 = \sum_{k=1}^\infty \lambda_k^s \mathbf{v}_k^2 < \infty \text{ with } \mathbf{v}_k = \int_\Omega \mathbf{v} \varphi_k \, dx \right\}.$$

Moreover, we denote by  $\mathbb{H}^{-s}(\Omega)$  the dual space of  $\mathbb{H}^s(\Omega)$ . Then, the spectral fractional Laplacian is defined for  $s \in (0, 1)$  on the space  $\mathbb{H}^s(\Omega)$  as the limit

$$(-\Delta)^s \mathbf{u} = \sum_{k=1}^\infty \lambda_k^s \mathbf{u}_k \varphi_k \in \mathbb{H}^{-s}(\Omega) \quad \text{with} \quad \mathbf{u}_k = \int_\Omega \mathbf{u} \varphi_k \, dx.$$

Due to the Cauchy criterion the limit exists for any  $\mathbf{u} \in \mathbb{H}^s(\Omega)$ . Thus, problem (1.1) has to be understood as follows: Given  $\mathbf{f} \in \mathbb{H}^{-s}(\Omega)$ , find  $\mathbf{u} \in \mathbb{H}^s(\Omega)$  such that

$$(2.1) \quad \sum_{k=1}^\infty \lambda_k^s \mathbf{u}_k \mathbf{v}_k = \int_\Omega \mathbf{f} \mathbf{v} \, dx \quad \forall \mathbf{v} \in \mathbb{H}^s(\Omega) \quad \text{with} \quad \mathbf{v}_k = \int_\Omega \mathbf{v} \varphi_k \, dx.$$

**PROPOSITION 2.1.** *For any  $\mathbf{f} \in \mathbb{H}^{-s}(\Omega)$ , problem (1.1) admits a unique solution  $\mathbf{u} \in \mathbb{H}^s(\Omega)$  fulfilling  $\|\mathbf{u}\|_{\mathbb{H}^s(\Omega)} = \|\mathbf{f}\|_{\mathbb{H}^{-s}(\Omega)}$ . Moreover, there is the series representation*

$$\mathbf{u} = \sum_{k=1}^\infty \mathbf{u}_k \varphi_k \quad \text{with} \quad \mathbf{u}_k = \lambda_k^{-s} \mathbf{f}_k \quad \text{and} \quad \mathbf{f}_k = \langle \mathbf{f}, \varphi_k \rangle_{\mathbb{H}^{-s}(\Omega), \mathbb{H}^s(\Omega)},$$

where  $\langle \cdot, \cdot \rangle_{\mathbb{H}^{-s}(\Omega), \mathbb{H}^s(\Omega)}$  denotes the duality pairing between  $\mathbb{H}^s(\Omega)$  and  $\mathbb{H}^{-s}(\Omega)$ .

*Proof.* The existence of a unique solution and the equality of the norms is a consequence of the Riesz representation theorem. The series representation of  $\mathbf{u}$  is obtained by testing (2.1) with  $\varphi_m \in \mathbb{H}^s(\Omega)$  and using the orthogonality of the eigenfunctions.  $\square$

*Remark 2.2.* We notice that the regularity of  $\mathbf{u}$  can be described in classical fractional Sobolev spaces as well, since

$$\mathbb{H}^s(\Omega) = \begin{cases} H^s(\Omega) & \text{for } 0 < s < \frac{1}{2}, \\ H_{00}^{\frac{1}{2}}(\Omega) & \text{for } s = \frac{1}{2}, \\ H_0^s(\Omega) & \text{for } \frac{1}{2} < s < 1. \end{cases}$$

For more details we refer to, e.g., [31].

*Remark 2.3.* Let  $A$  be the linear operator given by

$$Ay = - \sum_{i,j=1}^n \partial_{x_j} [a_{ij}(x) \partial_{x_i} y] + a_0(x)y$$

with  $a_{i,j} \in W^{1,\infty}(\Omega)$  satisfying  $a_{i,j} = a_{j,i}$  and  $a_0 \in L^\infty(\Omega)$  fulfilling  $a_0 \geq 0$  almost everywhere in  $\Omega$ . Moreover, we assume that there is a constant  $c > 0$  such that

$$\sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \geq c |\xi|^2 \quad \text{for a.a. } x \in \Omega \text{ and } \forall \xi \in \mathbb{R}^d.$$

This operator endowed with homogeneous Dirichlet boundary conditions possesses eigenpairs  $(\lambda_k, \varphi_k)_{k \in \mathbb{N}}$  with identical properties to those of the eigenpairs of the Dirichlet Laplacian. The fractional power  $s$  of such an elliptic operator can then analogously be defined by

$$A^s \mathbf{u} = \sum_{k=1}^\infty \lambda_k^s \mathbf{u}_k \varphi_k \in \mathbb{H}^{-s}(\Omega) \quad \text{with} \quad \mathbf{u}_k = \int_\Omega \mathbf{u} \varphi_k \, dx.$$

In the forthcoming analysis for the fractional Dirichlet Laplacian, we only use the series representation of the operator. For that reason, the results naturally extend to the case of fractional powers of general second order elliptic operators. Of course, the extended problem needs to be modified accordingly. For details we refer to [31, section 7], where this kind of generalization is outlined and error estimates are provided for the  $h$ -FEM variant.

Problem (1.1) can equivalently be posed on a semi-infinite cylinder. In  $\mathbb{R}^d$ , this is due to Caffarelli and Silvestre [13]. We also mention Molčanov and Ostrovskiĭ [29], who discovered this in the context of probabilistic theory. The restriction to bounded domains  $\Omega$  was considered by Stinga and Torrea in [35]; see also [12, 14]. This kind of extension is the basis for the computational approaches in the subsequent sections.

In order to state the extended problem, we first introduce the required notation. We denote by  $C = \Omega \times (0, \infty)$  the aforementioned semi-infinite cylinder and by  $\partial_L C = \partial\Omega \times [0, \infty)$  its lateral boundary. We also need to define a truncated cylinder: for  $Y > 0$ , the truncated cylinder is given by  $C_Y = \Omega \times (0, Y)$  with its lateral boundary  $\partial_L C_Y = \partial\Omega \times [0, Y]$ . As  $C$  and  $C_Y$  are objects in  $\mathbb{R}^{d+1}$ , we use  $y$  to denote

the extended variable, such that a vector  $(x, y) \in \mathbb{R}^{d+1}$  admits the representation  $(x, y) = (x_1, x_2, \dots, x_d, y)$ . Similarly, the gradient in  $\mathbb{R}^{d+1}$  has the representation  $\nabla = (\nabla_x, \partial_y) = (\partial_{x_1}, \partial_{x_2}, \dots, \partial_{x_d}, \partial_y)$ .

Next we introduce weighted Sobolev spaces with a weight function  $y^\alpha$  for  $\alpha \in (-1, 1)$ . In this regard, let  $D \subset \mathbb{R}^d \times (0, \infty)$  be an open set, such as  $C$  or  $C_Y$ . Then, we define the weighted space  $L^2(D, y^\alpha)$  as the space of all measurable functions on  $D$  with finite norm  $\|v\|_{L^2(D, y^\alpha)} = \|y^{\frac{\alpha}{2}}v\|_{L^2(D)}$ . Similarly, the space  $H^1(D, y^\alpha)$  denotes the space of all functions  $v \in L^2(D, y^\alpha)$  whose weak derivatives of first order belong to  $L^2(D, y^\alpha)$ .

To study the extended problems, we introduce the space

$$\mathring{H}_L^1(C, y^\alpha) = \{ v \in H^1(y^\alpha, C) \mid v = 0 \text{ on } \partial_L C \}.$$

The space  $\mathring{H}_L^1(C_Y, y^\alpha)$  is defined analogously but endowed with zero Dirichlet boundary conditions also on  $\Omega \times \{Y\}$ :

$$\mathring{H}_L^1(C_Y, y^\alpha) = \{ v \in H^1(y^\alpha, C_Y) \mid v = 0 \text{ on } \partial_L C_Y \cup (\Omega \times \{Y\}) \}.$$

For  $v \in \mathring{H}_L^1(C, y^\alpha)$ , we denote by  $\text{tr}_\Omega v$  the trace of  $v$  onto  $\Omega \times \{0\}$ , i.e.,  $\text{tr}_\Omega v = v(\cdot, 0)$ .

PROPOSITION 2.4. *For  $\alpha = 1 - 2s$ , it holds that*

$$\text{tr}_\Omega \mathring{H}_L^1(C, y^\alpha) = \mathbb{H}^s(\Omega) \quad \text{and} \quad \|\text{tr}_\Omega v\|_{\mathbb{H}^s(\Omega)} \leq c \|v\|_{\mathring{H}_L^1(C, y^\alpha)}.$$

*Proof.* See [12, Proposition 1.8] for  $s = \frac{1}{2}$  and [14, Proposition 2.1] for  $s \neq \frac{1}{2}$ .  $\square$

Now, we are able to state the extended problem: Given  $\mathfrak{f} \in \mathbb{H}^{-s}(\Omega)$ , find  $u \in \mathring{H}_L^1(C, y^\alpha)$  such that

$$(2.2) \quad \int_C y^\alpha \nabla u \cdot \nabla v \, d(x, y) = d_s \langle \mathfrak{f}, \text{tr}_\Omega v \rangle_{\mathbb{H}^{-s}(\Omega), \mathbb{H}^s(\Omega)} \quad \forall v \in \mathring{H}_L^1(C, y^\alpha)$$

with  $\alpha = 1 - 2s$  and  $d_s = 2^\alpha \frac{\Gamma(1-s)}{\Gamma(s)}$ . That is, the function  $u \in \mathring{H}_L^1(C, y^\alpha)$  is a weak solution of

$$\begin{aligned} \text{div}(y^\alpha \nabla u) &= 0 && \text{in } C, \\ u &= 0 && \text{on } \partial_L C, \\ \partial_{\nu^\alpha} u &= d_s \mathfrak{f} && \text{on } \Omega \times \{0\}, \end{aligned}$$

where we have set  $\partial_{\nu^\alpha} u(x, 0) = -\lim_{y \rightarrow 0} y^\alpha \partial_y u(x, y)$ . Note that subsequently, the parameter  $\alpha$  will always be equal to  $1 - 2s$  for the considered  $s \in (0, 1)$ .

In the remainder of this section, we discuss several properties of the solution  $u$  of (2.2). Due to the following proposition, it is reasonable to determine the solution  $u$  of (2.2) in order to get the solution  $\mathbf{u}$  of (1.1).

PROPOSITION 2.5. *For  $\mathfrak{f} \in \mathbb{H}^{-s}(\Omega)$ , the extended problem (2.2) admits a unique solution  $u \in \mathring{H}_L^1(C, y^\alpha)$ . Furthermore,  $\mathbf{u} = \text{tr}_\Omega u \in \mathbb{H}^s(\Omega)$  solves (1.1).*

*Proof.* See [14, Lemma 2.1].  $\square$

We have the following regularity result.

PROPOSITION 2.6. *Let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2) and  $\mathfrak{f} \in \mathbb{H}^{1-s}(\Omega)$ . Then, it holds that*

$$\|y^{\frac{\alpha}{2}} \nabla_x^2 u\|_{L^2(C)}^2 + \|y^{\frac{\alpha}{2}} \partial_y \nabla_x u\|_{L^2(C)}^2 \leq c \|\mathfrak{f}\|_{\mathbb{H}^{1-s}(\Omega)}^2.$$

*Proof.* Nchetto, Otárola, and Salgado [31, Theorem 2.7] yield

$$\|y^{\frac{\alpha}{2}} \Delta_x u\|_{L^2(C)}^2 + \|y^{\frac{\alpha}{2}} \partial_y \nabla_x u\|_{L^2(C)}^2 = d_s \|f\|_{\mathbb{H}^{1-s}(\Omega)}^2$$

with  $d_s$  given above. Convexity of  $\Omega$  then implies the assertion.  $\square$

*Remark 2.7.* Only for the validity of Proposition 2.6, and hence for the validity of Lemma 4.3 later on, we need to assume convexity. In nonconvex domains we do not have any longer that  $\Delta_x u(\cdot, y) \in L^2(\Omega)$  implies  $|\nabla_x^2 u(\cdot, y)| \in L^2(\Omega)$ . However, it is still possible to get  $u(\cdot, y) \in H^t(\Omega)$ , where  $t \in [1, 2]$  depends on corner and edge singularities included in the solution; see, e.g., [17]. It is even possible to obtain  $|\nabla_x^2 u(\cdot, y)| \in L^2_\beta(\Omega)$  for  $\Delta_x u(\cdot, y) \in L^2_\beta(\Omega)$ . Here,  $L^2_\beta(\Omega)$  denotes a weighted space which is defined such that the singularities in  $\Omega$  are compensated if  $\beta$  is appropriately chosen; see, for instance, [26].

For a series expansion of the solution to (2.2), we introduce  $\psi_s: [0, \infty) \rightarrow \mathbb{R}$  given by

$$(2.3) \quad \psi_s(z) = c_s z^s K_s(z) \quad \text{with} \quad c_s = \frac{2^{1-s}}{\Gamma(s)}.$$

Here,  $K_s(z)$  denotes the modified Bessel function of second kind (see, e.g., [1, section 9.6]) and  $\Gamma(s)$  denotes the gamma function.

**PROPOSITION 2.8.** *For  $f \in \mathbb{H}^{-s}(\Omega)$ , let  $u \in \mathbb{H}^s(\Omega)$  be the solution of (1.1) and  $u \in \dot{H}^1_L(C, y^\alpha)$  be the solution of (2.2). Then, it holds that*

$$u(x, y) = \sum_{k=1}^{\infty} u_k \varphi_k(x) \psi_{s,k}(y),$$

where  $u_k = \int_\Omega u \varphi_k dx$  and  $\psi_{s,k}(y) = \psi_s(\sqrt{\lambda_k} y)$ .

*Proof.* For  $s \neq \frac{1}{2}$ , see [14, Proposition 2.1]. For  $s = \frac{1}{2}$ , the result was proved with  $\psi_{\frac{1}{2}}(z) = e^{-z}$  in [12, Proposition 2.2]. However, by [1, eq. 9.6.23] and the relation  $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ , it holds that

$$c_{\frac{1}{2}} z^{\frac{1}{2}} K_{\frac{1}{2}}(z) = e^{-z} = \psi_{\frac{1}{2}}(z).$$

Hence,  $\psi_s(z) = c_s z^s K_s(z)$  holds also in the case  $s = \frac{1}{2}$ .  $\square$

We state an estimate of the derivatives of  $\psi_{s,k}$  which we will use later. The result is based on properties of  $\psi_s$  which are analyzed in the appendix.

**COROLLARY 2.9.** *Let  $r \in [0, 1]$ . There exists a constant  $c > 0$  depending only on  $s$  such that for any  $y > 0$  and  $n \in \mathbb{N}$  it holds that*

$$|y^n \psi_{s,k}^{(n)}(y)| \leq c 8^n n! \lambda_k^{s-\frac{r}{2}} y^{2s-r}.$$

*Proof.* Simple calculations yield

$$\psi_{s,k}^{(n)}(y) = \frac{d^n}{dy^n} \psi_s(\sqrt{\lambda_k} y) = (\sqrt{\lambda_k})^n \psi_s^{(n)}(\sqrt{\lambda_k} y).$$

Consequently, we obtain by means of Lemma A.3

$$|y^n \psi_{s,k}^{(n)}(y)| = |(\sqrt{\lambda_k} y)^n \psi_s^{(n)}(\sqrt{\lambda_k} y)| \leq c 8^n n! (\sqrt{\lambda_k} y)^{2s-r} = c 8^n n! \lambda_k^{s-\frac{r}{2}} y^{2s-r}. \quad \square$$

Next, we state a result about the exponential decay of  $\psi_{s,k}$  and its derivative. It is based on corresponding results for  $\psi_s$  and its derivative, proved in the appendix.

COROLLARY 2.10. *Let  $y \geq 1$ ,  $r_1 \geq \min(s, \frac{1}{2}) - s$ , and  $r_2 \geq \min(1 - s, \frac{1}{2}) - s$ . Then, there exists a constant  $c_1$  only depending on  $r_1$ ,  $s$ , and  $\lambda_1$  and a constant  $c_2$  only depending on  $r_2$ ,  $s$ , and  $\lambda_1$  such that*

$$|y^{r_1} \psi_{s,k}(y)| \leq c_1 \lambda_k^{-\frac{r_1}{2}} e^{-\frac{\sqrt{\lambda_k}}{2} y} \quad \text{and} \quad |y^{r_2} \psi'_{s,k}(y)| \leq c_2 \lambda_k^{-\frac{r_2}{2}} e^{-\frac{\sqrt{\lambda_k}}{2} y}.$$

*Proof.* Let  $s_0 = \min(s, \frac{1}{2})$ . According to [28, Theorem 5], we get for  $z \geq 0$  that  $z^{s_0} e^z K_s(z)$  is a decreasing function. Consequently, we obtain

$$(2.4) \quad (\sqrt{\lambda_k})^{s_0} e^{\sqrt{\lambda_k}} K_s(\sqrt{\lambda_k}) \leq (\sqrt{\lambda_1})^{s_0} e^{\sqrt{\lambda_1}} K_s(\sqrt{\lambda_1})$$

since the sequence  $(\lambda_k)_{k \in \mathbb{N}}$  is nondecreasing. Moreover, due to the definition of  $\psi_{s,k}$ , we deduce

$$|y^{r_1} \psi_{s,k}(y)| = \lambda_k^{-\frac{r_1}{2}} |(\sqrt{\lambda_k} y)^{r_1} \psi_s(\sqrt{\lambda_k} y)|.$$

Thus, by setting  $a = \sqrt{\lambda_k}$  and  $z = \sqrt{\lambda_k} y \geq a$  in Lemma A.4(a), we obtain the validity of the first inequality of the assertion by means of (2.4). The second inequality can be deduced in the same manner employing Lemma A.4(b).  $\square$

As already mentioned, for computational reasons, the semi-infinite cylinder  $C$  will be truncated to  $C_Y = \Omega \times (0, Y)$  for some  $Y > 0$  later on. Because of this, the behavior of  $\nabla u$  for  $y \rightarrow \infty$  will play a role. It can be estimated as follows.

PROPOSITION 2.11. *For  $f \in \mathbb{H}^{-s}(\Omega)$ , let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Then, there exists a constant  $c > 0$  such that for every  $Y \geq 1$ , it holds that*

$$\|y^{\frac{\alpha}{2}} \nabla u\|_{L^2(C \setminus C_Y)} \leq c e^{-\frac{\sqrt{\lambda_1}}{2} Y} \|f\|_{\mathbb{H}^{-s}(\Omega)}.$$

*Proof.* The result can be found in [31, Proposition 3.1].  $\square$

**3. Discretization.** Let  $\mathcal{T}_\Omega$  be a conforming and quasi-uniform triangulation of  $\Omega$  which is admissible in the sense of Ciarlet. For each  $\mathcal{T}_\Omega = \{K\}$ , let  $K \subset \mathbb{R}^d$  be an element that is isoparametrically equivalent either to the unit cube or to the unit simplex in  $\mathbb{R}^d$ . We introduce the global mesh parameter  $h_\Omega$  with respect to the triangulation of  $\Omega$  by  $h_\Omega = \max_{K \in \mathcal{T}_\Omega} \text{diam } K$ . We always assume that  $h_\Omega \leq \frac{1}{2}$ . On  $\mathcal{T}_\Omega$ , we define a finite element space  $V_h$  as

$$V_h = \{ \mathbf{v} \in C^0(\bar{\Omega}) \mid \mathbf{v}|_K \in \mathbb{P}_1(K), K \in \mathcal{T}_\Omega, \mathbf{v}|_{\partial\Omega} = 0 \}.$$

In the case that  $K$  is a simplex, then  $\mathbb{P}_1(K) = \mathcal{P}_1(K)$ , the set of polynomials on the element  $K$  of degree at most 1. If  $K$  is a cube, then  $\mathbb{P}_1(K)$  equals  $\mathcal{Q}_1(K)$ , the set of polynomials on  $K$  of degree at most 1 in each variable. The number of degrees of freedom in  $V_h$  is denoted by  $\mathcal{N}_\Omega$ . It holds that  $\mathcal{N}_\Omega = O(h_\Omega^{-d})$ .

Furthermore, let  $\mathcal{I}_Y = \{I_m\}$  be a triangulation of the interval  $(0, Y)$  in the sense that  $[0, Y] = \bigcup_{m=1}^M I_m$  with  $I_m = [y_{m-1}, y_m]$  and  $M \in \mathbb{N}$  exactly specified in sections 3.1 and 3.2. Moreover, let  $h_m = |I_m|$ . Next, we introduce a polynomial degree vector  $p = (p_1, p_2, \dots, p_M) \in \mathbb{N}^M$  which will assign to each element  $I_m \in \mathcal{I}_Y$  a maximal polynomial degree  $p_m$ . It will be exactly specified in sections 3.1 and 3.2. On  $\mathcal{I}_Y$ , we define the finite element space  $V_M$  as

$$V_M = \left\{ \xi \in C^0([0, Y]) \mid \xi|_{I_m} \in \mathcal{P}_{p_m}(I_m), I_m \in \mathcal{I}_Y, \xi(Y) = 0 \right\},$$

where  $\mathcal{P}_{p_m}(I_m)$  denotes the space of polynomials up to degree  $p_m$  on  $I_m$ .

Now, the triangulations  $\mathcal{T}_{\Omega,Y}$  of the cylinder  $C_Y$  are constructed as tensor product triangulations by means of  $\mathcal{T}_\Omega$  and  $\mathcal{I}_Y$ , i.e.,  $\mathcal{T}_{\Omega,Y} = \{T\}$  with  $T = K \times I_m$  for  $K \in \mathcal{T}_\Omega$  and  $I_m \in \mathcal{I}_Y$ . By means of the previous considerations, we define the finite element space  $V_{h,M}$  posed on the tensor product mesh  $\mathcal{T}_{\Omega,Y}$  by

$$V_{h,M} = V_h \otimes V_M = \text{span} \{ v \mid v(x, y) = \mathbf{v}(x)\xi(y), \mathbf{v} \in V_h, \xi \in V_M \} \subset \dot{H}_L^1(C_Y, y^\alpha).$$

Note that each function  $v_h \in V_{h,M}$  vanishes on the lateral boundary of  $C_Y$  and on its top. As a consequence, the extension by zero of  $v_h$  to the semi-infinite cylinder  $C$  belongs to  $\dot{H}_L^1(C, y^\alpha)$ . Without further mention, we consider this type of extension for each  $v_h \in V_{h,M}$  whenever needed.

With the just introduced notation, we define approximations to the solution  $u$  of (2.2) as follows: Find  $u_h \in V_{h,M}$  satisfying

$$(3.1) \quad \int_{C_Y} y^\alpha \nabla u_h \cdot \nabla v_h \, d(x, y) = d_s \langle \mathfrak{f}, \text{tr}_\Omega v_h \rangle_{\mathbb{H}^{-s}(\Omega), \mathbb{H}^s(\Omega)} \quad \forall v_h \in V_{h,M},$$

where we recall that  $\alpha = 1 - 2s$  and  $d_s = 2^\alpha \frac{\Gamma(1-s)}{\Gamma(s)}$ . Note that  $\text{tr}_\Omega u_h = u_h(\cdot, 0)$  will be used as an approximation of  $\mathbf{u}$ .

We distinguish two possible types of discretization in the artificial  $y$  direction, which will be defined in the following two sections.

**3.1. Graded meshes and h-FEM.** In this section, let  $M \in \mathbb{N}$  to be determined later. We set

$$y_m = \left( \frac{m}{M} \right)^{\frac{1}{\mu}} Y \quad \text{for } m = 0, 1, \dots, M \quad \text{and } p = (1, 1, \dots, 1) \in \mathbb{N}^M,$$

where  $\mu \in (0, 1]$  represents the grading parameter in direction  $y$ . Hence, for  $\mu < 1$  the triangulation  $\mathcal{T}_{\Omega,Y}$  is anisotropic. Moreover, due to the choice of the polynomial degree vector  $p$ , the discrete space  $V_{h,M}$  consists of globally continuous and piecewise multilinear functions on  $C_Y$ .

We start with an elementary result regarding the diameter  $h_m$  of the elements  $I_m$ .

LEMMA 3.1. *It holds that*

$$h_1 = M^{-\frac{1}{\mu}} Y \quad \text{and} \quad \frac{2^{\frac{\mu-1}{\mu}}}{\mu} y_m^{1-\mu} Y^\mu M^{-1} \leq h_m \leq \frac{1}{\mu} y_m^{1-\mu} Y^\mu M^{-1} \quad \text{for } m = 2, 3, \dots, M.$$

*Proof.* The first equality is obvious due to the definition of  $y_1$ . For the second, we observe that there exists a  $m_* \in (m - 1, m)$  such that

$$\begin{aligned} h_m &= y_m - y_{m-1} = (m^{\frac{1}{\mu}} - (m-1)^{\frac{1}{\mu}}) Y M^{-\frac{1}{\mu}} \\ &= \frac{1}{\mu} m_*^{\frac{1}{\mu}-1} Y M^{-\frac{1}{\mu}} = \frac{1}{\mu} \left( \left( \frac{m_*}{M} \right)^{\frac{1}{\mu}} Y \right)^{1-\mu} Y^\mu M^{-1} \end{aligned}$$

due to the mean value theorem. Since  $\frac{1}{2}m \leq m - 1 \leq m$  for  $m \geq 2$ , the result follows.  $\square$

Since we consider a discretization with linear polynomials in direction  $y$ , the number of degrees of freedom  $\mathcal{N}_Y$  in  $V_M$  is proportional to  $M$ , i.e., it holds that  $\mathcal{N}_Y = O(M)$ .



**3.2. Geometric meshes and hp-FEM.** For the second possibility of discretization presented here, the mesh in direction  $y$  is chosen as a geometric mesh. That is, for chosen  $\sigma \in (0, 1)$  and  $M \in \mathbb{N}$  to be determined later, the nodes  $y_0, y_1, \dots, y_M$  are given by

$$(3.2) \quad y_0 = 0 \quad \text{and} \quad y_m = \sigma^{M-m}Y \quad \text{for} \quad m = 1, 2, \dots, M.$$

Further, we define  $p = (p_1, p_2, \dots, p_M) \in \mathbb{N}^M$  to be a linear degree vector with slope  $\beta > 0$ . That is, there exists a constant  $c \geq 1$  such that for all  $m = 1, 2, \dots, M$  the following relation is fulfilled:

$$(3.3) \quad 1 + \beta \ln \frac{h_m}{h_1} \leq p_m \leq 1 + c\beta \ln \frac{h_m}{h_1}.$$

We start with collecting some basic results for the discretization considered in this subsection.

LEMMA 3.2. *For a geometric mesh given by (3.2), there holds for  $h_m = |I_m|$*

$$\begin{aligned} h_1 &= y_1 = \sigma^{M-1}Y, \\ h_m &= (1 - \sigma)y_m = (\sigma^{-1} - 1)y_{m-1} \quad \text{for } m = 2, 3, \dots, M, \\ h_m &= (1 - \sigma)\sigma^{1-m}h_1 \quad \text{for } m = 2, 3, \dots, M. \end{aligned}$$

*Proof.* The first equation of the assertion is obvious due to the definition of the geometric mesh. For  $m = 2, 3, \dots, M$ , we obtain

$$\begin{aligned} h_m &= y_m - y_{m-1} = (\sigma^{M-m} - \sigma^{M-m+1})Y \\ &= \begin{cases} \sigma^{M-m}(1 - \sigma)Y = (1 - \sigma)y_m, \\ \sigma^{M-m+1}(\sigma^{-1} - 1)Y = (\sigma^{-1} - 1)y_{m-1}. \end{cases} \end{aligned}$$

As a consequence, we get

$$\frac{h_m}{h_1} = \frac{\sigma^{M-m}(1 - \sigma)Y}{\sigma^{M-1}Y} = (1 - \sigma)\sigma^{1-m}. \quad \square$$

LEMMA 3.3. *For a geometric mesh given by (3.2) and a linear degree vector  $p \in \mathbb{N}^M$  in the sense of (3.3), it holds that  $p_1 = 1$  and there is a constant  $c \geq 1$  such that*

$$1 + \beta(\ln(1 - \sigma) + (1 - m) \ln \sigma) \leq p_m \leq 1 + \beta c(\ln(1 - \sigma) + (1 - m) \ln \sigma)$$

for all  $m = 2, 3, \dots, M$ .

*Proof.* Since  $\ln \frac{h_1}{h_1} = 0$ , there obviously holds that  $p_1 = 1$ . According to Lemma 3.2, we get for  $m = 2, 3, \dots, M - 1$

$$\ln \frac{h_m}{h_1} = \ln((1 - \sigma)\sigma^{1-m}) = \ln(1 - \sigma) + (1 - m) \ln \sigma,$$

which shows the second assertion. □

LEMMA 3.4. *Let  $\mathcal{I}_Y$  be a geometric triangulation of  $(0, Y)$  given by (3.2) and let  $p \in \mathbb{N}^M$  be a linear degree vector as defined in (3.3). Then, for the number of degrees of freedom  $\mathcal{N}_Y$  in  $V_M$ , it holds that*

$$\mathcal{N}_Y = 1 + \sum_{m=1}^M p_m = O(M^2).$$

*Proof.* On each interval  $I_m$  we locally have  $p_m + 1$  degrees of freedom. Moreover, two neighboring intervals always share one degree of freedom. Thus, we have

$$\mathcal{N}_Y = \sum_{m=1}^M (p_m + 1) - (M - 1) = 1 + \sum_{m=1}^M p_m,$$

which proves the assertion having in mind Lemma 3.3. □

**4. Error estimates.** We start with providing a general error estimate between the solutions  $u \in \dot{H}_L^1(C, y^\alpha)$  of (2.2) and  $u_h \in V_{h,M}$  of (3.1) in weighted norms. For a related result we refer to [31, Theorem 4.1].

LEMMA 4.1. *Let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2) and  $u_h \in V_{h,M}$  be the solution of (3.1). Then, it holds that*

$$\|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \leq \min_{v_h \in V_{h,M}} \|y^{\frac{\alpha}{2}} \nabla(u - v_h)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla u\|_{L^2(C \setminus C_Y)}.$$

*Proof.* Testing (2.2) with  $v_h \in V_{h,M}$  and then subtracting (3.1) from this equation yields

$$(4.1) \quad \int_{C_Y} y^\alpha \nabla(u - u_h) \cdot \nabla v_h \, d(x, y) = 0.$$

Based on this, we deduce for all  $v_h \in V_{h,M}$

$$\begin{aligned} & \|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)}^2 \\ &= \int_{C_Y} y^\alpha \nabla(u - u_h) \cdot \nabla(u - v_h) \, d(x, y) + \int_{C \setminus C_Y} y^\alpha \nabla(u - u_h) \cdot \nabla u \, d(x, y) \\ &\leq \|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \{ \|y^{\frac{\alpha}{2}} \nabla(u - v_h)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla u\|_{L^2(C \setminus C_Y)} \}. \end{aligned}$$

Dividing by  $\|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)}$  ends the proof. □

Whereas the last term in the estimate of Lemma 4.1 can be treated by means of Proposition 2.11, we have to estimate the first term. To this end, we introduce the following approximation operators separately for the  $x$  and  $y$  variables.

By  $\pi_x: L^2(\Omega) \rightarrow V_h$ , we denote the  $L^2$  projection with respect to the  $x$  variable on  $\Omega$ . For the interpolation with respect to the  $y$  direction, we consider each interval  $I_m$  separately. For fixed  $m \in \{1, 2, \dots, M\}$ , let  $q \in \mathbb{N}$  and  $y_{m-1} = x_0, x_1, \dots, x_q = y_m$  be the Gauss–Lobatto points in  $I_m$  and let  $l_{i,q}$  denote the corresponding Lagrange polynomials of order  $q$ . Then, we define the Gauss–Lobatto interpolant  $i_q: C^0(\overline{I_m}) \rightarrow \mathcal{P}_q(I_m)$  by

$$(i_q \xi)(y) = \sum_{i=0}^q \xi(x_i) l_{i,q}(y).$$

Further, we define an interpolant  $\tilde{i}_q$  which admits  $(\tilde{i}_q \xi)(y_m) = 0$  given by

$$(\tilde{i}_q \xi)(y) = \sum_{i=0}^{q-1} \xi(x_i) l_{i,q}(y).$$

Based on this, we define the interpolation  $i_y^p: C^0((0, Y]) \rightarrow V_M$  for a linear degree vector  $p = (p_1, p_2, \dots, p_M) \in \mathbb{N}^M$  by

$$(4.2) \quad (i_y^p \xi)|_{I_m} = \begin{cases} \xi(y_1) & \text{on } I_1, \\ i_{p_m} \xi|_{I_m} & \text{on } I_m, \quad m = 2, 3, \dots, M - 1, \\ \tilde{i}_{p_M} \xi|_{I_M} & \text{on } I_M. \end{cases}$$

In particular, it holds that  $i_y^p \xi$  is constant on  $I_1$  and  $(i_y^p \xi)(Y) = 0$ . For any function  $v \in \mathring{H}_L^1(C, y^\alpha)$ , we set

$$(\pi_x v)(\cdot, y) = \pi_x v(\cdot, y) \quad \text{for a.a. } y \in (0, \infty).$$

Moreover, for the solution  $u$  of (2.2), the application of  $i_y^p$  is defined as

$$(4.3) \quad (i_y^p u)(x, \cdot) = \sum_{k=1}^{\infty} u_k \varphi_k(x) i_y^p \psi_{s,k}(\cdot) \quad \text{for a.a. } x \in \Omega,$$

which is well-defined because  $\psi_{s,k} \in C^0(\mathbb{R}_+)$ . Thus, by construction, we have  $\pi_x i_y^p u \in V_{h,M}$ .

For the first term on the right-hand side of the estimate in Lemma 4.1, we have the following result.

LEMMA 4.2. *Let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Then, it holds that*

$$\min_{v_h \in V_{h,M}} \|y^{\frac{\alpha}{2}} \nabla(u - v_h)\|_{L^2(C_Y)} \leq \|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)} + c \|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(C_Y)}.$$

*Proof.* First, we set  $v_h = \pi_x i_y^p u \in V_{h,M}$ . Then, by introducing  $\pi_x u$  as an intermediate function, we deduce

$$\min_{v_h \in V_{h,M}} \|y^{\frac{\alpha}{2}} \nabla(u - v_h)\|_{L^2(C_Y)} \leq \|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla \pi_x(u - i_y^p u)\|_{L^2(C_Y)}.$$

According to the definition of  $\pi_x$ , we have  $\partial_y \pi_x(u - i_y^p u) = \pi_x \partial_y(u - i_y^p u)$  almost everywhere in  $C_Y$ . As a consequence, we deduce by well-known stability estimates for the  $L^2$  projection  $\pi_x$

$$\begin{aligned} \|y^{\frac{\alpha}{2}} \nabla \pi_x(u - i_y^p u)\|_{L^2(C_Y)} &= \int_0^Y y^\alpha \int_\Omega \{|\nabla_x \pi_x(u - i_y^p u)|^2 + |\partial_y \pi_x(u - i_y^p u)|^2\} dx dy \\ &= \int_0^Y y^\alpha \int_\Omega \{|\nabla_x \pi_x(u - i_y^p u)|^2 + |\pi_x \partial_y(u - i_y^p u)|^2\} dx dy \\ &\leq c \int_0^Y y^\alpha \int_\Omega \{|\nabla_x(u - i_y^p u)|^2 + |\partial_y(u - i_y^p u)|^2\} dx dy, \end{aligned}$$

which shows the assertion. □

LEMMA 4.3. *For  $f \in \mathbb{H}^{1-s}(\Omega)$ , let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Then, it holds that*

$$\|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)} \leq ch_\Omega \|f\|_{\mathbb{H}^{1-s}(\Omega)}.$$

*Proof.* Analogously to the foregoing proof, by classical estimates for the  $L^2$  projection  $\pi_x$ , we obtain

$$\begin{aligned} \|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)}^2 &= \int_0^Y y^\alpha \int_\Omega \{|\nabla_x(u - \pi_x u)|^2 + |\partial_y(u - \pi_x u)|^2\} dx dy \\ &= \int_0^Y y^\alpha \int_\Omega \{|\nabla_x(u - \pi_x u)|^2 + |\partial_y u - \pi_x(\partial_y u)|^2\} dx dy \\ &\leq ch_\Omega^2 \int_0^Y y^\alpha \int_\Omega \{|\nabla_x^2 u|^2 + |\nabla_x \partial_y u|^2\} dx dy \\ &= ch_\Omega^2 \{ \|y^{\frac{\alpha}{2}} \nabla_x^2 u\|_{L^2(C_Y)}^2 + \|y^{\frac{\alpha}{2}} \partial_y \nabla_x u\|_{L^2(C_Y)}^2 \}. \end{aligned}$$

Then, Proposition 2.6 yields the assertion. □

*Remark 4.4.* Only for the validity of Lemma 4.3, we assume the convexity of the domain; see also Proposition 2.6 and Remark 2.7. In nonconvex domains, using the  $H^t(\Omega)$  regularity of  $u(\cdot, y)$ , we simply obtain a convergence rate of  $t - 1 \in [0, 1]$ , which then dominates the overall error. As a remedy, one could use the regularity of  $u(\cdot, y)$  in  $\Omega$ -weighted Sobolev spaces and, for instance, graded meshes, which are appropriately refined to resolve the singularities in  $\Omega$ , in order to retain a convergence rate of one; see, e.g., [33, 6].

Next, we are concerned with estimates for the second term in Lemma 4.2. To this end, we consider the interpolation error on each subinterval  $I_m \in \mathcal{I}_Y$ . Employing the decomposition from Proposition 2.8 and the definition of the eigenfunctions  $\varphi_k$ , we obtain

$$\begin{aligned}
 & \|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2 \\
 &= \int_{I_m} y^\alpha \int_\Omega \{|\nabla_x(u - i_y^p u)|^2 + |\partial_y(u - i_y^p u)|^2\} dx dy \\
 (4.4) \quad &= \int_{I_m} y^\alpha \sum_{k=1}^\infty \mathbf{u}_k^2 \{ \lambda_k (\psi_{s,k} - i_y^p \psi_{s,k})^2 + ((\psi_{s,k} - i_y^p \psi_{s,k})')^2 \} dy \\
 &= \sum_{k=1}^\infty \mathbf{u}_k^2 \{ \lambda_k \|y^{\frac{\alpha}{2}} (\psi_{s,k} - i_y^p \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}} (\psi_{s,k} - i_y^p \psi_{s,k})'\|_{L^2(I_m)}^2 \}.
 \end{aligned}$$

By using this identity in the following two subsections, we will estimate the terms

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2, \quad i = 1, 2, \dots, M,$$

for the two types of triangulations  $\mathcal{I}_Y$  and polynomial spaces  $V_M$  introduced in sections 3.1 and 3.2. We already note that we will track the dependence on  $Y$  in the corresponding estimates explicitly since as a last step  $Y$  will be chosen  $h$ -dependent.

**4.1. Graded meshes and  $h$ -FEM.** In the present section, we will mainly recover the results of [31]. The reason for doing this is twofold. First, we are able to slightly improve the grading condition from  $\mu < \frac{2}{3}s$  (in our notation) of [31, section 5.2] to  $\mu < s$ . However, the main reason to analyze the  $h$ -FEM on graded meshes before developing the analysis for the considered  $hp$ -method is that the techniques we use are almost identical for both cases, but the details are simpler for  $h$ -FEM, of course.

As announced in the preface of this section, we are primarily concerned with estimates for (4.4) for the discretization defined in section 3.1. By these and the foregoing considerations, we will obtain the main result for the  $h$ -FEM on graded meshes. For simplicity, in this subsection, we will write  $i_y u$  for  $i_y^p u$ , since we have  $p = (1, 1, \dots, 1)$  here.

**LEMMA 4.5** (estimate on  $I_1$ ). *For  $\mathfrak{f} \in L^2(\Omega)$ , let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2) and let  $M \geq h_\Omega^{-1}$ . Then, it holds that*

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(\Omega \times I_1)}^2 \leq ch_\Omega^{\frac{2s}{\mu}} Y^{2s} \|\mathfrak{f}\|_{L^2(\Omega)}^2.$$

*Proof.* First, we observe that the interpolant  $i_y \psi_{s,k}$  is constant on  $I_1$ . By its definition (4.2), it holds that  $(i_y \psi_{s,k})|_{I_1} = \psi_{s,k}(y_1)$ . Integration by parts and noting that  $y^{\alpha+1}(\psi_{s,k} - \psi_{s,k}(y_1))^2|_0^{y_1} = 0$  yields

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)}^2 &= \int_{I_1} y^\alpha (\psi_{s,k} - \psi_{s,k}(y_1))^2 dy \\ &= -\frac{2}{\alpha + 1} \int_{I_1} y^{\alpha+1} (\psi_{s,k} - \psi_{s,k}(y_1)) \psi'_{s,k} dy \\ &\leq \frac{2}{\alpha + 1} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)} \|y^{\frac{\alpha}{2}+1} \psi'_{s,k}\|_{L^2(I_1)}. \end{aligned}$$

Then, dividing by  $\|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_y \psi_{s,k})\|_{L^2(I_1)}$  implies

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)} \leq c \|y^{\frac{\alpha}{2}+1} \psi'_{s,k}\|_{L^2(I_1)}$$

with  $c = \frac{1}{1-s}$ . By means of Corollary 2.9 with  $n = 1$  and  $r = 1$ , we obtain

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)} \leq c \|y^{\frac{\alpha}{2}+1} \psi'_{s,k}\|_{L^2(I_1)} \leq c \lambda_k^{s-\frac{1}{2}} \|y^{s-\frac{1}{2}}\|_{L^2(I_1)}.$$

Hence, we get by Lemma 3.1 together with the assumption on  $M$  that

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)}^2 \leq c \lambda_k^{2s} h_1^{2s} \leq c \lambda_k^{2s} h_\Omega^{\frac{2s}{\mu}} Y^{2s}.$$

In a similar fashion, we obtain by Corollary 2.9 with  $n = 1$  and  $r = 0$  the relation

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \psi_{s,k}(y_1))'\|_{L^2(I_1)}^2 &= \|y^{\frac{\alpha}{2}} \psi'_{s,k}\|_{L^2(I_1)}^2 \leq c \lambda_k^{2s} \|y^{s-\frac{1}{2}}\|_{L^2(I_1)}^2 \leq c \lambda_k^{2s} h_1^{2s} \\ &\leq c \lambda_k^{2s} h_\Omega^{\frac{2s}{\mu}} Y^{2s}, \end{aligned}$$

where we again used Lemma 3.1 in the last step.

The previous estimates together with (4.4) and Proposition 2.1 yield

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(\Omega \times I_1)}^2 \leq c h_\Omega^{\frac{2s}{\mu}} Y^{2s} \sum_{k=1}^\infty \lambda_k^{2s} u_k^2 = c h_\Omega^{\frac{2s}{\mu}} Y^{2s} \sum_{k=1}^\infty f_k^2,$$

which implies the assertion. □

LEMMA 4.6 (estimates on  $I_m$  for  $2 \leq m \leq M - 1$ ). For  $f \in L^2(\Omega)$ , let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $\mu \neq s$  and  $M \geq h_\Omega^{-1}$ . Then, it holds that

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(\Omega \times I_m)}^2 \leq c h_\Omega^2 Y^{2\mu} \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\} \|f\|_{L^2(\Omega)}^2.$$

*Proof.* For  $m \geq 2$ , we have that  $y_{m-1} \leq y_m \leq 2^{\frac{1}{\mu}} y_{m-1}$ . On  $I_m$ , there holds  $i_y \psi_{s,k} = i_1 \psi_{s,k}$  such that we conclude with standard estimates for the linear Lagrange interpolant  $i_1$ , Lemma 3.1, and the assumption on  $M$  that

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_m)}^2 &\leq c y_m^\alpha \|\psi_{s,k} - i_1 \psi_{s,k}\|_{L^2(I_m)}^2 \leq c y_m^\alpha h_m^2 \|\psi'_{s,k}\|_{L^2(I_m)}^2 \\ &\leq c y_m^{\alpha+2-2\mu} h_\Omega^2 Y^{2\mu} \|\psi'_{s,k}\|_{L^2(I_m)}^2 \leq c h_\Omega^2 Y^{2\mu} \|y^{\frac{\alpha}{2}+1-\mu} \psi'_{s,k}\|_{L^2(I_m)}^2. \end{aligned}$$

By using Corollary 2.9 with  $n = 1$  and  $r = 1$ , this implies

$$\begin{aligned} \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_m)}^2 &\leq c \lambda_k^{2s} h_\Omega^2 Y^{2\mu} \|y^{s-\mu-\frac{1}{2}}\|_{L^2(I_m)}^2 \\ &= c \lambda_k^{2s} h_\Omega^2 Y^{2\mu} \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\}, \end{aligned}$$

where we used that  $\mu \neq s$ . Similarly, using Corollary 2.9 with  $n = 1$  and  $r = 0$ , we obtain for the term involving the derivative

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1\psi_{s,k})'\|_{L^2(I_m)}^2 &\leq ch_\Omega^2 Y^{2\mu} \|y^{\frac{\alpha}{2}+1-\mu}\psi_{s,k}''\|_{L^2(I_m)}^2 \\ &\leq c\lambda_k^{2s} h_\Omega^2 Y^{2\mu} \|y^{s-\mu-\frac{1}{2}}\|_{L^2(I_m)}^2 \\ &= c\lambda_k^{2s} h_\Omega^2 Y^{2\mu} \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\}. \end{aligned}$$

The previous estimates in combination with (4.4) yield

$$\|y^{\frac{\alpha}{2}}\nabla(u - i_y u)\|_{L^2(\Omega \times I_m)}^2 \leq ch_\Omega^2 Y^{2\mu} \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\} \sum_{k=1}^\infty \lambda_k^{2s} u_k^2.$$

Finally, applying Proposition 2.1, we get

$$\|y^{\frac{\alpha}{2}}\nabla(u - i_y u)\|_{L^2(\Omega \times I_m)}^2 \leq ch_\Omega^2 Y^{2\mu} \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\} \sum_{k=1}^\infty f_k^2,$$

which states the assertion. □

LEMMA 4.7 (estimate on  $I_M$ ). For  $f \in L^2(\Omega)$ , let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $\mu \neq s$ ,

$$h_\Omega^{-1} \geq M \geq h_\Omega^{-1} \quad \text{and} \quad Y \geq \max\left(\frac{3|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right).$$

Then, it holds that

$$\|y^{\frac{\alpha}{2}}\nabla(u - i_y u)\|_{L^2(\Omega \times I_M)}^2 \leq ch_\Omega^2 \left(Y^{2\mu} \{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\} + 1\right) \|f\|_{L^2(\Omega)}^2.$$

*Proof.* We recall that  $Y = y_M$  and  $i_y \psi_{s,k} = \tilde{i}_1 \psi_{s,k}$  on  $I_M$ . We introduce the Lagrange interpolation  $i_1$  on  $I_M$  as an intermediate function such that

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_1 \psi_{s,k})\|_{L^2(I_M)} &\leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_M)} + \|y^{\frac{\alpha}{2}}(i_1 \psi_{s,k} - \tilde{i}_1 \psi_{s,k})\|_{L^2(I_M)} \\ &= \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_M)} + \psi_{s,k}(Y) \|y^{\frac{\alpha}{2}} l_{1,1}\|_{L^2(I_M)} \\ &\leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_M)} + cY^{\frac{\alpha+1}{2}} \psi_{s,k}(Y), \end{aligned}$$

where we used that  $\|l_{1,1}\|_{L^\infty(I_M)} = \|\frac{y-y_{M-1}}{h_M}\|_{L^\infty(I_M)} = 1$  in the last step. As in the proof of Lemma 4.6, we deduce

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1 \psi_{s,k})\|_{L^2(I_M)}^2 \leq c\lambda_k^{2s} h_\Omega^2 Y^{2\mu} \{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\}.$$

Since  $Y \geq 1$  by assumption, we obtain using Corollary 2.10 with  $r_1 = \frac{\alpha+1}{2} = 1 - s$  together with the monotonicity of  $e^{-\sqrt{\lambda_k} y}$

$$Y^{\frac{\alpha+1}{2}} \psi_{s,k}(Y) \leq c\lambda_k^{\frac{s-1}{2}} e^{-\frac{\sqrt{\lambda_k}}{2} Y} \leq c\lambda_k^{s-\frac{1}{2}} e^{-\frac{\sqrt{\lambda_1}}{2} Y},$$

where we notice that  $(\lambda_k)_{k \in \mathbb{N}}$  is a nondecreasing sequence. Combining the previous results yields

$$(4.5) \quad \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_1 \psi_{s,k})\|_{L^2(I_M)}^2 \leq c\lambda_k^{2s} \left(h_\Omega^2 Y^{2\mu} \{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\} + e^{-\sqrt{\lambda_1} Y}\right).$$

Similarly, we deduce

$$\begin{aligned} & \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_1\psi_{s,k})'\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1\psi_{s,k})'\|_{L^2(I_M)} + \|y^{\frac{\alpha}{2}}(i_1\psi_{s,k} - \tilde{i}_1\psi_{s,k})'\|_{L^2(I_M)} \\ & = \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1\psi_{s,k})'\|_{L^2(I_M)} + \psi_{s,k}(Y)\|y^{\frac{\alpha}{2}}l'_{1,1}\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1\psi_{s,k})'\|_{L^2(I_M)} + ch_M^{-\frac{1}{2}}Y^{\frac{\alpha}{2}}\psi_{s,k}(Y), \end{aligned}$$

where we used that  $\|l'_{1,1}\|_{L^\infty(I_M)} = \|h_M^{-1}\|_{L^\infty(I_M)} = h_M^{-1}$  in the last step. The first term can again be estimated as in the proof of Lemma 4.6 such that

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_1\psi_{s,k})'\|_{L^2(I_m)}^2 \leq c\lambda_k^{2s}h_\Omega^2Y^{2\mu}\{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\}.$$

Employing Corollary 2.10 with  $r_1 = \frac{\alpha}{2} = \frac{1}{2} - s$  together with the monotonicity of  $e^{-\sqrt{\lambda_k}y}$  yields for  $Y \geq 1$

$$h_M^{-\frac{1}{2}}Y^{\frac{\alpha}{2}}\psi_{s,k}(Y) \leq ch_M^{-\frac{1}{2}}\lambda_k^{\frac{s}{2}-\frac{1}{4}}e^{-\frac{\sqrt{\lambda_k}}{2}Y} \leq ch_M^{-\frac{1}{2}}\lambda_k^s e^{-\frac{\sqrt{\lambda_1}}{2}Y},$$

where we used once again that the sequence  $(\lambda_k)_{k \in \mathbb{N}}$  is nondecreasing. Due to the previous results, we arrive at

(4.6)

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_1\psi_{s,k})'\|_{L^2(I_M)}^2 \leq c\lambda_k^{2s}\left(h_\Omega^2Y^{2\mu}\{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\} + h_M^{-1}e^{-\sqrt{\lambda_1}Y}\right).$$

By combining (4.4), (4.5), and (4.6), we obtain

$$\begin{aligned} & \|y^{\frac{\alpha}{2}}\nabla(u - i_yu)\|_{L^2(\Omega \times I_M)}^2 \\ & \leq c\left(h_\Omega^2Y^{2\mu}\{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\} + \{1 + h_M^{-1}\}e^{-\sqrt{\lambda_1}Y}\right)\sum_{k=1}^\infty \lambda_k^{2s}u_k^2 \\ & = c\left(h_\Omega^2Y^{2\mu}\{Y^{2(s-\mu)} - y_{M-1}^{2(s-\mu)}\} + \{1 + h_M^{-1}\}e^{-\sqrt{\lambda_1}Y}\right)\|f\|_{L^2(\Omega)}^2, \end{aligned}$$

where we used  $\sum_{k=1}^\infty \lambda_k^{2s}u_k^2 = \sum_{k=1}^\infty f_k^2$  (see Proposition 2.1) and the definition of  $\|f\|_{L^2(\Omega)}^2$ . According to Lemma 3.1, there holds  $h_M^{-1} \leq cY^{\mu-1}Y^{-\mu}M = cY^{-1}M$ . Since

$$Y \geq \frac{3|\ln h_\Omega|}{\sqrt{\lambda_1}} \quad \text{and} \quad M \leq 2h_\Omega^{-1}$$

by assumption, we obtain

$$\{1 + h_M^{-1}\}e^{-\sqrt{\lambda_1}Y} \leq h_\Omega^3 + ch_\Omega^2|\ln h_\Omega|^{-1} \leq ch_\Omega^2,$$

which ends the proof. □

**COROLLARY 4.8.** For  $f \in L^2(\Omega)$ , let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $\mu < s$ ,

$$2h_\Omega^{-1} \geq M \geq h_\Omega^{-1} \quad \text{and} \quad 2\max\left(\frac{3|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right) \geq Y \geq \max\left(\frac{3|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right).$$

Then, it holds that

$$\|y^{\frac{\alpha}{2}}\nabla(u - i_yu)\|_{L^2(C_Y)} \leq ch_\Omega|\ln h_\Omega|^s\|f\|_{L^2(\Omega)}.$$

*Proof.* By Lemmas 4.5, 4.6, and 4.7, we obtain

$$\begin{aligned} & \|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(C_Y)}^2 \\ &= \sum_{m=1}^M \|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(\Omega \times I_m)}^2 \\ &\leq \left( ch_{\Omega}^{\frac{2s}{\mu}} Y^{2s} + ch_{\Omega}^2 Y^{2\mu} \sum_{m=2}^M \{y_m^{2(s-\mu)} - y_{m-1}^{2(s-\mu)}\} + ch_{\Omega}^2 \right) \|f\|_{L^2(\Omega)}^2 \\ &\leq ch_{\Omega}^2 \left( Y^{2s} - Y^{2\mu} y_1^{2(s-\mu)} + 1 \right) \|f\|_{L^2(\Omega)}^2 \leq ch_{\Omega}^2 |\ln h_{\Omega}|^{2s} \|f\|_{L^2(\Omega)}^2, \end{aligned}$$

where we have used  $\mu < s$  and the upper bound on  $Y$ . □

Now, we are able to state the main result for this subsection analyzing the  $h$ -FEM on graded meshes.

**THEOREM 4.9.** *For  $f \in \mathbb{H}^{1-s}(\Omega)$ , let  $u \in \mathbb{H}^s(\Omega)$  and  $u \in \dot{H}_L^1(C, y^{\alpha})$  be the solutions of (1.1) and (2.2), respectively, and let  $u_h \in V_{h,M}$  be the solution of (3.1). Moreover, let  $\mu < s$ ,*

$$(4.7) \quad 2h_{\Omega}^{-1} \geq M \geq h_{\Omega}^{-1} \quad \text{and} \quad 2 \max\left(\frac{3|\ln h_{\Omega}|}{\sqrt{\lambda_1}}, 1\right) \geq Y \geq \max\left(\frac{3|\ln h_{\Omega}|}{\sqrt{\lambda_1}}, 1\right).$$

*Then, it holds that*

$$\|u - \text{tr}_{\Omega} u_h\|_{\mathbb{H}^s(\Omega)} \leq c \|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \leq ch_{\Omega} |\ln h_{\Omega}|^s \|f\|_{\mathbb{H}^{1-s}(\Omega)}.$$

*Proof.* The first inequality of the assertion is due to Propositions 2.4 and 2.5. Using Lemmas 4.1 and 4.2, we get

$$\begin{aligned} & \|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \\ & \leq \|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla u\|_{L^2(C \setminus C_Y)}. \end{aligned}$$

The three terms on the right-hand side are estimated in Lemma 4.3, Corollary 4.8, and Proposition 2.11. Hence, we get

$$\|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \leq ch_{\Omega} \|f\|_{\mathbb{H}^{1-s}(\Omega)} + ch_{\Omega} |\ln h_{\Omega}|^s \|f\|_{L^2(\Omega)} + ce^{-\frac{\sqrt{\lambda_1}}{2} Y} \|f\|_{\mathbb{H}^{-s}(\Omega)}.$$

Then, the lower bound on  $Y$  yields  $e^{-\frac{\sqrt{\lambda_1}}{2} Y} \leq ch_{\Omega}^{\frac{3}{2}} \leq ch_{\Omega}$ , which implies the assertion. □

**THEOREM 4.10.** *The total number degrees of freedom  $\mathcal{N}_{\Omega,Y}$  in  $V_{h,M}$  to achieve the order of convergence given in Theorem 4.9 behaves like*

$$\mathcal{N}_{\Omega,Y} = O(\mathcal{N}_{\Omega}^{1+\frac{1}{d}}),$$

where  $d$  denotes the dimension of  $\Omega$ .

*Proof.* For the number of degrees of freedom  $\mathcal{N}_{\Omega,Y}$  of the discretization considered in this section, it holds that  $\mathcal{N}_{\Omega,Y} = \mathcal{N}_{\Omega} \mathcal{N}_Y = \mathcal{N}_{\Omega} M$ . Then, the assertion follows from  $M = O(h_{\Omega}^{-1}) = O(\mathcal{N}_{\Omega}^{1/d})$ . □

*Remark 4.11.* We notice that the factor 2 in each of the upper bounds of (4.7) is chosen rather arbitrarily. Any constant factor greater than one (independent of  $h_{\Omega}$ ) is sufficient for the validity of the error estimates and for the validity of the result about the complexity. As a consequence, once one has a guess or an approximation for  $\lambda_1$ , it is easily possible to choose  $Y$ .



**4.2. Geometric meshes and hp-FEM.** In this section, we derive discretization error estimates for the *hp*-method described in section 3.2, which results in a slightly improved rate of convergence ( $h_\Omega$  versus  $h_\Omega |\ln h_\Omega|^s$ ) compared to the previous subsection. However, we will have a drastically reduced computational complexity of  $O(\mathcal{N}_\Omega (\ln \mathcal{N}_\Omega)^2)$  compared to  $O(\mathcal{N}_\Omega^{1+1/d})$  in terms of the number of degrees of freedom  $\mathcal{N}_\Omega$ . To this end, we fix neither the number of elements  $M$  in direction  $y$  nor the slope  $\beta$  of the linear degree vector  $p$  yet. These will be set below. As before, we start with estimates for  $\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2$  based on (4.4).

LEMMA 4.12 (estimate on  $I_1$ ). For  $\mathfrak{f} \in L^2(\Omega)$ , let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2) and let

$$M \geq \frac{(1 + \varepsilon) |\ln h_\Omega|}{s |\ln \sigma|}$$

for some  $\varepsilon \geq 0$ . Then, it holds that

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_1)}^2 \leq ch_\Omega^{2+2\varepsilon} Y^{2s} \|\mathfrak{f}\|_{L^2(\Omega)}^2.$$

*Proof.* Notice that  $i_y^p \psi_{s,k} = \psi_{s,k}(y_1)$  on the first interval  $I_1$  as in the previous section. Thus, as in the proof of Lemma 4.5 but using

$$h_1 = \sigma^{M-1} Y \leq ch_\Omega^{\frac{1+\varepsilon}{s}} Y$$

from Lemma 3.2 and the assumption on  $M$ , we get

$$\begin{aligned} \lambda_k \|y^{\frac{\alpha}{2}} (\psi_{s,k} - \psi_{s,k}(y_1))\|_{L^2(I_1)}^2 &\leq c\lambda_k^{2s} h_1^{2s} \leq c\lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}, \\ \|y^{\frac{\alpha}{2}} (\psi_{s,k} - \psi_{s,k}(y_1))'\|_{L^2(I_1)}^2 &\leq c\lambda_k^{2s} h_1^{2s} \leq c\lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}. \end{aligned}$$

Hence, we obtain

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_1)}^2 \leq ch_\Omega^{2+2\varepsilon} Y^{2s} \sum_{k=1}^\infty \lambda_k^{2s} u_k^2.$$

As in the proof of Lemma 4.5, this yields the assertion. □

In order to derive estimates on  $I_m$  for  $2 \leq m \leq M - 1$ , we recall the following result, which is a direct consequence of [27, Lemma 3.2.6].

PROPOSITION 4.13. Let  $w$  be analytic on  $\hat{I} = (0, 1)$  and satisfy for some  $c_w, \delta > 0$  the estimate

$$\|w^{(n)}\|_{L^\infty(\hat{I})} \leq c_w \delta^n n! \quad \forall n \in \mathbb{N}.$$

Then, there are constants  $c, b > 0$  depending only on  $\delta$  such that the Gauss-Lobatto interpolant  $i_q w$  of degree  $q \in \mathbb{N}$  on  $\hat{I}$  satisfies

$$\|w - i_q w\|_{L^\infty(\hat{I})} + \|(w - i_q w)'\|_{L^\infty(\hat{I})} \leq cc_w e^{-bq}.$$

LEMMA 4.14 (estimates on  $I_m$  for  $2 \leq m \leq M - 1$ ). For  $\mathfrak{f} \in L^2(\Omega)$ , let  $u \in \dot{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $p \in \mathbb{N}^M$  be a linear degree vector as in (3.3) with some  $\beta > 0$  and let

$$M \geq \frac{(1 + \varepsilon) |\ln h_\Omega|}{\min(s, \beta b) |\ln \sigma|}$$

for some  $\varepsilon \geq 0$ , where  $b > 0$  is a constant depending on  $\sigma$  only. Then, it holds that

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2 \leq ch_\Omega^{2+2\varepsilon} Y^{2s} \|\mathfrak{f}\|_{L^2(\Omega)}^2.$$

*Proof.* For  $m \geq 2$  it holds that  $y_{m-1} \leq y_m \leq \sigma^{-1}y_{m-1}$ . By transforming to the reference element  $\hat{I} = (0, 1)$ , we obtain for  $(i_y^p \psi_{s,k})|_{I_m} = i_{p_m} \psi_{s,k}|_{I_m}$  that

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)} &\leq cy_{m-1}^{\frac{\alpha}{2}} h_m^{\frac{1}{2}} \|\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k}\|_{L^2(\hat{I})} \\ &\leq cy_{m-1}^{\frac{\alpha}{2}} h_m^{\frac{1}{2}} \|\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k}\|_{L^\infty(\hat{I})}. \end{aligned}$$

By means of Corollary 2.9, applied with  $r = 1$ , and Lemma 3.2, we have

$$\begin{aligned} \|\hat{\psi}_{s,k}^{(n)}\|_{L^\infty(\hat{I})} &= h_m^n \|\psi_{s,k}^{(n)}\|_{L^\infty(I_m)} \leq c\lambda_k^{s-\frac{1}{2}} h_m^n y_{m-1}^{2s-1-n} 8^n n! \\ &\leq c\lambda_k^{s-\frac{1}{2}} y_{m-1}^{2s-1} (8(\sigma^{-1} - 1))^n n!. \end{aligned}$$

Moreover, due to well-known series representations of  $K_s$  from [1, eqs. 9.6.2 and 9.6.10], we directly conclude that  $\hat{\psi}_{s,k}$  is analytic on  $\hat{I}$ . Hence, Proposition 4.13 implies with  $\delta = 8(\sigma^{-1} - 1)$  that

$$\|\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k}\|_{L^\infty(\hat{I})} \leq c\lambda_k^{s-\frac{1}{2}} y_{m-1}^{2s-1} e^{-bp_m}.$$

Then, we get by Lemma 3.2

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)} \leq cy_{m-1}^{s-\frac{1}{2}} h_m^{\frac{1}{2}} \lambda_k^{s-\frac{1}{2}} e^{-bp_m} \leq ch_m^s \lambda_k^{s-\frac{1}{2}} e^{-bp_m}.$$

Analogously, we obtain

$$\begin{aligned} \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)} &\leq cy_{m-1}^{\frac{\alpha}{2}} h_m^{-\frac{1}{2}} \|(\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k})'\|_{L^2(\hat{I})} \\ &\leq cy_{m-1}^{\frac{\alpha}{2}} h_m^{-\frac{1}{2}} \|(\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k})'\|_{L^\infty(\hat{I})}. \end{aligned}$$

By means of Corollary 2.9 with  $r = 0$  and Lemma 3.2, we have

$$\|\hat{\psi}_{s,k}^{(n)}\|_{L^\infty(\hat{I})} = h_m^n \|\psi_{s,k}^{(n)}\|_{L^\infty(I_m)} \leq c\lambda_k^s h_m^n y_{m-1}^{2s-n} 8^n n! \leq c\lambda_k^s y_{m-1}^{2s} (8(\sigma^{-1} - 1))^n n!.$$

Consequently, Proposition 4.13 yields with  $\delta = 8(\sigma^{-1} - 1)$  that

$$\|(\hat{\psi}_{s,k} - \hat{i}_{p_m} \hat{\psi}_{s,k})'\|_{L^\infty(\hat{I})} \leq c\lambda_k^s y_{m-1}^{2s} e^{-bp_m}.$$

This implies

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)} \leq cy_{m-1}^{s+\frac{1}{2}} h_m^{-\frac{1}{2}} \lambda_k^s e^{-bp_m} \leq c\lambda_k^s h_m^s e^{-bp_m}.$$

Collecting the previous results yields

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \leq c\lambda_k^{2s} h_m^{2s} e^{-2bp_m}.$$

Relation (3.3) implies

$$e^{-2bp_m} \leq ch_1^{2\beta b} h_m^{-2\beta b}.$$

Thus, we deduce

$$(4.8) \quad \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \leq c\lambda_k^{2s} h_m^{2(s-\beta b)} h_1^{2\beta b}.$$

Let us now distinguish two cases:

- $s \leq \beta b$ : Since  $h_m \geq h_1$ , we then obtain

$$\begin{aligned} \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \\ \leq c\lambda_k^{2s} h_1^{2(s-\beta b)} h_1^{2\beta b} = c\lambda_k^{2s} h_1^{2s}. \end{aligned}$$

As before, the relation  $M \geq \frac{(1+\varepsilon)|\ln h_\Omega|}{s|\ln \sigma|}$  together with Lemma 3.2 implies  $h_1 \leq ch_\Omega^{\frac{1+\varepsilon}{s}} Y$ . Hence, we get

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \leq c\lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}.$$

- $s > \beta b$ : With  $h_m \leq Y$ , we get from (4.8) that

$$\begin{aligned} \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \\ \leq c\lambda_k^{2s} Y^{2(s-\beta b)} h_1^{2\beta b}. \end{aligned}$$

Similarly as before, the relation  $M \geq \frac{(1+\varepsilon)|\ln h_\Omega|}{\beta b|\ln \sigma|}$  together with Lemma 3.2 implies  $h_1 \leq ch_\Omega^{\frac{1+\varepsilon}{\beta b}} Y$ . Thus, we get

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})\|_{L^2(I_m)}^2 + \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_m} \psi_{s,k})'\|_{L^2(I_m)}^2 \leq c\lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}.$$

The previous results in combination with (4.4) imply

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2 \leq ch_\Omega^{2+2\varepsilon} Y^{2s} \sum_{k=1}^{\infty} \lambda_k^{2s} u_k^2.$$

Finally, applying Proposition 2.1 yields the assertion. □

LEMMA 4.15 (estimate on  $I_M$ ). For  $f \in L^2(\Omega)$ , let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $p \in \mathbb{N}^M$  be a linear degree vector as in (3.3) with some  $\beta > 0$  and let

$$2 \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|} \geq M \geq \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|} \quad \text{and} \quad Y \geq \max\left(\frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right)$$

for some  $\varepsilon \geq 0$ , where  $b > 0$  is a constant depending on  $\sigma$  only. Then, it holds that

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_M)}^2 \leq c(h_\Omega^{2+2\varepsilon} Y^{2s} + h_\Omega^2) \|f\|_{L^2(\Omega)}^2.$$

*Proof.* We proceed as in the proof of Lemma 4.7 and recall that  $Y = y_M = \sigma^{-1}y_{M-1}$  and  $h_M = (1 - \sigma)Y = (\sigma^{-1} - 1)y_{M-1}$ . Moreover, according to [19], the Lagrange basis functions  $l_{i,q}$  of order  $q \in \mathbb{N}$  on  $I_M$  have the property

$$(4.9) \quad \|l_{i,q}\|_{L^\infty(I_M)} \leq 1 \quad \text{for } i = 0, 1, \dots, q.$$

As a consequence, we obtain by means of an inverse inequality (see, e.g., [27, Lemma 3.2.2])

$$(4.10) \quad \|l'_{i,q}\|_{L^\infty(I_M)} \leq ch_M^{-1} q^2 \|l_{i,q}\|_{L^\infty(I_M)} \leq cq^2 Y^{-1}.$$

Noting that  $i_y^p \psi_{s,k} = \tilde{i}_{p_M} \psi_{s,k}$  on  $I_M$ , we introduce the Gauss–Lobatto interpolant  $i_{p_M} \psi_{s,k}$  on  $I_M$  as an intermediate function such that

$$\begin{aligned} & \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})\|_{L^2(I_M)} + \|y^{\frac{\alpha}{2}}(i_{p_M} \psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})\|_{L^2(I_M)} \\ & = \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})\|_{L^2(I_M)} + \psi_{s,k}(Y) \|y^{\frac{\alpha}{2}} l_{p_M, p_M}\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})\|_{L^2(I_M)} + Y^{\frac{\alpha+1}{2}} \psi_{s,k}(Y), \end{aligned}$$

where we used (4.9) in the last step. As in the proof of Lemma 4.14, we deduce for  $M \geq \frac{(1+\varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|}$  that

$$\lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})\|_{L^2(I_M)}^2 \leq c \lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}.$$

Since  $Y \geq 1$  by assumption, we obtain using Corollary 2.10 with  $r_1 = \frac{\alpha+1}{2} = 1 - s$  together with the monotonicity of  $e^{-\sqrt{\lambda_k} y}$

$$Y^{\frac{\alpha+1}{2}} \psi_{s,k}(Y) \leq c \lambda_k^{\frac{s-1}{2}} e^{-\frac{\sqrt{\lambda_k}}{2} Y} \leq c \lambda_k^{s-\frac{1}{2}} e^{-\frac{\sqrt{\lambda_1}}{2} Y},$$

where we notice that  $(\lambda_k)_{k \in \mathbb{N}}$  is a nondecreasing sequence. Combining the previous results yields

$$(4.11) \quad \lambda_k \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})\|_{L^2(I_M)}^2 \leq c \lambda_k^{2s} \left( h_\Omega^{2+2\varepsilon} Y^{2s} + e^{-\sqrt{\lambda_1} Y} \right).$$

Similarly, we deduce by means of (4.10)

$$\begin{aligned} & \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})'\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})'\|_{L^2(I_M)} + \|y^{\frac{\alpha}{2}}(i_{p_M} \psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})'\|_{L^2(I_M)} \\ & = \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})'\|_{L^2(I_M)} + \psi_{s,k}(Y) \|y^{\frac{\alpha}{2}} l'_{p_M, p_M}\|_{L^2(I_M)} \\ & \leq \|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})'\|_{L^2(I_M)} + c p_M^2 Y^{-2} Y^{\frac{\alpha+1}{2}+1} \psi_{s,k}(Y). \end{aligned}$$

Using  $M \geq \frac{(1+\varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|}$ , the first term can again be estimated as in the proof Lemma 4.14 such that

$$\|y^{\frac{\alpha}{2}}(\psi_{s,k} - i_{p_M} \psi_{s,k})'\|_{L^2(I_M)}^2 \leq c \lambda_k^{2s} h_\Omega^{2+2\varepsilon} Y^{2s}.$$

Employing Corollary 2.10 with  $r_1 = \frac{\alpha+1}{2} + 1 = 2 - s$  together with the monotonicity of  $e^{-\sqrt{\lambda_k} y}$  yields for  $Y \geq 1$

$$Y^{\frac{\alpha+1}{2}+1} \psi_{s,k}(Y) \leq c \lambda_k^{\frac{s-2}{2}} e^{-\frac{\sqrt{\lambda_k}}{2} Y} \leq c \lambda_k^s e^{-\frac{\sqrt{\lambda_1}}{2} Y},$$

where we used once again that the sequence  $(\lambda_k)_{k \in \mathbb{N}}$  is nondecreasing. Due to the previous results, we arrive at

$$(4.12) \quad \|y^{\frac{\alpha}{2}}(\psi_{s,k} - \tilde{i}_{p_M} \psi_{s,k})'\|_{L^2(I_M)}^2 \leq c \lambda_k^{2s} \left( h_\Omega^{2+2\varepsilon} Y^{2s} + p_M^4 Y^{-4} e^{-\sqrt{\lambda_1} Y} \right).$$

By combining (4.4), (4.11), and (4.12), we obtain as in the proof of Lemma 4.7

$$\begin{aligned} & \|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_M)}^2 \leq c \left( h_\Omega^{2+2\varepsilon} Y^{2s} + \{1 + p_M^4 Y^{-4}\} e^{-\sqrt{\lambda_1} Y} \right) \sum_{k=1}^{\infty} \lambda_k^{2s} \mathbf{u}_k^2 \\ & = c \left( h_\Omega^{2+2\varepsilon} Y^{2s} + \{1 + p_M^4 Y^{-4}\} e^{-\sqrt{\lambda_1} Y} \right) \|\mathbf{f}\|_{L^2(\Omega)}^2. \end{aligned}$$

According to Lemma 3.3, there holds that  $p_M \leq cM$ . Since

$$Y \geq \frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}} \quad \text{and} \quad M \leq \frac{2(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|}$$

by assumption, we obtain

$$\{1 + p_M^4 Y^{-4}\} e^{-\sqrt{\lambda_1} Y} \leq h_\Omega^2 + ch_\Omega^2 M^4 |\ln h_\Omega|^{-4} \leq ch_\Omega^2,$$

which ends the proof. □

**COROLLARY 4.16.** For  $\mathbf{f} \in L^2(\Omega)$ , let  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solution of (2.2). Moreover, let  $p \in \mathbb{N}^M$  be a linear degree vector as in (3.3) with some  $\beta > 0$  and let

$$\begin{aligned} 2 \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|} &\geq M \geq \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|}, \\ 2 \max\left(\frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right) &\geq Y \geq \max\left(\frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right) \end{aligned}$$

for some  $\varepsilon > 0$ , where  $b > 0$  is a constant depending on  $\sigma$  only. Then, it holds that

$$\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(C_Y)} \leq ch_\Omega \|\mathbf{f}\|_{L^2(\Omega)}.$$

*Proof.* By Lemmas 4.12, 4.14, and 4.15, we obtain

$$\begin{aligned} &\|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(C_Y)}^2 \\ &= \sum_{m=1}^M \|y^{\frac{\alpha}{2}} \nabla(u - i_y^p u)\|_{L^2(\Omega \times I_m)}^2 \leq c(h_\Omega^{2+2\varepsilon} Y^{2s} M + h_\Omega^2) \|\mathbf{f}\|_{L^2(\Omega)}^2 \\ &\leq c(h_\Omega^{2+2\varepsilon} |\ln h_\Omega|^{2s+1} + h_\Omega^2) \|\mathbf{f}\|_{L^2(\Omega)}^2 \leq ch_\Omega^2 \|\mathbf{f}\|_{L^2(\Omega)}^2, \end{aligned}$$

where we have used the bounds on  $M$  and  $Y$  and the boundedness of  $h_\Omega^{2\varepsilon} |\ln h_\Omega|^{2s+1}$  for  $\varepsilon > 0$ . □

Now, we are able to state the main result for this subsection analyzing the  $hp$ -FEM on geometric meshes.

**THEOREM 4.17.** For  $\mathbf{f} \in \mathbb{H}^{1-s}(\Omega)$ , let  $\mathbf{u} \in \mathbb{H}^s(\Omega)$  and  $u \in \mathring{H}_L^1(C, y^\alpha)$  be the solutions of (1.1) and (2.2), respectively, and let  $u_h \in V_{h,M}$  be the solution of (3.1). Moreover, let  $p \in \mathbb{N}^M$  be a linear degree vector as in (3.3) with some  $\beta > 0$  and let

$$(4.13) \quad \begin{aligned} 2 \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|} &\geq M \geq \frac{(1 + \varepsilon)|\ln h_\Omega|}{\min(s, \beta b)|\ln \sigma|}, \\ 2 \max\left(\frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right) &\geq Y \geq \max\left(\frac{2|\ln h_\Omega|}{\sqrt{\lambda_1}}, 1\right) \end{aligned}$$

for some  $\varepsilon > 0$ , where  $b > 0$  is a constant depending on  $\sigma$  only. Then, it holds that

$$\|\mathbf{u} - \text{tr}_\Omega u_h\|_{\mathbb{H}^s(\Omega)} \leq c \|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \leq ch_\Omega \|\mathbf{f}\|_{\mathbb{H}^{1-s}(\Omega)}.$$

*Proof.* The first inequality of the assertion is due to Propositions 2.4 and 2.5. Using Lemmas 4.1 and 4.2, we get

$$\begin{aligned} &\|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \\ &\leq \|y^{\frac{\alpha}{2}} \nabla(u - \pi_x u)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla(u - i_y u)\|_{L^2(C_Y)} + \|y^{\frac{\alpha}{2}} \nabla u\|_{L^2(C \setminus C_Y)}. \end{aligned}$$

The three terms on the right-hand side are estimated in Lemma 4.3, Corollary 4.16, and Proposition 2.11. Hence, we get

$$\|y^{\frac{\alpha}{2}} \nabla(u - u_h)\|_{L^2(C)} \leq ch_{\Omega} \|f\|_{\mathbb{H}^{1-s}(\Omega)} + ch_{\Omega} \|f\|_{L^2(\Omega)} + ce^{-\frac{\sqrt{\lambda_1}}{2} Y} \|f\|_{\mathbb{H}^{-s}(\Omega)}.$$

Then, the lower bound on  $Y$  yields  $e^{-\frac{\sqrt{\lambda_1}}{2} Y} \leq ch_{\Omega}$ , which implies the assertion.  $\square$

**THEOREM 4.18.** *The total number degrees of freedom  $\mathcal{N}_{\Omega, Y}$  in  $V_{h, M}$  to achieve the order of convergence given in Theorem 4.17 behaves like*

$$\mathcal{N}_{\Omega, Y} = O(\mathcal{N}_{\Omega}(\ln \mathcal{N}_{\Omega})^2).$$

*Proof.* As a direct consequence of Lemma 3.4, we obtain that the number of degrees of freedom  $\mathcal{N}_{\Omega, Y}$  of the discretization considered in this section fulfills  $\mathcal{N}_{\Omega, Y} = \mathcal{N}_{\Omega} \mathcal{N}_Y = O(\mathcal{N}_{\Omega} M^2)$ . Then, the assertion follows from  $M = O(|\ln h_{\Omega}|) = O(\ln \mathcal{N}_{\Omega})$ .  $\square$

*Remark 4.19.* We note that the factor 2, which appears in each of the inequalities of (4.13) as an upper bound, is chosen arbitrarily in the sense that any factor greater than one (independent of  $h_{\Omega}$ ) is sufficient for the validity of the error estimates and for the validity of the result about the complexity. Thus, once one has a guess or an approximation for  $\lambda_1$  and  $b$ , it is straightforward to choose  $Y$  and  $M$ .

### 5. Numerical experiments.

**5.1. Implementation.** For the discretization with respect to  $y$  in  $h$ -FEM and  $hp$ -FEM, we use hierarchical Lobatto polynomials (see, e.g., [34]) as local shape functions on  $\hat{I} = (0, 1)$ , which then are transformed onto each interval  $I_m$ . Without the weight, i.e., for  $s = \frac{1}{2}$ , this would result in a very sparse structure of the local stiffness matrix, since those shape functions are orthogonal. For  $s \neq \frac{1}{2}$  the latter does not hold; nevertheless the global matrix is of course still sparse.

Let  $\eta_i$  for  $i = 1, 2, \dots, \mathcal{N}_{\Omega}$  be the ansatz functions for the discretization of  $\Omega$  and  $\tau_j$  for  $j = 1, 2, \dots, \mathcal{N}_Y$  ansatz functions for the discretization of  $(0, Y)$ . On the cylinder  $C_Y$  we then use ansatz functions of the form  $\varphi_{i,j}(x, y) = \eta_i(x)\tau_j(y)$  with  $i = 1, 2, \dots, \mathcal{N}_{\Omega}$  and  $j = 1, \dots, \mathcal{N}_Y$ .

Due to this special structure, the system matrix  $S \in \mathbb{R}^{\mathcal{N}_{\Omega, Y} \times \mathcal{N}_{\Omega, Y}}$  for solving (3.1) can be expressed by means of the Kronecker product as

$$S = B^{\text{mass}} \otimes A^{\text{stiff}} + B^{\text{stiff}} \otimes A^{\text{mass}}.$$

Here,  $A \in \mathbb{R}^{\mathcal{N}_{\Omega} \times \mathcal{N}_{\Omega}}$  denotes matrices arising from discretization of  $\Omega$  and  $B \in \mathbb{R}^{\mathcal{N}_Y \times \mathcal{N}_Y}$  denotes matrices arising from discretization of  $(0, Y)$  given as

$$\begin{aligned} A_{ik}^{\text{mass}} &= \int_{\Omega} \eta_i(x)\eta_k(x) dx, & A_{ik}^{\text{stiff}} &= \int_{\Omega} \nabla \eta_i(x)\nabla \eta_k(x) dx, \quad i, k = 1, 2, \dots, \mathcal{N}_{\Omega}, \\ B_{jl}^{\text{mass}} &= \int_0^Y y^{\alpha} \tau_j(y)\tau_l(y) dy, & B_{jl}^{\text{stiff}} &= \int_0^Y y^{\alpha} \tau_j'(y)\tau_l'(y) dy, \quad j, l = 1, 2, \dots, \mathcal{N}_Y. \end{aligned}$$

We observe that one can assemble the matrices  $A$  and  $B$  completely independent from each other. This is advantageous since the weight  $y^{\alpha}$  only affects the  $B$  matrices, while the  $A$  matrices are standard FEM matrices, which can be computed by any FEM software.

Using the special structure of  $S$ , one can implement memory efficient solvers for the algebraic systems without ever fully assembling  $S$ . Indeed, the linear system

of equations corresponding to (3.1) can be reformulated as a generalized Sylvester matrix equation. For such equations efficient solvers are known. For instance, by using a simultaneous diagonalization of  $B^{\text{mass}}$  and  $B^{\text{stiff}}$ , which can be realized by solving a generalized eigenvalue problem of dimension  $O(\ln \mathcal{N}_\Omega)$ , the algebraic system can be decoupled. The arising  $O(\ln \mathcal{N}_\Omega)$  linear systems of equations of dimension  $\mathcal{N}_\Omega$  can be solved in parallel by standard solvers. This will be the topic of a forthcoming paper.

**5.2. Numerical results.** We take the following configuration from [31, section 6.1]. For  $\Omega = (0, 1)^2 \subset \mathbb{R}^2$  the eigenfunctions of the Dirichlet-Laplacian are known to be  $\varphi_{k,l}(x) = \sin(k\pi x_1) \sin(l\pi x_2)$  with corresponding eigenvalues  $\lambda_{k,l} = \pi^2(k^2 + l^2)$  for  $k, l = 1, 2, \dots$ . For the right-hand side  $f(x) = \lambda_{1,1}^s \varphi_{1,1}(x) = (2\pi^2)^s \sin(\pi x_1) \sin(\pi x_2)$ , the solutions  $\mathbf{u}$  of (1.1) and  $u$  of (2.2) are then given by

$$\mathbf{u}(x) = \sin(\pi x_1) \sin(\pi x_2) \quad \text{and} \quad u(x, y) = \frac{2^{1-\frac{s}{2}} \pi^s}{\Gamma(s)} \sin(\pi x_1) \sin(\pi x_2) y^s K_s(\sqrt{2}\pi y).$$

For the discretization by means of  $h$ -FEM (cf. sections 3.1 and 4.1), we choose the parameters

$$\mu = 0.8s, \quad M = \lceil h_\Omega^{-1} \rceil, \quad \text{and} \quad Y = \max\left(\frac{3|\ln h_\Omega|}{\sqrt{2}\pi}, 1\right),$$

whereas for the discretization by means of  $hp$ -FEM (cf. sections 3.2 and 4.2), we choose the following parameters:

$$\beta = 0.7, \quad \sigma = 0.125, \quad M = \left\lceil \frac{1.75|\ln h_\Omega|}{s|\ln \sigma|} \right\rceil, \quad \text{and} \quad Y = \max\left(\frac{3|\ln h_\Omega|}{\sqrt{2}\pi}, 1\right).$$

The orders of convergence stated by Theorems 4.9 and 4.17 in terms of  $h_\Omega$  are confirmed by the results of the numerical computations given in Figure 1. Note that the error  $\|y^{\frac{s}{2}} \nabla(u - u_h)\|_{L^2(C)}$  is evaluated by means of the identity

$$\|y^{\frac{s}{2}} \nabla(u - u_h)\|_{L^2(C)}^2 = d_s \int_\Omega (f \operatorname{tr}_\Omega u - f \operatorname{tr}_\Omega u_h) \, dx,$$

which holds due to the Galerkin orthogonality (4.1).

In Figure 2, we depict the errors for both types of discretizations over the total numbers of degrees of freedom  $\mathcal{N}_{\Omega,Y}$ . Thereby, the slower growth of  $\mathcal{N}_{\Omega,Y}$  for the  $hp$ -discretization given by Theorem 4.18 in comparison to Theorem 4.10 clearly leads to a drastic reduction of the number of degrees of freedom compared to  $h$ -FEM on a graded mesh. For instance, the number of degrees of freedom to achieve an error of less than  $9 \cdot 10^{-3}$  in the case  $s = 0.8$  reduces from  $\mathcal{N}_{\Omega,Y} = 1,072,692,225$  for  $h$ -FEM to  $\mathcal{N}_{\Omega,Y} = 9,661,477$  for  $hp$ -FEM, which is a factor of about 111.

**Appendix A. Estimates for  $\psi_s$  and its derivatives.** We begin with a representation of the derivatives of the expression  $z^s K_s(z)$ , where  $K_s$  are the modified Bessel functions of second kind. It will be used in what follows to derive estimates for the derivatives of  $\psi_s$ .

LEMMA A.1. *The derivatives of  $z^s K_s(z)$  of order  $n \in \mathbb{N}_0$  can be calculated as*

$$(A.1) \quad (z^s K_s(z))^{(n)} = \sum_{m=0}^n a_m^n z^{s-m} K_{s-(n-m)}(z),$$




FIG. 1.  $\|y^{\frac{s}{2}} \nabla(u - u_h)\|_{L^2(C)}$  for  $s = 0.2$  (left) and  $s = 0.8$  (right) over  $h_\Omega$ .




FIG. 2.  $\|y^{\frac{s}{2}} \nabla(u - u_h)\|_{L^2(C)}$  for  $s = 0.2$  (left) and  $s = 0.8$  (right) over the total number of degrees of freedom  $N_{\Omega,Y}$ .

where the coefficients  $a_m^n$  are given by

$$(A.2) \quad a_0^n = (-1)^n,$$

$$(A.3) \quad a_m^n = (-1)^{n+m} \frac{1}{2^m} \frac{n!}{m!(n-2m)!} \quad \text{for } 1 \leq m \leq \lfloor \frac{n}{2} \rfloor,$$

$$(A.4) \quad a_m^n = 0 \quad \text{for } \lfloor \frac{n}{2} \rfloor < m \leq n.$$

*Proof.* We prove this assertion by induction. To this end, we first collect some basic results for the modified Bessel function of second kind. In [1, eq. 9.6.28], we find for all  $\nu \in \mathbb{R}$

$$\frac{1}{z} \frac{d}{dz} (z^\nu K_\nu(z)) = -z^{\nu-1} K_{\nu-1}(z).$$

As a consequence, there holds

$$(A.5) \quad (z^\nu K_\nu(z))' = -z^\nu K_{\nu-1}(z).$$



Using the latter result, we get the following formula for  $m, l \in \mathbb{N}_0$ :

$$\begin{aligned} (z^{s-m}K_{s-l}(z))' &= (z^{l-m}z^{s-l}K_{s-l}(z))' \\ &= -z^{l-m}z^{s-l}K_{s-l-1}(z) + (l-m)z^{l-m-1}z^{s-l}K_{s-l}(z) \\ &= -z^{s-m}K_{s-(l+1)}(z) + (l-m)z^{s-m-1}K_{s-l}(z). \end{aligned}$$

By means of this, we obtain for  $m, n \in \mathbb{N}_0$  with  $m \leq n$  by setting  $l = n - m \geq 0$

$$(A.6) \quad (z^{s-m}K_{s-(n-m)}(z))' = -z^{s-m}K_{s-(n+1-m)}(z) + (n-2m)z^{s-m-1}K_{s-(n-m)}(z).$$

These elementary results build the basis for the induction: The hypothesis (A.1) clearly holds for  $n = 0$ .

Assuming that (A.1) holds for some  $n \in \mathbb{N}_0$ , we deduce

$$(z^sK_s(z))^{(n+1)} = \sum_{m=0}^n a_m^n (z^{s-m}K_{s-(n-m)}(z))'.$$

Employing (A.6), we continue with

$$\begin{aligned} (z^sK_s(z))^{(n+1)} &= \sum_{m=0}^n a_m^n (-z^{s-m}K_{s-(n+1-m)}(z) + (n-2m)z^{s-m-1}K_{s-(n-m)}(z)) \\ &= \sum_{m=0}^n -a_m^n z^{s-m}K_{s-(n+1-m)}(z) \\ &\quad + \sum_{m=1}^{n+1} a_{m-1}^n (n-2(m-1))z^{s-m}K_{s-(n+1-m)}(z) \\ &= -a_0^n z^s K_{s-(n+1)}(z) - na_n^n z^{s-n-1}K_s(z) \\ &\quad + \sum_{m=1}^n (-a_m^n + a_{m-1}^n (n-2(m-1)))z^{s-m}K_{s-(n+1-m)}(z). \end{aligned}$$

It remains to show that

$$a_0^{n+1} = -a_0^n, \quad a_m^{n+1} = -a_m^n + a_{m-1}^n (n-2(m-1)) \text{ for } 1 \leq m \leq n, \quad a_{n+1}^{n+1} = -na_n^n.$$

The first and third equations are obviously due to (A.2) and (A.4). Thus, we only elaborate on the second. We distinguish three cases for  $1 \leq m \leq n$ :

- $m \geq \lfloor \frac{n+1}{2} \rfloor + 2$ : Again due to (A.4), we have  $a_m^n = a_{m-1}^n = 0$ , since  $m, m-1 > \lfloor \frac{n}{2} \rfloor$ . Hence, it holds that

$$-a_m^n + a_{m-1}^n (n-2(m-1)) = 0 = a_m^{n+1}.$$

- $m = \lfloor \frac{n+1}{2} \rfloor + 1$ : Here, it holds that  $m > \lfloor \frac{n}{2} \rfloor$  and we already know that  $a_m^n = 0$ . Moreover, in the case that  $n$  is even, we deduce  $m = \frac{n}{2} + 1$  and

$$n - 2(m - 1) = n - 2\left(\frac{n}{2} + 1 - 1\right) = 0.$$

If  $n$  is odd, we obtain  $a_{m-1}^n = 0$  since  $m - 1 = \frac{n+1}{2} > \lfloor \frac{n}{2} \rfloor$ . As a consequence, we get

$$-a_m^n + a_{m-1}^n (n - 2(m - 1)) = 0 = a_m^{n+1}.$$

- $m \leq \lfloor \frac{n+1}{2} \rfloor$ : Here, we again distinguish between  $n$  even and  $n$  odd. In the first case, we have  $\lfloor \frac{n+1}{2} \rfloor = \lfloor \frac{n}{2} \rfloor$ . Hence, it holds that  $m - 1, m \leq \lfloor \frac{n}{2} \rfloor$  and by means of (A.3), we get

$$\begin{aligned} & -a_m^n + (n - 2(m - 1))a_{m-1}^n \\ &= (-1)^{n+m+1} \frac{1}{2^m} \frac{n!}{m!(n - 2m)!} \\ & \quad + (n - 2(m - 1))(-1)^{n+m-1} \frac{1}{2^{m-1}} \frac{n!}{(m - 1)!(n - 2(m - 1))!} \\ &= (-1)^{n+1+m} \frac{1}{2^m} \frac{(n + 1)!}{m!(n + 1 - 2m)!} = a_m^{n+1}. \end{aligned}$$

In the case that  $n$  is odd, we have that  $\lfloor \frac{n+1}{2} \rfloor = \lfloor \frac{n}{2} \rfloor + 1$ . Thus, for  $m \leq \lfloor \frac{n+1}{2} \rfloor - 1$ , we can reuse the calculations from before. If  $m = \lfloor \frac{n+1}{2} \rfloor = \frac{n+1}{2}$ , we have  $a_m^n = 0$  such that

$$\begin{aligned} -a_m^n + (n - 2(m - 1))a_{m-1}^n &= a_{m-1}^n = (-1)^{n+\frac{n-1}{2}} \frac{1}{2^{\frac{n-1}{2}}} \frac{n!}{(\frac{n-1}{2})!} \\ &= (-1)^{n+1+\frac{n+1}{2}} \frac{1}{2^{\frac{n+1}{2}}} \frac{(n + 1)!}{(\frac{n+1}{2})!} = a_m^{n+1}. \end{aligned}$$

This ends the proof. □

We next analyze  $\psi_s$  defined in (2.3) and its derivatives with respect to its boundedness properties.

LEMMA A.2. *For  $z \geq 0$ , it holds*

$$0 < \psi_s(z) \leq \psi_s(0) = 1.$$

*Proof.* Since  $z^\nu K_\nu(z) > 0$  for all  $z > 0$  and  $\nu > -1$  (see, e.g., [1, section 9.6]), and due to (A.5), the function  $\psi_s$  is positive and monotone decreasing such that  $\psi_s(z) \leq \psi_s(0)$  for all  $z \geq 0$ .

In [1, eq. 9.6.9] one can find for  $\nu > 0$  the following behavior of the modified Bessel function of the second kind for  $z \rightarrow 0$ :

$$K_\nu(z) \sim \frac{\Gamma(\nu)}{2^{1-\nu}} z^{-\nu}.$$

As a consequence, we obtain by (2.3)

$$(A.7) \quad \lim_{z \rightarrow 0} \psi_s(z) = \lim_{z \rightarrow 0} c_s z^s K_s(z) = 1,$$

which yields together with the foregoing observations the assertion. □

LEMMA A.3. *Let  $r \in [0, 1]$ . There exists a constant  $c > 0$  depending only on  $s$  such that for any  $z > 0$  and  $n \in \mathbb{N}$  it holds that*

$$|z^n \psi_s^{(n)}(z)| \leq c 8^n n! z^{2s-r}.$$

*Proof.* In order to deduce the bounds for the derivatives of  $\psi_s$ , we continue with collecting some auxiliary results. As before in the proof of Lemma A.2, we have that  $z^r z^\nu K_\nu(z)$  is positive for all  $z > 0$  and  $\nu > -1$ . Let  $\nu_0 = \min(\nu, \frac{1}{2})$ . From

[28, Theorem 5], we obtain for  $z > 0$  that  $z^{\nu_0} e^z K_\nu(z)$  is a decreasing function for all  $\nu \geq 0$ . To employ this, we consider the product

$$z^{\nu+r} K_\nu(z) = z^{\nu+r-\nu_0} e^{-z} \cdot z^{\nu_0} e^z K_\nu(z)$$

and note that

$$\arg \max_{z \geq 0} z^{\nu+r-\nu_0} e^{-z} = \nu + r - \nu_0 \quad \text{and} \quad (z^{\nu+r-\nu_0} e^{-z})' < 0 \quad \text{for} \quad z > \nu + r - \nu_0.$$

Hence,  $z^{\nu+r} K_\nu(z)$  admits its maximum in the interval  $[0, \nu+r-\nu_0]$ . Due to Lemma A.2 we consequently get by (2.3)

$$(A.8) \quad z^{\nu+r} K_\nu(z) = \frac{z^r}{c_\nu} \psi_\nu(z) \leq \frac{(\nu + r - \nu_0)^r}{c_\nu} \psi_\nu(0) = \frac{(\nu + r - \nu_0)^r}{c_\nu}.$$

Next, we are concerned with the bounds for the derivatives of  $\psi_s$ . Employing Lemma A.1 and the relation  $K_\nu(z) = K_{-\nu}(z)$  for  $\nu \in \mathbb{R}$  (see [1, eq. 9.6.6]), we obtain by (2.3)

$$\begin{aligned} |z^{n-2s+r} \psi_s^{(n)}(z)| &= \left| z^{n-2s+r} \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} c_s a_m^n z^{s-m} K_{s-(n-m)}(z) \right| \\ &\leq \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} c_s |a_m^n| |z^{n-m-s+r} K_{n-m-s}(z)|, \end{aligned}$$

where the coefficients are given by Lemma A.1. Let  $\nu(m, n) = n - m - s$ . We observe that  $\nu(m, n) > 0$  since  $m \leq \lfloor \frac{n}{2} \rfloor$ . Consequently, (A.8) yields

$$|z^{n-2s+r} \psi_s^{(n)}(z)| \leq \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \frac{c_s}{c_{\nu(m,n)}} |a_m^n| (\nu(m, n) + r - \nu_0(m, n))^r$$

with  $\nu_0(m, n) = \min(\nu(m, n), \frac{1}{2})$ . Since  $n - m \geq 1$ , it holds that

$$(A.9) \quad (\nu(m, n) + r - \nu_0(m, n))^r \leq (n - m + r)^r \leq n - m + 1.$$

Further, using

$$\Gamma(l + \rho) \leq \frac{\Gamma(l + 1)}{(l + \frac{\rho}{2})^{1-\rho}}$$

from [25, estimate (8)], which holds for all  $l \in \mathbb{N}_0$  and  $0 < \rho < 1$ , we get by choosing  $l = n - m - 1$  and  $\rho = 1 - s$

$$(A.10) \quad \begin{aligned} \frac{c_s}{c_{\nu(m,n)}} &= \frac{2^\nu \Gamma(\nu(m, n))}{2^s \Gamma(s)} = \frac{1}{4^s \Gamma(s)} 2^{n-m} \Gamma(\nu(m, n)) \leq \frac{1}{4^s \Gamma(s)} \frac{2^{n-m} \Gamma(n - m)}{(n - m - 1 + \frac{1-s}{2})^s} \\ &\leq \frac{1}{2^s \Gamma(s) (1 - s)^s} 2^{n-m} (n - m - 1)! = c 2^{n-m} (n - m - 1)! \end{aligned}$$

with a constant  $c$  depending only on  $s$ . Using (A.9) and (A.10), we get

$$(A.11) \quad |z^{n-2s+r} \psi_s^{(n)}(z)| \leq c \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} 2^{n-m} (n - m + 1)! |a_m^n|.$$

Estimating each summand separately yields by means of Lemma A.1

$$\begin{aligned}
 2^{n-m}(n-m+1)!|a_m^n| &= 2^{n-m}(n-m+1)! \frac{1}{2^m} \frac{n!}{m!(n-2m)!} \\
 &= 2^{n-2m}(m+1)n! \frac{(n-m+1)!}{(m+1)!(n-m+1-(m+1))!} \\
 &= 2^{n-2m}(m+1)n! \binom{n-m+1}{m+1} \\
 &\leq 4^n n!.
 \end{aligned}
 \tag{A.12}$$

For the last step, notice that

$$2^{-2m}(m+1) \leq 1 \quad \text{and} \quad \binom{n-m+1}{m+1} \leq 2^n.$$

Finally, (A.11) and (A.12) yield the assertion since  $\lfloor \frac{n}{2} \rfloor + 1 \leq 2^n$ . □

Finally, we state a result about the exponential decay of  $\psi_s$  and its derivative.

LEMMA A.4. *The following assertions hold:*

- (a) *Let  $s_0 = \min(s, \frac{1}{2})$ . Moreover, let  $z \geq a > 0$  and  $r \geq s_0 - s$ . Then there exists a constant  $c$  only depending on  $r$  and  $s$  such that*

$$|z^r \psi_s(z)| \leq ce^{-\frac{z}{2}} \cdot c_s a^{s_0} e^a K_s(a).$$

- (b) *Let  $s'_0 = \min(1-s, \frac{1}{2})$ . Moreover, let  $z \geq a > 0$  and  $r \geq s'_0 - s$ . Then there exists a constant  $c$  only depending on  $r$  and  $s$  such that*

$$|z^r \psi'_s(z)| \leq ce^{-\frac{z}{2}} \cdot c_s a^{s'_0} e^a K_{1-s}(a).$$

*Proof.* We start as in the proof of Lemma A.3. According to [28, Theorem 5], we get for  $z \geq 0$  that  $z^{s_0} e^z K_s(z)$  is a decreasing function. Consequently, having in mind the definition of  $\psi_s$  and that  $z^\nu K_\nu(z) > 0$  for all  $z \geq 0$  and  $\nu > -1$  (see the proof of Lemma A.2), we obtain

$$|z^r \psi_s(z)| = z^{r+s-s_0} e^{-z} \cdot c_s z^{s_0} e^z K_s(z) \leq z^{r+s-s_0} e^{-z} \cdot c_s a^{s_0} e^a K_s(a).$$

This is already the desired result for  $r = s_0 - s$  noticing that  $e^{-\frac{z}{2}} < 1 = c$  for  $z \geq 0$ . For  $r > s_0 - s$ , we observe that

$$z^{r+s-s_0} e^{-z} = z^{r+s-s_0} e^{-\frac{z}{2}} e^{-\frac{z}{2}} \leq ce^{-\frac{z}{2}},$$

where we used that

$$\max_{z \geq 0} (z^{r+s-s_0} e^{-\frac{z}{2}}) = \left( \frac{2(r+s-s_0)}{e} \right)^{r+s-s_0} = c.$$

Combining the previous results yields the first inequality of the assertion. Next, we deduce by means of the definition of  $\psi_s$  and (A.5)

$$z^r \psi'_s(z) = \frac{c_s}{c_{1-s}} z^{r+2s-1} c_{1-s} z^{1-s} K_{1-s}(z) = \frac{c_s}{c_{1-s}} z^{r+2s-1} \psi_{1-s}(z)$$

such that the second inequality of the assertion follows from the first one noting that  $r+2s-1 \geq s'_0 - s + 2s - 1 = \min(1-s, \frac{1}{2}) - (1-s)$  by the assumption on  $r$ . □

**Acknowledgments.** The authors acknowledge Christian Kahle's and Martin Stoll's support in the efficient solution of the arising linear systems.

## REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series 55, Superintendent of Documents, U.S. Government Printing Office, Washington, DC, 1964.
- [2] G. ACOSTA, F. M. BERSETCHE, AND J. P. BORTHAGARAY, *Finite Element Approximations for Fractional Evolution Problems*, arXiv:1705.09815, 2017.
- [3] G. ACOSTA, F. M. BERSETCHE, AND J. P. BORTHAGARAY, *A short FE implementation for a 2D homogeneous Dirichlet problem of a fractional Laplacian*, *Comput. Math. Appl.*, 74 (2017), pp. 784–816.
- [4] G. ACOSTA AND J. P. BORTHAGARAY, *A fractional Laplace equation: Regularity of solutions and finite element approximations*, *SIAM J. Numer. Anal.*, 55 (2017), pp. 472–495.
- [5] G. ACOSTA, J. P. BORTHAGARAY, O. BRUNO, AND M. MAAS, *Regularity theory and high order numerical methods for the (1D)-Fractional Laplacian*, *Math. Comp.*, 87 (2018), pp. 1821–1857.
- [6] T. APEL, A.-M. SÄNDIG, AND J. R. WHITEMAN, *Graded mesh refinement and error estimates for finite element solutions of elliptic boundary value problems in non-smooth domains*, *Math. Methods Appl. Sci.*, 19 (1996), pp. 63–85.
- [7] A. BONITO, W. LEI, AND J. E. PASCIAK, *The approximation of parabolic equations involving fractional powers of elliptic operators*, *J. Comput. Appl. Math.*, 315 (2017), pp. 32–48.
- [8] A. BONITO, W. LEI, AND J. E. PASCIAK, *Numerical Approximation of Space-Time Fractional Parabolic Equations*, arXiv:1704.04254, 2017.
- [9] A. BONITO AND J. E. PASCIAK, *Numerical approximation of fractional powers of elliptic operators*, *Math. Comp.*, 84 (2015), pp. 2083–2110.
- [10] A. BONITO AND J. E. PASCIAK, *Numerical approximation of fractional powers of regularly accretive operators*, *IMA J. Numer. Anal.*, 37 (2017), pp. 1245–1273.
- [11] J. P. BORTHAGARAY, L. M. DEL PEZZO, AND S. MARTÍNEZ, *Finite Element Approximation for the Fractional Eigenvalue Problem*, arXiv:1603.00317, 2016.
- [12] X. CABRÉ AND J. TAN, *Positive solutions of nonlinear problems involving the square root of the Laplacian*, *Adv. Math.*, 224 (2010), pp. 2052–2093.
- [13] L. CAFFARELLI AND L. SILVESTRE, *An extension problem related to the fractional Laplacian*, *Comm. Partial Differential Equations*, 32 (2007), pp. 1245–1260.
- [14] A. CAPELLA, J. DÁVILA, L. DUPAIGNE, AND Y. SIRE, *Regularity of radial extremal solutions for some non-local semilinear equations*, *Comm. Partial Differential Equations*, 36 (2011), pp. 1353–1384.
- [15] L. CHEN, R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *A PDE approach to fractional diffusion: A posteriori error analysis*, *J. Comput. Phys.*, 293 (2015), pp. 339–358.
- [16] L. CHEN, R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *Multilevel methods for nonuniformly elliptic operators and fractional diffusion*, *Math. Comp.*, 85 (2016), pp. 2583–2607.
- [17] M. DAUGE, *Elliptic Boundary Value Problems on Corner Domains: Smoothness and Asymptotics of Solutions*, *Lecture Notes in Math.*, Springer-Verlag, Berlin, 1988.
- [18] M. D'ELIA AND M. GUNZBURGER, *The fractional Laplacian operator on bounded domains as a special case of the nonlocal diffusion operator*, *Comput. Math. Appl.*, 66 (2013), pp. 1245–1260.
- [19] L. FEJÉR, *Bestimmung derjenigen Abszissen eines Intervalles, für welche die Quadratsumme der Grundfunktionen der Lagrangeschen Interpolation im Intervalle ein Möglichst kleines Maximum Besitzt*, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2)*, 1 (1932), pp. 263–276.
- [20] Q. GUAN AND M. GUNZBURGER,  *$\theta$  schemes for finite element discretization of the space-time fractional diffusion equations*, *J. Comput. Appl. Math.*, 288 (2015), pp. 264–273.
- [21] Q. GUAN AND M. GUNZBURGER, *Analysis and approximation of a nonlocal obstacle problem*, *J. Comput. Appl. Math.*, 313 (2017), pp. 102–118.
- [22] Y. HUANG AND A. OBERMAN, *Numerical methods for the fractional Laplacian: A finite difference–quadrature approach*, *SIAM J. Numer. Anal.*, 52 (2014), pp. 3056–3084.
- [23] M. ILIĆ, F. LIU, I. TURNER, AND V. ANH, *Numerical approximation of a fractional-in-space diffusion equation. I*, *Fract. Calc. Appl. Anal.*, 8 (2005), pp. 323–341.
- [24] M. ILIĆ, F. LIU, I. TURNER, AND V. ANH, *Numerical approximation of a fractional-in-space diffusion equation. II. With nonhomogeneous boundary conditions*, *Fract. Calc. Appl. Anal.*, 9 (2006), pp. 333–349.

- [25] L. LORCH, *Inequalities for ultraspherical polynomials and the gamma function*, J. Approx. Theory, 40 (1984), pp. 115–120.
- [26] V. MAZ'YA AND J. ROSSMANN, *Elliptic Equations in Polyhedral Domains*, Math. Surveys Monogr., AMS, Providence, RI, 2010.
- [27] J. M. MELENK, *hp-Finite Element Methods for Singular Perturbations*, Lecture Notes in Math., Springer-Verlag, Berlin, 2002.
- [28] K. S. MILLER AND S. G. SAMKO, *Completely monotonic functions*, Integral Transform. Spec. Funct., 12 (2001), pp. 389–402.
- [29] S. A. MOLČANOV AND E. OSTROVSKIĬ, *Symmetric stable processes as traces of degenerate diffusion processes*, Teor. Verojatnost. i Primenen., 14 (1969), pp. 127–130.
- [30] R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *Convergence rates for the classical, thin and fractional elliptic obstacle problems*, Philos. Trans. Roy. Soc. A, 373 (2015), pp. 20140449, 14.
- [31] R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *A PDE approach to fractional diffusion in general domains: A priori error analysis*, Found. Comput. Math., 15 (2015), pp. 733–791.
- [32] R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *A PDE approach to space-time fractional parabolic problems*, SIAM J. Numer. Anal., 54 (2016), pp. 848–873.
- [33] L. A. OGANESJAN AND L. A. RUHOVEC, *Variatsionno-raznostnye metody resheniya ellipticheskikh uravnenii*, Akad. Nauk Armyan. SSR, Erevan, 1979.
- [34] P. ŠOLÍN, K. SEGETH, AND I. DOLEŽEL, *Higher-Order Finite Element Methods*, Stud. Adv. Math., Chapman & Hall/CRC, Boca Raton, FL 2004.
- [35] P. R. STINGA AND J. L. TORREA, *Extension problem and Harnack's inequality for some fractional operators*, Comm. Partial Differential Equations, 35 (2010), pp. 2092–2122.
- [36] Q. YANG, I. TURNER, F. LIU, AND M. ILIĆ, *Novel numerical methods for solving the time-space fractional diffusion equation in two dimensions*, SIAM J. Sci. Comput., 33 (2011), pp. 1159–1180.